Parma International School of Theoretical Physics

September 3 - September 8, 2012

Parma, Italy

Scattering Amplitudes in QCD, Supersymmetric Gauge Theories and Supergravity

One-loop calculations in QCD and N=4 supersymmetric gauge theories

Zoltan Kunszt, ETH, Zurich

Reviews:

R. Keith Ellis, Zoltan Kunszt, Kirilll Melnikov, Giulia Zanderighi,
One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts.
Phys.Rep. (in print), arXiv:1105.4319 [hep-ph]

Harald Ita, **Susy Theories in QCD: Numerical Approaches, J.Phys.A A44 (2011) 454005**

Books:

R. Keith Ellis, James W. Stirling, Bryan Webber, **QCD and Collider Physics,** Cambridge University Press, 2003

Taizo Muta, **Foundation of Quantum Chromodynamics,** World Scientific, 1997

M. Creutz, **Lattice**

Papers:

Z. Bern, L. Dixon, D. Dunbar and D. Kosower,
Fusing gauge theory tree amplitudes into loop amplitudes, **Nucl.Phys. B435 (1995) 59-101**

One-loop calculations in QCD and N=4 supersymmetric gauge theories

OUTLINE

Lecture 1: QCD basics

Lecture 2: One loop tensor integrals and their reduction

Lecture 3: Unitarity method and amplitudes

Lecture 4: Analytic and numerical computations

Lecture 5: Outlook

QCD basics

QCD basics

SU(3) vector Yang-Mills theory coupled to quarks:

QCD basics

SU(3) vector Yang-Mills theory coupled to quarks:

$$\mathcal{L} = -\frac{1}{4}\mathbf{G}^{\mathbf{a}}_{\mu\nu}\mathbf{G}^{\mu\nu}_{\mathbf{a}} + \sum_{\mathbf{i,j,f}} \bar{\mathbf{q}}^{(\mathbf{f})}_{\mathbf{i}}(\mathbf{i}\hat{\mathbf{D}} - \mathbf{m})_{\mathbf{ij}}\mathbf{q}^{(\mathbf{f})}_{\mathbf{j}} - \frac{\mathbf{1}}{\mathbf{2}\lambda}(\partial^{\mu}\mathbf{G}^{\mathbf{a}}_{\mu})^{\mathbf{2}} + \partial_{\mu}\mathbf{c}^{\mathbf{a}\,\dagger}\mathbf{D}^{\mu}_{\mathbf{abj}}\mathbf{c}^{\mathbf{b}} \ ,$$

$$(\mathbf{D}_{\mu})_{\mathbf{ij}} = \partial_{\mu}\delta_{\mathbf{ij}} - \mathbf{igT}^{\mathbf{a}}_{\mathbf{ij}}\mathbf{G}^{\mathbf{a}}_{\mu}, \quad \mathbf{i,j} = \mathbf{1,2,3}\,; \quad \mathbf{a} = \mathbf{1},\ldots,\mathbf{8}$$

$$(\mathbf{D}_{\mu})_{\mathbf{ab}} = \partial_{\mu}\delta_{\mathbf{ab}} - \mathbf{igF}^{\mathbf{c}}_{\mathbf{ab}}\mathbf{G}^{\mathbf{c}}_{\mu}, \quad \mathbf{a,b,c} = \mathbf{1},\ldots,\mathbf{8}$$

$$\mathbf{G}^{\mathbf{a}}_{\mu\nu} = \partial_{\mu}\mathbf{G}^{\mathbf{a}}_{\nu} - \partial_{\nu}\mathbf{G}^{\mathbf{a}}_{\mu} + \mathbf{gf}^{\mathbf{abc}}\mathbf{G}^{\mathbf{b}}_{\mu}\mathbf{G}^{\mathbf{c}}_{\nu} \ ,$$

QCD basics



SU(3) vector Yang-Mills theory coupled to quarks:

$$\mathcal{L} = -\frac{1}{4}\mathbf{G}^{\mathbf{a}}_{\mu\nu}\mathbf{G}^{\mu\nu}_{\mathbf{a}} + \sum_{\mathbf{i,j,f}} \bar{\mathbf{q}}^{(\mathbf{f})}_{\mathbf{i}}(\mathbf{i}\hat{\mathbf{D}} - \mathbf{m})_{\mathbf{ij}}\mathbf{q}^{(\mathbf{f})}_{\mathbf{j}} - \frac{\mathbf{1}}{\mathbf{2}\lambda}(\partial^{\mu}\mathbf{G}^{\mathbf{a}}_{\mu})^{\mathbf{2}} + \partial_{\mu}\mathbf{c}^{\mathbf{a}\,\dagger}\mathbf{D}^{\mu}_{\mathbf{abj}}\mathbf{c}^{\mathbf{b}} \ ,$$

$$(\mathbf{D}_{\mu})_{\mathbf{ij}} = \partial_{\mu}\delta_{\mathbf{ij}} - \mathbf{igT}^{\mathbf{a}}_{\mathbf{ij}}\mathbf{G}^{\mathbf{a}}_{\mu}, \quad \mathbf{i,j} = \mathbf{1,2,3}\,; \quad \mathbf{a} = \mathbf{1},\ldots,\mathbf{8}$$

$$(\mathbf{D}_{\mu})_{\mathbf{ab}} = \partial_{\mu}\delta_{\mathbf{ab}} - \mathbf{igF}^{\mathbf{c}}_{\mathbf{ab}}\mathbf{G}^{\mathbf{c}}_{\mu}, \quad \mathbf{a,b,c} = \mathbf{1},\ldots,\mathbf{8}$$

$$\mathbf{G}^{\mathbf{a}}_{\mu\nu} = \partial_{\mu}\mathbf{G}^{\mathbf{a}}_{\nu} - \partial_{\nu}\mathbf{G}^{\mathbf{a}}_{\mu} + \mathbf{gf}^{\mathbf{abc}}\mathbf{G}^{\mathbf{b}}_{\mu}\mathbf{G}^{\mathbf{c}}_{\nu} \ ,$$

QCD basics



SU(3) vector Yang-Mills theory coupled to quarks:

$$\mathcal{L} = -\frac{1}{4}\mathbf{G}^{\mathbf{a}}_{\mu\nu}\mathbf{G}^{\mu\nu}_{\mathbf{a}} + \sum_{\mathbf{i,j,f}} \bar{\mathbf{q}}^{(\mathbf{f})}_{\mathbf{i}}(\mathbf{i}\hat{\mathbf{D}} - \mathbf{m})_{\mathbf{ij}}\mathbf{q}^{(\mathbf{f})}_{\mathbf{j}} - \frac{\mathbf{1}}{\mathbf{2}\lambda}(\partial^{\mu}\mathbf{G}^{\mathbf{a}}_{\mu})^{\mathbf{2}} + \partial_{\mu}\mathbf{c}^{\mathbf{a}\,\dagger}\mathbf{D}^{\mu}_{\mathbf{abj}}\mathbf{c}^{\mathbf{b}} \ ,$$

$$(\mathbf{D}_{\mu})_{\mathbf{ij}} = \partial_{\mu}\delta_{\mathbf{ij}} - \mathbf{igT}^{\mathbf{a}}_{\mathbf{ij}}\mathbf{G}^{\mathbf{a}}_{\mu}, \quad \mathbf{i,j} = \mathbf{1,2,3}\,; \quad \mathbf{a} = \mathbf{1},\ldots,\mathbf{8}$$

$$(\mathbf{D}_{\mu})_{\mathbf{ab}} = \partial_{\mu}\delta_{\mathbf{ab}} - \mathbf{igF}^{\mathbf{c}}_{\mathbf{ab}}\mathbf{G}^{\mathbf{c}}_{\mu}, \quad \mathbf{a,b,c} = \mathbf{1},\ldots,\mathbf{8}$$

$$\mathbf{G}^{\mathbf{a}}_{\mu\nu} = \partial_{\mu}\mathbf{G}^{\mathbf{a}}_{\nu} - \partial_{\nu}\mathbf{G}^{\mathbf{a}}_{\mu} + \mathbf{gf}^{\mathbf{abc}}\mathbf{G}^{\mathbf{b}}_{\mu}\mathbf{G}^{\mathbf{c}}_{\nu} \ ,$$

Quarks come in six different flavors (u,d,s,c,b,t) and three colors (red, blue, green).

$$\hat{\mathbf{G}}^{\mathbf{\Omega}}_{\mu}(\mathbf{x}) = \mathbf{\Omega}(\mathbf{x})\hat{\mathbf{G}}_{\mu}\mathbf{\Omega}^{-\mathbf{1}}(\mathbf{x}) + \mathbf{\Omega}(\mathbf{x})\partial_{\mu}\mathbf{\Omega}(\mathbf{x})^{-\mathbf{1}} \ , \quad \hat{\mathbf{G}}_{\mu} = \mathbf{T}^{\mathbf{a}}\mathbf{G}^{\mathbf{a}}_{\mu}$$

$$\mathbf{q}(\mathbf{x})_{\mathbf{i}} = \left(\mathbf{e}^{\{\mathbf{i}\sum_{\mathbf{a=1}}^{\mathbf{8}}\mathbf{T}^{\mathbf{a}}\xi^{\mathbf{a}}(\mathbf{x})\}}\right)_{\mathbf{ij}}\mathbf{q_j}(\mathbf{x}) \ = \mathbf{\Omega}(\mathbf{x})_{\mathbf{ij}}\mathbf{q_j}(\mathbf{x})$$

$$\delta\mathbf{G}^{\mathbf{a}}_{\mu}(\mathbf{x}) = -\frac{\mathbf{1}}{\mathbf{g}}\partial_{\mu}\xi^{\mathbf{a}}(\mathbf{x}) - \sum_{\mathbf{b,c=1}}^{\mathbf{8}}\mathbf{f}^{\mathbf{abc}}\xi^{\mathbf{b}}(\mathbf{x})\mathbf{G}^{\mathbf{c}}_{\mu}(\mathbf{x})$$

} SU(3) gauge transformations

QCD is a consistent QFT with rich dynamical structure:

QCD is a consistent QFT with rich dynamical structure:

- **Asymptotic freedom:**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

QCD is a consistent QFT with rich dynamical structure:

- **Asymptotic freedom:**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **Confinement Hypothesis:**
  Spectrum of the theory is given by colorless states (hadrons).
  The color degrees of freedom are permanently confined.
  Hadrons are made from quarks and gluons, hold together by the color force mediated by colored gluons.

  1 000 000 $ Milleneum Problem, Clay Mathematics Institute, Cambridge, MA
  Existence of Yang-Mills Theories and proof of mass gap.

It is generally believed that the hypothesis is correct.
It is supported by large scale computer simulations of lattice QCD.

QCD is a consistent QFT with rich dynamical structure:

- **Asymptotic freedom:**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **Confinement Hypothesis:**
  Spectrum of the theory is given by colorless states (hadrons).
  The color degrees of freedom are permanently confined.
  Hadrons are made from quarks and gluons, hold together by the color force mediated by colored gluons.

  1 000 000 $ Milleneum Problem, Clay Mathematics Institute, Cambridge, MA
  Existence of Yang-Mills Theories and proof of mass gap.

  It is generally believed that the hypothesis is correct.
  It is supported by large scale computer simulations of lattice QCD.

- **Spontaneous breaking of approximate chiral symmetry:**
  u,d,(s) quarks are massless: accidental SU(2)_L X SU(2)_R global symmetry
  Pions are the Nambu-Goldstone bosons.

QCD is a consistent QFT with rich dynamical structure:

- **Asymptotic freedom:**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **Confinement Hypothesis:**
  Spectrum of the theory is given by colorless states (hadrons).
  The color degrees of freedom are permanently confined.
  Hadrons are made from quarks and gluons, hold together by the color force mediated by colored gluons.

  1 000 000 $ Milleneum Problem, Clay Mathematics Institute, Cambridge, MA
  Existence of Yang-Mills Theories and proof of mass gap.

  It is generally believed that the hypothesis is correct.
  It is supported by large scale computer simulations of lattice QCD.

- **Spontaneous breaking of approximate chiral symmetry:**
  u,d,(s) quarks are massless: accidental SU(2)_L X SU(2)_R global symmetry
  Pions are the Nambu-Goldstone bosons.

- **Non-trivial topological gauge configurations:**
  Instantons, $\mathbf{U(1)_A}$ anomaly, Theta term in Lagrangian: $-\Theta \frac{g^2}{64\pi^2} \epsilon^{\mu\nu\lambda\sigma} G^i_{\mu\nu} G^i_{\lambda\sigma}$

- ....

Weak coupling perturbation theory

Perturbative series of renormalizable field theory in terms of gluons an quarks with standard Feynman rules and Feynman diagrams.

Propagators:



$$\delta_{ab}\frac{-i}{k^2+i\epsilon}\left[g^{\mu\nu}+(\lambda-1)\frac{k^\mu k^\nu}{k^2+i\epsilon}\right]$$

$$\delta_{ij}\left(\frac{i}{\hat{k}-m+i\epsilon}\right)_{\alpha\beta}$$

$$\delta_{ab}\frac{i}{k^2+i\epsilon}$$

Vertices:



$$gf^{a_1a_2a_3}\Big[(k_1-k_2)^{\mu_3}g^{\mu_1\mu_2}+(k_2-k_3)^{\mu_1}g^{\mu_2\mu_3}+(k_3-k_1)^{\mu_2}g^{\mu_3\mu_1}\Big]$$



$$-ig^2f^{a_1a_3b}f^{a_2a_4b}\left(g^{\mu_1\mu_2}g^{\mu_3\mu_4}-g^{\mu_1\mu_4}g^{\mu_2\mu_3}\right)+(1\leftrightarrow 2)+(2\leftrightarrow 3)$$



$$ig(T^a)_{ij}(\gamma^\mu)_{\alpha\beta}$$



$$-gf^{a_1a_2a_3}k_3^\mu$$

Weak coupling perturbation theory:

The diagrammatic rules mathematically are not always well defined. In higher orders diagrams with closed loops can appear when the expressions given by the Feynman diagrams have to be integrated over the internal loop momenta. Loop integrals can become divergent both in the ultraviolet region (at large values of the loop momenta) and in the the infrared region (soft and collinear regions of the loop momenta).

Weak coupling dimensional regularization scheme:

Perturbative series of renormalizable field theory in terms of gluons and quarks with standard Feynman rules and Feynman diagrams.

Weak coupling dimensional regularization scheme:

Perturbative series of renormalizable field theory in terms of gluons and quarks with standard Feynman rules and Feynman diagrams.

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires the introduction of some convenient regularization scheme for the divergent integrals and some conventions concerning the renormalization.

In QCD dimensional regularization is particularly convenient since it is a gauge invariant and Lorentz invariant regularization scheme.

Weak coupling dimensional regularization scheme:

Perturbative series of renormalizable field theory in terms of gluons and quarks with standard Feynman rules and Feynman diagrams.

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires the introduction of some convenient regularization scheme for the divergent integrals and some conventions concerning the renormalization.

In QCD dimensional regularization is particularly convenient since it is a gauge invariant and Lorentz invariant regularization scheme.

Integrals which are divergent in four dimensions are carried out in different dimensions where the integrals are finite and the singularities of the integrals are exhibited as poles in ϵ with analytic continuation into d=4-2ϵ dimensions.

$$\int \frac{\mathbf{d}^{\mathbf{4-2\epsilon}}\mathbf{dk}}{(\mathbf{2\pi})^{\mathbf{4-2\epsilon}}} \frac{\mathbf{1}}{[\mathbf{-k^2+C-i\epsilon}]^{\mathbf{s}}} = \mathbf{i}(\mathbf{4\pi})^{\mathbf{-2+\epsilon}}[\mathbf{C-i\epsilon}]^{\mathbf{2-s-\epsilon}} \frac{\mathbf{\Gamma(s-2+\epsilon)}}{\mathbf{\Gamma(s)}}$$

Weak coupling renormalization with counter terms:

Weak coupling renormalization with counter terms:

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires regularization scheme for the divergent integrals and some conventions concerning the renormalization.

The counter terms are generated with the help of the renormalization factors Z_i

Weak coupling renormalization with counter terms:

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires regularization scheme for the divergent integrals and some conventions concerning the renormalization.

The counter terms are generated with the help of the renormalization factors $\mathbf{Z_i}$

$$\mathbf{G}_\mu = \mathbf{Z_3^{-1/2}} \mathbf{G}_\mu^{(0)}, \quad \mathbf{q} = \mathbf{Z_2^{-1/2}} \mathbf{q}^{(0)}, \quad \mathbf{c_a} = \tilde{\mathbf{Z}}_\mathbf{3}^{\mathbf{-1/2}} \mathbf{c}_\mathbf{a}^{(0)}, \quad \mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g}^{(0)}(\epsilon), \quad \mathbf{m} = \mathbf{Z_m^{-1}} \mathbf{m}^{(0)}$$

Weak coupling renormalization with counter terms:

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires regularization scheme for the divergent integrals and some conventions concerning the renormalization.

The counter terms are generated with the help of the renormalization factors $\mathbf{Z_i}$

$$\mathbf{G}_\mu = \mathbf{Z_3^{-1/2}} \mathbf{G}_\mu^{(0)}, \quad \mathbf{q} = \mathbf{Z_2^{-1/2}} \mathbf{q}^{(0)}, \quad \mathbf{c_a} = \tilde{\mathbf{Z}}_\mathbf{3}^{\mathbf{-1/2}} \mathbf{c}_\mathbf{a}^{(0)}, \quad \mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g}^{(0)}(\epsilon)\,, \quad \mathbf{m} = \mathbf{Z_m^{-1}} \mathbf{m}^{(0)}$$

The difference of the bare Lagrangian and the renormalized Lagrangian gives the counter terms:

$$\mathcal{L}_\mathbf{ct} = -(\mathbf{Z_3}-\mathbf{1}) \frac{\mathbf{1}}{\mathbf{4}} \left(\partial_\mu \mathbf{G}_\nu^\mathbf{i} - \partial_\nu \mathbf{G}_\mu^\mathbf{i}\right)^\mathbf{2} - (\mathbf{Z_1}-\mathbf{1}) \frac{\mathbf{g}}{\mathbf{2}} \mu^\epsilon \mathbf{f^{ijk}} (\partial_\mu \mathbf{G}_\nu^\mathbf{i} - \partial_\nu \mathbf{G}_\mu^\mathbf{i}) \mathbf{G}^{\mathbf{j},\mu} \mathbf{G}^{\mathbf{k},\nu} + \cdots$$

Weak coupling renormalization with counter terms:

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires regularization scheme for the divergent integrals and some conventions concerning the renormalization.

The counter terms are generated with the help of the renormalization factors $\mathbf{Z_i}$

$$\mathbf{G}_\mu = \mathbf{Z_3^{-1/2}} \mathbf{G}_\mu^{(0)}, \quad \mathbf{q} = \mathbf{Z_2^{-1/2}} \mathbf{q}^{(0)}, \quad \mathbf{c_a} = \tilde{\mathbf{Z}}_\mathbf{3}^{\mathbf{-1/2}} \mathbf{c}_\mathbf{a}^{(0)}, \quad \mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g}^{(0)}(\epsilon), \quad \mathbf{m} = \mathbf{Z_m^{-1}} \mathbf{m}^{(0)}$$

The difference of the bare Lagrangian and the renormalized Lagrangian gives the counter terms:

$$\mathcal{L}_{\mathbf{ct}} = -(\mathbf{Z_3}-\mathbf{1}) \frac{\mathbf{1}}{\mathbf{4}} \left(\partial_\mu \mathbf{G}_\nu^{\mathbf{i}} - \partial_\nu \mathbf{G}_\mu^{\mathbf{i}}\right)^{\mathbf{2}} - (\mathbf{Z_1}-\mathbf{1}) \frac{\mathbf{g}}{\mathbf{2}} \mu^\epsilon \mathbf{f^{ijk}} (\partial_\mu \mathbf{G}_\nu^{\mathbf{i}} - \partial_\nu \mathbf{G}_\mu^{\mathbf{i}}) \mathbf{G}^{\mathbf{j},\mu} \mathbf{G}^{\mathbf{k},\nu} + \cdots$$

$$\mathbf{Z_1} - \mathbf{1} = \frac{\mathbf{g^2}}{\mathbf{16\pi^2}} \frac{\mathbf{1}}{\epsilon} \left[\mathbf{N_C} \left(\frac{\mathbf{17}}{\mathbf{12}} - \frac{\mathbf{3}\lambda}{\mathbf{4}}\right) - \frac{\mathbf{4}}{\mathbf{3}} \mathbf{n_f T_R}\right], \qquad \mathbf{Z_3} - \mathbf{1} = \frac{\mathbf{g^2}}{\mathbf{16\pi^2}} \frac{\mathbf{1}}{\epsilon} \left[\mathbf{N_C} \left(\frac{\mathbf{13}}{\mathbf{6}} - \frac{\lambda}{\mathbf{2}}\right) - \frac{\mathbf{4}}{-} \mathbf{3 n_f T_R}\right], \cdots$$

Weak coupling renormalization with counter terms:

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires regularization scheme for the divergent integrals and some conventions concerning the renormalization.

The counter terms are generated with the help of the renormalization factors $\mathbf{Z_i}$

$$\mathbf{G}_\mu = \mathbf{Z_3^{-1/2}} \mathbf{G}_\mu^{(0)}, \quad \mathbf{q} = \mathbf{Z_2^{-1/2}} \mathbf{q}^{(0)}, \quad \mathbf{c_a} = \tilde{\mathbf{Z}}_\mathbf{3}^{\mathbf{-1/2}} \mathbf{c}_\mathbf{a}^{(0)}, \quad \mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g}^{(0)}(\epsilon), \quad \mathbf{m} = \mathbf{Z_m^{-1}} \mathbf{m}^{(0)}$$

The difference of the bare Lagrangian and the renormalized Lagrangian gives the counter terms:

$$\mathcal{L}_{\mathbf{ct}} = -(\mathbf{Z_3}-\mathbf{1}) \frac{\mathbf{1}}{\mathbf{4}} \left(\partial_\mu \mathbf{G}_\nu^{\mathbf{i}} - \partial_\nu \mathbf{G}_\mu^{\mathbf{i}}\right)^{\mathbf{2}} - (\mathbf{Z_1}-\mathbf{1}) \frac{\mathbf{g}}{\mathbf{2}} \mu^\epsilon \mathbf{f^{ijk}} (\partial_\mu \mathbf{G}_\nu^{\mathbf{i}} - \partial_\nu \mathbf{G}_\mu^{\mathbf{i}}) \mathbf{G}^{\mathbf{j},\mu} \mathbf{G}^{\mathbf{k},\nu} + \cdots$$

$$\mathbf{Z_1} - \mathbf{1} = \frac{\mathbf{g^2}}{\mathbf{16\pi^2}} \frac{\mathbf{1}}{\epsilon} \left[\mathbf{N_C} \left(\frac{\mathbf{17}}{\mathbf{12}} - \frac{\mathbf{3}\lambda}{\mathbf{4}}\right) - \frac{\mathbf{4}}{\mathbf{3}} \mathbf{n_f T_R}\right], \qquad \mathbf{Z_3} - \mathbf{1} = \frac{\mathbf{g^2}}{\mathbf{16\pi^2}} \frac{\mathbf{1}}{\epsilon} \left[\mathbf{N_C} \left(\frac{\mathbf{13}}{\mathbf{6}} - \frac{\lambda}{\mathbf{2}}\right) - \frac{\mathbf{4}}{\mathbf{3}} \mathbf{n_f T_R}\right], \cdots$$

Renormalization prescription: subtract the pole only (MS-scheme)

Weak coupling renormalization with counter terms:

Ultraviolet divergences can be cancelled by renormalizing the parameters of the theory. It requires regularization scheme for the divergent integrals and some conventions concerning the renormalization.

The counter terms are generated with the help of the renormalization factors $\mathbf{Z_i}$

$$\mathbf{G}_\mu = \mathbf{Z_3^{-1/2}} \mathbf{G}_\mu^{(0)}, \quad \mathbf{q} = \mathbf{Z_2^{-1/2}} \mathbf{q}^{(0)}, \quad \mathbf{c_a} = \tilde{\mathbf{Z}}_\mathbf{3}^{\mathbf{-1/2}} \mathbf{c}_\mathbf{a}^{(0)}, \quad \mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g}^{(0)}(\epsilon), \quad \mathbf{m} = \mathbf{Z_m^{-1}} \mathbf{m}^{(0)}$$

The difference of the bare Lagrangian and the renormalized Lagrangian gives the counter terms:

$$\mathcal{L}_{\mathbf{ct}} = -(\mathbf{Z_3}-\mathbf{1}) \frac{\mathbf{1}}{\mathbf{4}} \left(\partial_\mu \mathbf{G}_\nu^{\mathbf{i}} - \partial_\nu \mathbf{G}_\mu^{\mathbf{i}}\right)^{\mathbf{2}} - (\mathbf{Z_1}-\mathbf{1}) \frac{\mathbf{g}}{\mathbf{2}} \mu^\epsilon \mathbf{f^{ijk}} (\partial_\mu \mathbf{G}_\nu^{\mathbf{i}} - \partial_\nu \mathbf{G}_\mu^{\mathbf{i}}) \mathbf{G}^{\mathbf{j},\mu} \mathbf{G}^{\mathbf{k},\nu} + \cdots$$

$$\mathbf{Z_1} - \mathbf{1} = \frac{\mathbf{g^2}}{\mathbf{16\pi^2}} \frac{\mathbf{1}}{\epsilon} \left[\mathbf{N_C} \left(\frac{\mathbf{17}}{\mathbf{12}} - \frac{\mathbf{3}\lambda}{\mathbf{4}}\right) - \frac{\mathbf{4}}{\mathbf{3}} \mathbf{n_f T_R}\right], \qquad \mathbf{Z_3} - \mathbf{1} = \frac{\mathbf{g^2}}{\mathbf{16\pi^2}} \frac{\mathbf{1}}{\epsilon} \left[\mathbf{N_C} \left(\frac{\mathbf{13}}{\mathbf{6}} - \frac{\lambda}{\mathbf{2}}\right) - \frac{\mathbf{4}}{-} \mathbf{3 n_f T_R}\right], \cdots$$

Renormalization prescription: subtract the pole only (MS-scheme)

Additional diagrams with counter term vertices cancel the UV divergences coming from loop integrals order by order of perturbation theory iteratively.

Weak coupling: running coupling constant, asymptotic freedom

Weak coupling: running coupling constant, asymptotic freedom

$$\mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g^{(0)}}(\epsilon)$$

Weak coupling: running coupling constant, asymptotic freedom

$$\mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g^{(0)}}(\epsilon)$$

In the massless limit the classical Lagrangian of QCD is scale invariant. It is destroyed at the quantum level : performing renormalization a scale parameter had to be introduced.

If we change the parameter μ , a change is generated,in the renormalized parameters $\mathbf{g}(\mu)$.

$$\mu^2 \frac{\mathbf{d}\alpha_\mathbf{s}(\mu)}{\mathbf{d}\mu^2} = \beta(\alpha_\mathbf{s}) \ , \quad \alpha_\mathbf{s} = \frac{\mathbf{g^2}(\mu)}{\mathbf{4\pi}} \ , \ \beta(\alpha_\mathbf{s}) = -\alpha_\mathbf{s}{}^2 \mathbf{b_0} - \alpha_\mathbf{s}{}^3 \mathbf{b_1} + \mathcal{O}(\alpha_\mathbf{s}{}^4)$$

Weak coupling: running coupling constant, asymptotic freedom

$$\mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g^{(0)}}(\epsilon)$$

In the massless limit the classical Lagrangian of QCD is scale invariant. It is destroyed at the quantum level : performing renormalization a scale parameter had to be introduced.

If we change the parameter μ , a change is generated,in the renormalized parameters $\mathbf{g}(\mu)$.

$$\mu^2 \frac{\mathbf{d}\alpha_\mathbf{s}(\mu)}{\mathbf{d}\mu^2} = \beta(\alpha_\mathbf{s}) \ , \quad \alpha_\mathbf{s} = \frac{\mathbf{g^2}(\mu)}{4\pi} \ , \ \beta(\alpha_\mathbf{s}) = -\alpha_\mathbf{s}{}^2 \mathbf{b_0} - \alpha_\mathbf{s}{}^3 \mathbf{b_1} + \mathcal{O}(\alpha_\mathbf{s}{}^4)$$

$$\mathbf{b_0} = \frac{11\mathbf{N_C} - 4\mathbf{n_f}\mathbf{T_R}}{12\pi}$$

$$\mathbf{b_0} \geq \mathbf{0} \quad \text{for} \quad \mathbf{N_c} = \mathbf{3}, \mathbf{T_R} = \frac{\mathbf{1}}{\mathbf{2}} \quad \text{provided} \quad \mathbf{n_f} \leq \mathbf{16.5}$$

$$\alpha_\mathbf{S}(\mu, \mathbf{n_f}, \mathbf{\Lambda}) = \frac{\mathbf{1}}{\mathbf{b_0} \ln \frac{\mu^2}{\mathbf{\Lambda^2}}} + \mathcal{O}\left(\frac{\ln \ln \frac{\mu^2}{\mathbf{\Lambda^2}}}{\ln \frac{\mu^2}{\mathbf{\Lambda^2}}} \right)$$

Weak coupling: running coupling constant, asymptotic freedom

$$\mathbf{g}(\mu) = \mu^{-\epsilon} \mathbf{Z_1^{-1}} \mathbf{Z_3^{3/2}} \mathbf{g^{(0)}}(\epsilon)$$

In the massless limit the classical Lagrangian of QCD is scale invariant. It is destroyed at the quantum level : performing renormalization a scale parameter had to be introduced.

If we change the parameter μ, a change is generated,in the renormalized parameters $\mathbf{g}(\mu)$.

$$\mu^2 \frac{\mathbf{d}\alpha_\mathbf{s}(\mu)}{\mathbf{d}\mu^2} = \beta(\alpha_\mathbf{s}) \ , \quad \alpha_\mathbf{s} = \frac{\mathbf{g^2}(\mu)}{\mathbf{4}\pi} \ , \ \beta(\alpha_\mathbf{s}) = -\alpha_\mathbf{s}{}^2 \mathbf{b_0} - \alpha_\mathbf{s}{}^3 \mathbf{b_1} + \mathcal{O}(\alpha_\mathbf{s}{}^4)$$

$$\mathbf{b_0} = \frac{\mathbf{11 N_C - 4 n_f T_R}}{\mathbf{12}\pi}$$

$$\mathbf{b_0} \geq \mathbf{0} \ \text{ for } \ \mathbf{N_c = 3, T_R} = \frac{\mathbf{1}}{\mathbf{2}} \ \text{ provided } \ \mathbf{n_f} \leq \mathbf{16.5}$$

$$\alpha_\mathbf{S}(\mu, \mathbf{n_f}, \mathbf{\Lambda}) = \frac{\mathbf{1}}{\mathbf{b_0} \ln \frac{\mu^2}{\mathbf{\Lambda^2}}} + \mathcal{O}\left(\frac{\ln \ln \frac{\mu^2}{\mathbf{\Lambda^2}}}{\ln \frac{\mu^2}{\mathbf{\Lambda^2}}} \right)$$

Experiment: $\alpha_\mathbf{s}(\mathbf{m_Z}, \mathbf{5}, \mathbf{\Lambda}^{(5)}_{\overline{\mathbf{MS}}}) = \mathbf{0.1183 \pm 0.0010}$, $\mathbf{\Lambda}^{(5)}_{\overline{\mathbf{MS}}} \approx \mathbf{300}\,\mathrm{MeV}$

$\mathbf{\Lambda}^{(5)}_{\overline{\mathbf{MS}}}$: hidden fundamental scale of QCD

Measured values of $\alpha_s(M_Z)$





Asymptotic freedom, color confinement, lattice QCD

Asymptotic freedom, color confinement, lattice QCD

Asymptotically free QFT's can be studied in the non-perturbative regions (strong coupling) using lattice regularization.

Asymptotic freedom, color confinement, lattice QCD

Asymptotically free QFT's can be studied in the non-perturbative regions (strong coupling) using lattice regularization.

Wilson-formulation:

4-dimensional Euclidean hyper cubic lattice with periodic boundary conditions

Asymptotic freedom, color confinement, lattice QCD

Asymptotically free QFT's can be studied in the non-perturbative regions (strong coupling) using lattice regularization.

Wilson-formulation:

4-dimensional Euclidean hyper cubic lattice with periodic boundary conditions

Euclidean action is obtained by Wick rotation: $\mathbf{x^4 = ix^0}, \quad \partial^4 = -\mathbf{i}\partial^0 \qquad \mathbf{i}\int \mathbf{d^4x}\, \mathcal{L} \rightarrow -\mathbf{S_E}$

Asymptotic freedom, color confinement, lattice QCD

Asymptotically free QFT's can be studied in the non-perturbative regions (strong coupling) using lattice regularization.

Wilson-formulation:

4-dimensional Euclidean hyper cubic lattice with periodic boundary conditions

Euclidean action is obtained by Wick rotation: $\mathbf{x^4 = ix^0}, \quad \partial^4 = -\mathbf{i}\partial^0 \qquad \mathbf{i}\int \mathbf{d^4x}\, \mathcal{L} \rightarrow -\mathbf{S_E}$



Asymptotic freedom, color confinement, lattice QCD

Asymptotically free QFT's can be studied in the non-perturbative regions (strong coupling) using lattice regularization.

Wilson-formulation:

4-dimensional Euclidean hyper cubic lattice with periodic boundary conditions

Euclidean action is obtained by Wick rotation: $\mathbf{x^4 = ix^0}, \quad \partial^4 = -\mathbf{i}\partial^0 \qquad \mathbf{i}\int \mathbf{d^4x}\,\mathcal{L} \rightarrow -\mathbf{S_E}$

Finite volume: infrared-cut, finite lattice spacing: ultraviolet cut-off
very large but finite degree of freedom.



Asymptotic freedom, color confinement, lattice QCD

Asymptotically free QFT's can be studied in the non-perturbative regions (strong coupling) using lattice regularization.

Wilson-formulation:

4-dimensional Euclidean hyper cubic lattice with periodic boundary conditions

Euclidean action is obtained by Wick rotation: $\mathbf{x^4 = ix^0}, \quad \partial^4 = -\mathbf{i}\partial^0 \qquad \mathbf{i}\int \mathbf{d^4x}\,\mathcal{L} \rightarrow -\mathbf{S_E}$



Finite volume: infrared-cut, finite lattice spacing: ultraviolet cut-off
very large but finite degree of freedom.

$$\mathcal{Z} = \int \mathcal{D}\mathbf{A}_\mu \int \mathcal{D}\psi\mathcal{D}\bar{\psi}\mathbf{e}^{-(\bar{\psi}\mathbf{M}\psi+\mathbf{S_G})}$$

Lattice QCD defines the theory with infrared and ultraviolet cut-off

$x_\mu = n_\mu a \quad \text{with} \quad n_\mu \in \{0, \ldots, N_\mu - 1\}$



The parallel transport $\mathbf{U}_\mu(\mathbf{x})$ is the discretized version of the path ordered product of continuum gauge fields $\hat{\mathbf{G}}_\mu$

$$\mathbf{U}_\mu(\mathbf{x}) = \mathcal{P}\mathbf{e}^{\mathbf{ig}\int_{\mathbf{x}}^{\mathbf{x}+\hat{\mu}} \mathbf{dx}'_\mu \hat{\mathbf{G}}_\mu(\mathbf{x}')}$$

$$\mathbf{U}_\mu(\mathbf{x}) \to \mathbf{\Omega}(\mathbf{x})\mathbf{U}_\mu(\mathbf{x})\mathbf{\Omega}^\dagger(\mathbf{x}+\hat{\mu})$$

The trace over a closed loop of parallel transports is gauge invariant. The simplest of these loops is the plaquette

$$\mathbf{U}_{\mu\nu}(\mathbf{x}) = \mathbf{U}_\mu(\mathbf{x})\mathbf{U}_\nu(\mathbf{x}+\hat{\mu})\mathbf{U}^\dagger_\mu(\mathbf{x}+\hat{\nu})\mathbf{U}^\dagger_\nu(\mathbf{x})$$

$$\mathbf{S_G} = \beta \sum_{\mathbf{x},\mu>\nu} \left(\mathbf{1} - \frac{\mathbf{1}}{\mathbf{6}}\mathrm{Tr}(\mathbf{U}^\dagger_{\mu\nu}(\mathbf{x}) + \mathbf{U}_{\mu\nu}(\mathbf{x}))\right) \quad , \quad \beta = \frac{\mathbf{6}}{\mathbf{g^2}}$$

The path integral

$$\langle \mathbf{O} \rangle = \frac{\mathbf{1}}{\mathcal{Z}} \int \mathcal{D}\mathbf{U}_\mu \int \mathcal{D}\mathbf{\Psi}\mathcal{D}\bar{\mathbf{\Psi}}\mathbf{O}\mathbf{e}^{-(\bar{\mathbf{\Psi}}\mathbf{M}(\mathbf{U})\mathbf{\Psi} + \mathbf{S_G}(\mathbf{U}))}$$

is carried out by numerical Monte Carlo integration

Lattice QCD defines the theory with infrared and ultraviolet cut-off

Taking the continuum limit.

Let us imagine that measuring the correlation function of proton creation operators on the lattice with Monte Carlo simulation, we extract the value of the proton mass.The cut-off scale on the lattice is the lattice distance “a” therefore all physical quantities are obtained in its physical units.The value of the proton mass in such a lattice measurement depends on the coupling constant and on the lattice distance as

$$\mathbf{m_P} = \frac{\mathbf{1}}{\mathbf{a}}\mathbf{f}(\mathbf{g}(\mathbf{a}))\,, \qquad \lim_{\mathbf{a}\to\mathbf{0}} \mathbf{m_P} = \frac{\mathbf{1}}{\mathbf{a}}\mathbf{f}(\mathbf{g}(\mathbf{a})) \longrightarrow \text{const} \quad , \quad \mathbf{a}\frac{\mathbf{d}}{\mathbf{da}}\mathbf{m_P} = \mathbf{0}$$

$$\mathbf{a} \ll \mathbf{R_P} \ll \mathbf{L} = \mathbf{N}\,\mathbf{a}$$



$$\mathbf{f}(\mathbf{g}) = \mathbf{a}\frac{\partial \mathbf{g}}{\partial \mathbf{a}}\frac{\partial}{\partial \mathbf{g}}\mathbf{f}(\mathbf{g})$$

$$\mathbf{a}\frac{\partial \mathbf{g}}{\partial \mathbf{a}} = \frac{\mathbf{b}}{\mathbf{4\pi}}\mathbf{g^3}\left(\mathbf{1} + \mathbf{b'}\frac{\mathbf{g^2}}{\mathbf{4\pi}}\right)$$

$$\mathbf{\Lambda_L} = \frac{\mathbf{1}}{\mathbf{a}}\exp\left(-\frac{\mathbf{2\pi}}{\mathbf{g_L^2}}\right)\left(\frac{\mathbf{4\pi} + \mathbf{b'g_L^2}}{\mathbf{4\pi g_L^2}}\right)^{\frac{\mathbf{b'}}{\mathbf{2b}}}$$

$$\mathbf{m_P} = \mathbf{c\Lambda_L}$$

dimensional transmutation

$$\mathbf{\Lambda_L} \approx \mathbf{1.7}\text{MeV}$$

$$\mathbf{g_L^2} = \mathbf{1}\,,\ \mathbf{a^{-1}} \approx \mathbf{2}\,\text{GeV}\,,\ \mathbf{a} \approx \mathbf{0.1}\text{fermi}$$

The relative magnitude of $\mathbf{\Lambda}_{\text{QCD}}$ with respect to the quark masses has important physical significance. In particular, it is very important that the value of $\mathbf{\Lambda}_{\text{QCD}}$ is much bigger than the value of the light quark masses $\mathbf{m_u}, \mathbf{m_d}$. It is also important that in high energy experiments the typical energy scale of is much higher than the value of $\mathbf{\Lambda}_{\text{QCD}}$.

Lattice QCD defines the theory with infrared and ultraviolet cut-off

30 years of effort of a large community

Hadron spectrum from lattice QCD



S. Dürr *et. al.*, Science 322:1224-1227,2008

How to use perturbation theory for predictions for observables?

How to use perturbation theory for predictions for observables?

- **Asymptotic freedom (full control of UV region)**

  Clue to resolve the apparent contradiction namely that

  QCD is the theory of hadrons and their interactions

  QCD is the theory of quark and gluon interactions

How to use perturbation theory for predictions for observables?

- **Asymptotic freedom (full control of UV region)**

  Clue to resolve the apparent contradiction namely that

  QCD is the theory of hadrons and their interactions

  QCD is the theory of quark and gluon interactions

- **QCD pertubation theory is plagued by infrared singularities:**

  Soft gluon singularities and collinear singularities provided by large distance contributions of perturbation theory. They are controlled by the powerful cancellation and factorization theorem.

How to use perturbation theory for predictions for observables?

- **Asymptotic freedom (full control of UV region)**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **QCD pertubation theory is plagued by infrared singularities:**
  Soft gluon singularities and collinear singularities provided by large distance contributions of perturbation theory. They are controlled by the powerful cancellation and factorization theorem.

- **Simplest example:** The total cross section of e^+e^- annihilation into hadrons

How to use perturbation theory for predictions for observables?

- **Asymptotic freedom (full control of UV region)**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **QCD pertubation theory is plagued by infrared singularities:**
  Soft gluon singularities and collinear singularities provided by large distance contributions of perturbation theory. They are controlled by the powerful cancellation and factorization theorem.

- **Simplest example:** The total cross section of e^+e^- annihilation into hadrons

  Only one large scale. Soft and collinear singularities **cancel** in all order of PT.

How to use perturbation theory for predictions for observables?

- **Asymptotic freedom (full control of UV region)**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **QCD pertubation theory is plagued by infrared singularities:**
  Soft gluon singularities and collinear singularities provided by large distance contributions of perturbation theory. They are controlled by the powerful cancellation and factorization theorem.

- **Simplest example:** The total cross section of e^+e^- annihilation into hadrons

Only one large scale. Soft and collinear singularities **cancel** in all order of PT.

Assuming that the unitarity sum over all hadronic final state can be replaced with the unitarity sum over all final state quark and gluons appearing in the perturbative treatment, we get direct prediction for the measured cross-section.

How to use perturbation theory for predictions for observables?

- **Asymptotic freedom (full control of UV region)**
  Clue to resolve the apparent contradiction namely that
  QCD is the theory of hadrons and their interactions
  QCD is the theory of quark and gluon interactions

- **QCD pertubation theory is plagued by infrared singularities:**
  Soft gluon singularities and collinear singularities provided by large distance contributions of perturbation theory. They are controlled by the powerful cancellation and factorization theorem.

- **Simplest example:** The total cross section of $\mathbf{e^+e^-}$ annihilation into hadrons

Only one large scale. Soft and collinear singularities **cancel** in all order of PT.

Assuming that the unitarity sum over all hadronic final state can be replaced with the unitarity sum over all final state quark and gluons appearing in the perturbative treatment, we get direct prediction for the measured cross-section.

$$\sum_{\mathbf{h}} |\mathbf{h}><\mathbf{h}| = \sum_{\mathbf{q,g}} |\mathrm{gluons, quarks}><\mathrm{gluons, quarks}|$$

$$\sqrt{\mathbf{s}} = \mathbf{Q}, \quad \mathbf{Q}, \gg, \mathbf{m_a}, \quad \mathbf{a} = \mathbf{u, d, s, ...}$$

$e^+e^- \longrightarrow q\bar{q} \longrightarrow$ hadrons



$$\mathbf{R(Q^2)_{e^+e^-}} = \frac{\sigma(\mathbf{e^+e^-} \to \text{hadrons})}{\sigma(\mathbf{e^+e^-} \to \mu^+\mu^-)}$$

$$\sum_{\mathbf{h}} |\mathbf{h}><\mathbf{h}| = \sum_{\mathbf{q,g}} |\text{gluons, quarks}><\text{gluons, quarks}|$$

$$\sqrt{\mathbf{s}} = \mathbf{Q}, \quad \mathbf{Q}\,, \gg, \mathbf{m_a}, \quad \mathbf{a} = \mathbf{u}, \mathbf{d}, \mathbf{s}, \ldots$$

$$\mathbf{R(s)} = \mathbf{12}\,\pi\,\mathrm{Im}\,\mathbf{\Pi}^{\mathrm{EM}}(\mathbf{s}) = \mathbf{3}\sum_{\mathbf{i=1}}^{\mathbf{n_f}} \mathbf{Q_i^2}\left(\mathbf{1} + \frac{\alpha_\mathbf{s}}{\pi} + \ldots\right)$$

The measured values of $\mathbf{R(Q^2)_{e^+e^-}} = \dfrac{\sigma(\mathbf{e^+e^-} \to \text{hadrons})}{\sigma(\mathbf{e^+e^-} \to \mu^+\mu^-)}$



The measured values of $\mathbf{R(Q^2)_{e^+e^-}} = \dfrac{\sigma(\mathbf{e^+e^-} \to \text{hadrons})}{\sigma(\mathbf{e^+e^-} \to \mu^+\mu^-)}$



and its precise theoretical calculation to NNNLO order in perturbative QCD

The measured values of $\mathbf{R(Q^2)_{e^+e^-}} = \dfrac{\sigma(\mathbf{e^+e^-} \to \text{hadrons})}{\sigma(\mathbf{e^+e^-} \to \mu^+\mu^-)}$



and its precise theoretical calculation to NNNLO order in perturbative QCD

$$R(s) = 1 + a_s + 1.4097a_s^2 - 12.76703a_s^3 - 80.0075a_s^4..$$

$$\alpha_s(M_z)^{\text{NNNLO}} = 0.1190 \pm 0.0026_{\text{exp}}$$

Baikov, Chetyrkin,Kuhn

Hadrons in the initial state; QCD improved parton model

Hadrons in the initial state; QCD improved parton model

Infinite momentum frame: hadrons are beams of partons

Hadrons in the initial state; QCD improved parton model

Infinite momentum frame: hadrons are beams of partons

parton number densities
extracted from the data +evolution

hard parton x-section expanded
in the QCD coupling constant

$$\frac{d\sigma_{\mathrm{pp}\to\mathrm{hadrons}}}{dX} = \sum_{a,b} \int dx_1 dx_2 f_a(x_1,\mu_F) f_b(x_2,\mu_F) \times \frac{d\hat{\sigma}_{\mathrm{ab}\to\mathrm{partons}}(\alpha_s(\mu_R),\mu_R,\mu_F)}{dX} + \mathcal{O}\left(\frac{\Lambda^n_{\mathrm{QCD}}}{Q^n}\right)$$

Hadrons in the initial state; QCD improved parton model

Infinite momentum frame: hadrons are beams of partons

parton number densities
extracted from the data +evolution

hard parton x-section expanded
in the QCD coupling constant

$$\frac{d\sigma_{\mathrm{pp}\to\mathrm{hadrons}}}{dX} = \sum_{a,b} \int dx_1 dx_2 f_a(x_1,\mu_F) f_b(x_2,\mu_F) \times \frac{d\hat{\sigma}_{\mathrm{ab}\to\mathrm{partons}}(\alpha_s(\mu_R),\mu_R,\mu_F)}{dX} + \mathcal{O}\left(\frac{\Lambda^n_{\mathrm{QCD}}}{Q^n}\right)$$



Drell-Yan process

Hadrons in the initial state; QCD improved parton model

Infinite momentum frame: hadrons are beams of partons

parton number densities
extracted from the data +evolution

hard parton x-section expanded
in the QCD coupling constant

$$\frac{d\sigma_{\mathrm{pp}\to\mathrm{hadrons}}}{dX} = \sum_{a,b} \int dx_1 dx_2 f_a(x_1,\mu_F) f_b(x_2,\mu_F) \times \frac{d\hat{\sigma}_{\mathrm{ab}\to\mathrm{partons}}(\alpha_s(\mu_R),\mu_R,\mu_F)}{dX} + \mathcal{O}\left(\frac{\Lambda^n_{\mathrm{QCD}}}{Q^n}\right)$$



Drell-Yan process

Many processes, many different multi particle final states

Schematic view of hard processes at the LHC



Z. Bern

Hadrons in the initial state; QCD improved parton model





The hard parton cross-section:

$$\mathbf{d}\hat{\sigma}_{\mathbf{ab}\to\mathrm{partons}}(\alpha_{\mathbf{s}}(\mu_{\mathbf{R}}), \mu_{\mathbf{R}}, \mu_{\mathbf{F}})$$

requires “collinear infrared renormalization” .

The hard parton cross-section:

$$\mathbf{d}\hat{\sigma}_{\mathbf{ab}\to\mathrm{partons}}(\alpha_{\mathbf{s}}(\mu_{\mathbf{R}}),\mu_{\mathbf{R}},\mu_{\mathbf{F}})$$

requires "collinear infrared renormalization" .

In d-dimension the parton cross section is defined by a set of Feynman diagrams together with the ultraviolet renormalization counter terms to cancel the ultraviolet divergences appearing as poles in epsilon in the loop integrals.
The cross-section also have infrared singularities. They can also be regularized by performing the calculation in d-dimension. For infrared safe observables the final state soft and collinear singularities cancel. But the initial state collinear singularities do not cancel. However they factorize into process indepenedent functions $\mathbf{\Gamma_{i/a}(x,\epsilon)}$

The hard parton cross-section:

$$\mathbf{d}\hat{\sigma}_{\mathbf{ab}\to\mathrm{partons}}(\alpha_{\mathbf{s}}(\mu_{\mathbf{R}}),\mu_{\mathbf{R}},\mu_{\mathbf{F}})$$

requires "collinear infrared renormalization" .

In d-dimension the parton cross section is defined by a set of Feynman diagrams together with the ultraviolet renormalization counter terms to cancel the ultraviolet divergences appearing as poles in epsilon in the loop integrals.
The cross-section also have infrared singularities. They can also be regularized by performing the calculation in d-dimension. For infrared safe observables the final state soft and collinear singularities cancel. But the initial state collinear singularities do not cancel. However they factorize into process indepenedent functions $\mathbf{\Gamma_{i/a}(x,\epsilon)}$

$$\mathbf{d}\sigma^{(\mathbf{bare})}_{\mathbf{ab}}(\mathbf{p_a},\mathbf{p_b},\mathbf{d}) = \sum_{\mathbf{ij}}\int_{\mathbf{0}}^{\mathbf{1}}\mathbf{dx_a}\int_{\mathbf{0}}^{\mathbf{1}}\mathbf{dx_b}\mathbf{\Gamma_{i/a}}(\mathbf{x_a},\epsilon)\mathbf{\Gamma_{j/b}}(\mathbf{x_b},\epsilon)\mathbf{d}\hat{\sigma}_{\mathbf{ij}}(\mathbf{q},\mathbf{x_a p_a},\mathbf{x_b p_b},\alpha_{\mathbf{s}}(\mu),\mu)$$

$$\mathbf{\Gamma_{i/a}(x)} = \delta_{\mathbf{ia}}\delta(\mathbf{1}-\mathbf{x}) - \frac{\alpha_{\mathbf{s}}}{\mathbf{2}\pi}\left(\frac{\mathbf{1}}{\epsilon}\mathbf{P_{i/a}(x,0)} - \mathbf{K_{i/a}(x)}\right) + \mathcal{O}\left(\alpha_{\mathbf{s}}^{\mathbf{2}}\right),$$

The hard parton cross-section:

$$\mathbf{d}\hat{\sigma}_{\mathbf{ab}\to\mathrm{partons}}(\alpha_{\mathbf{s}}(\mu_{\mathbf{R}}),\mu_{\mathbf{R}},\mu_{\mathbf{F}})$$

requires "collinear infrared renormalization" .

In d-dimension the parton cross section is defined by a set of Feynman diagrams together with the ultraviolet renormalization counter terms to cancel the ultraviolet divergences appearing as poles in epsilon in the loop integrals.
The cross-section also have infrared singularities. They can also be regularized by performing the calculation in d-dimension. For infrared safe observables the final state soft and collinear singularities cancel. But the initial state collinear singularities do not cancel. However they factorize into process indepenedent functions $\mathbf{\Gamma_{i/a}(x,\epsilon)}$

$$\mathbf{d}\sigma^{(\mathbf{bare})}_{\mathbf{ab}}(\mathbf{p_a},\mathbf{p_b},\mathbf{d}) = \sum_{\mathbf{ij}}\int_{\mathbf{0}}^{\mathbf{1}}\mathbf{dx_a}\int_{\mathbf{0}}^{\mathbf{1}}\mathbf{dx_b}\mathbf{\Gamma_{i/a}}(\mathbf{x_a},\epsilon)\mathbf{\Gamma_{j/b}}(\mathbf{x_b},\epsilon)\mathbf{d}\hat{\sigma}_{\mathbf{ij}}(\mathbf{q},\mathbf{x_a p_a},\mathbf{x_b p_b},\alpha_{\mathbf{s}}(\mu),\mu)$$

$$\mathbf{\Gamma_{i/a}(x)} = \delta_{\mathbf{ia}}\delta(\mathbf{1}-\mathbf{x}) - \frac{\alpha_{\mathbf{s}}}{\mathbf{2}\pi}\left(\frac{\mathbf{1}}{\epsilon}\mathbf{P_{i/a}(x,0)} - \mathbf{K_{i/a}(x)}\right) + \mathcal{O}\left(\alpha_{\mathbf{s}}^{\mathbf{2}}\right),$$

Adding appropriate collinear counter terms one can obtain the finite short distance cross section in any finite order of perturbation theory

The hard parton cross-section:

$$\mathbf{d}\hat{\sigma}_{\mathbf{ab}\to\mathrm{partons}}(\alpha_{\mathbf{s}}(\mu_{\mathbf{R}}),\mu_{\mathbf{R}},\mu_{\mathbf{F}})$$

requires "collinear infrared renormalization" .

In d-dimension the parton cross section is defined by a set of Feynman diagrams together with the ultraviolet renormalization counter terms to cancel the ultraviolet divergences appearing as poles in epsilon in the loop integrals.
The cross-section also have infrared singularities. They can also be regularized by performing the calculation in d-dimension. For infrared safe observables the final state soft and collinear singularities cancel. But the initial state collinear singularities do not cancel. However they factorize into process indepenedent functions $\mathbf{\Gamma_{i/a}(x,\epsilon)}$

$$\mathbf{d}\sigma^{(\mathbf{bare})}_{\mathbf{ab}}(\mathbf{p_a},\mathbf{p_b},\mathbf{d}) = \sum_{\mathbf{ij}}\int_{\mathbf{0}}^{\mathbf{1}}\mathbf{dx_a}\int_{\mathbf{0}}^{\mathbf{1}}\mathbf{dx_b}\mathbf{\Gamma_{i/a}}(\mathbf{x_a},\epsilon)\mathbf{\Gamma_{j/b}}(\mathbf{x_b},\epsilon)\mathbf{d}\hat{\sigma}_{\mathbf{ij}}(\mathbf{q},\mathbf{x_a p_a},\mathbf{x_b p_b},\alpha_{\mathbf{s}}(\mu),\mu)$$

$$\mathbf{\Gamma_{i/a}(x)} = \delta_{\mathbf{ia}}\delta(\mathbf{1}-\mathbf{x}) - \frac{\alpha_{\mathbf{s}}}{\mathbf{2}\pi}\left(\frac{\mathbf{1}}{\epsilon}\mathbf{P_{i/a}(x,0)} - \mathbf{K_{i/a}(x)}\right) + \mathcal{O}\left(\alpha_{\mathbf{s}}^{\mathbf{2}}\right),$$

Adding appropriate collinear counter terms one can obtain the finite short distance cross section in any finite order of perturbation theory

$$\mathbf{d}\hat{\sigma}_{\mathbf{ab}}(\mathbf{p_a},\mathbf{p_b},...,\mu) = \mathbf{d}\hat{\sigma}^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_a},\mathbf{p_b},...) + \mathbf{d}\hat{\sigma}^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_a},\mathbf{p_b},...,\mu) + \ ... \quad .$$

$$\mathbf{d}\sigma_{\mathbf{ab}} = \mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{ab}} + \mathbf{d}\sigma^{(\mathbf{1})}_{\mathbf{ab}}\,, \quad \mathbf{d}\hat{\sigma}_{\mathbf{ab}} = \mathbf{d}\hat{\sigma}^{(\mathbf{0})}_{\mathbf{ab}} + \mathbf{d}\hat{\sigma}^{(\mathbf{1})}_{\mathbf{ab}}\,,$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1}, \mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1}, \mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) + \mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) + \mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \frac{\alpha_\mathbf{s}}{\mathbf{2}\pi}\sum_{\mathbf{d}}\int \mathbf{dx}\left(\frac{\mathbf{1}}{\overline{\epsilon}}\mathbf{P_{d/a}(x,0)} - \mathbf{K_{d/a}(x)}\right)\mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{db}}(\mathbf{xp_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(\mathbf{1})}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) + \mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \frac{\alpha_\mathbf{s}}{\mathbf{2}\pi}\sum_{\mathbf{d}}\int \mathbf{dx}\left(\frac{\mathbf{1}}{\bar{\epsilon}}\mathbf{P_{d/a}}(\mathbf{x},\mathbf{0}) - \mathbf{K_{d/a}}(\mathbf{x})\right)\mathbf{d}\sigma^{(\mathbf{0})}_{\mathbf{db}}(\mathbf{x p_1},\mathbf{p_2})$$

Similar to UV renormalization. Initial state collinear singularities do not cancel. But they can be cancelled by counter terms generated from bare parton number densities. This “infrared renormalization” requires regularization and renormalization prescription. The renormalized parton densities depend on the corresponding factorization scale and obey the evolution equation

$$\mathbf{d}\hat{\sigma}^{(0)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(0)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(1)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(1)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) + \mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \frac{\alpha_\mathbf{s}}{\mathbf{2}\pi}\sum_{\mathbf{d}}\int \mathbf{dx}\left(\frac{\mathbf{1}}{\bar{\epsilon}}\mathbf{P_{d/a}}(\mathbf{x},\mathbf{0}) - \mathbf{K_{d/a}}(\mathbf{x})\right)\mathbf{d}\sigma^{(0)}_{\mathbf{db}}(\mathbf{xp_1},\mathbf{p_2})$$

Similar to UV renormalization. Initial state collinear singularities do not cancel. But they can be cancelled by counter terms generated from bare parton number densities. This “infrared renormalization” requires regularization and renormalization prescription. The renormalized parton densities depend on the corresponding factorization scale and obey the evolution equation

$$\mu\frac{\mathbf{d}}{\mathbf{d}\mu}\mathbf{df_{a/A}}(\mathbf{x},\mu) = \sum_{\mathbf{b}}\int_{\mathbf{x}}^{\mathbf{1}}\frac{\mathbf{dz}}{\mathbf{z}}\mathbf{P_{a/b}}(\mathbf{z},\alpha_\mathbf{s}(\mu))\mathbf{f_{b/A}}(\mathbf{x/z},\mu)$$

$$\mathbf{d}\hat{\sigma}^{(0)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(0)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(1)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(1)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) + \mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \frac{\alpha_\mathbf{s}}{\mathbf{2}\pi}\sum_{\mathbf{d}}\int \mathbf{dx}\left(\frac{\mathbf{1}}{\bar{\epsilon}}\mathbf{P_{d/a}}(\mathbf{x},\mathbf{0}) - \mathbf{K_{d/a}}(\mathbf{x})\right)\mathbf{d}\sigma^{(0)}_{\mathbf{db}}(\mathbf{xp_1},\mathbf{p_2})$$

Similar to UV renormalization. Initial state collinear singularities do not cancel. But they can be cancelled by counter terms generated from bare parton number densities. This “infrared renormalization” requires regularization and renormalization prescription. The renormalized parton densities depend on the corresponding factorization scale and obey the evolution equation

$$\mu\frac{\mathbf{d}}{\mathbf{d}\mu}\mathbf{df_{a/A}}(\mathbf{x},\mu) = \sum_{\mathbf{b}}\int_{\mathbf{x}}^{\mathbf{1}}\frac{\mathbf{dz}}{\mathbf{z}}\mathbf{P_{a/b}}(\mathbf{z},\alpha_\mathbf{s}(\mu))\mathbf{f_{b/A}}(\mathbf{x/z},\mu)$$

Physical cross section: the scale dependence of the parton number densitiesis cancelled by the scale dependence of the hard scattering cross-section

$$\mathbf{d}\hat{\sigma}^{(0)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(0)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\hat{\sigma}^{(1)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \mathbf{d}\sigma^{(1)}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) + \mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2})$$

$$\mathbf{d}\sigma^{\mathrm{count}}_{\mathbf{ab}}(\mathbf{p_1},\mathbf{p_2}) = \frac{\alpha_\mathbf{s}}{\mathbf{2}\pi}\sum_{\mathbf{d}}\int \mathbf{dx}\left(\frac{\mathbf{1}}{\bar{\epsilon}}\mathbf{P_{d/a}}(\mathbf{x},\mathbf{0}) - \mathbf{K_{d/a}}(\mathbf{x})\right)\mathbf{d}\sigma^{(0)}_{\mathbf{db}}(\mathbf{xp_1},\mathbf{p_2})$$

Similar to UV renormalization. Initial state collinear singularities do not cancel. But they can be cancelled by counter terms generated from bare parton number densities. This “infrared renormalization” requires regularization and renormalization prescription. The renormalized parton densities depend on the corresponding factorization scale and obey the evolution equation

$$\mu\frac{\mathbf{d}}{\mathbf{d}\mu}\mathbf{df_{a/A}}(\mathbf{x},\mu) = \sum_{\mathbf{b}}\int_{\mathbf{x}}^{\mathbf{1}}\frac{\mathbf{dz}}{\mathbf{z}}\mathbf{P_{a/b}}(\mathbf{z},\alpha_\mathbf{s}(\mu))\mathbf{f_{b/A}}(\mathbf{x/z},\mu)$$

Physical cross section: the scale dependence of the parton number densitiesis cancelled by the scale dependence of the hard scattering cross-section

$$\mathbf{d}\sigma_{\mathbf{pp->}\mu^+\mu^-\mathbf{+hadrons}} = \int \mathbf{f_{a/A}}(\mathbf{x_a},\mu)\mathbf{f_{b/B}}(\mathbf{x_b},\mu)\mathbf{d}\hat{\sigma}_{\mathbf{ab}}(\mathbf{x_a x_b s},\mu,\alpha_\mathbf{s}(\mu))$$

Cancellation and factorization theorems:

They constitute the theoretical basis of description of scattering processes of hadrons in perturbative QCD. These theorems are valid for infrared safe measurables.

An observable is infrared safe if it is insensitive to collinear splittings of partons and/or emission of soft gluons.

Cancellation and factorization theorems:

They constitute the theoretical basis of description of scattering processes of hadrons in perturbative QCD. These theorems are valid for infrared safe measurables.

An observable is infrared safe if it is insensitive to collinear splittings of partons and/or emission of soft gluons.

In the case of infrared safe quantities the perturbative QCD predictions are well defined in terms of partons.

The fundamental assumption of the QCD improved parton model:

Predictions made for infrared save quantities in terms of partons partons give good approximation to the same quantities measured in terms of hadrons (the power corrections are small at high momentum scales).

The spectacular success of perturbative QCD:

In all the cases considered, the comparison of observables that are calculable in perturbative QCD with experimental results improved if next-to-leading order (NLO) QCD computations were used. This fact establishes perturbative QCD as a systematic framework to describe the physics of hard hadronic collisions. It also suggests that, ideally, the theoretical toolkit for the LHC should contain next-to-leading computation for a large variety of processes.

The spectacular success of perturbative QCD:

In all the cases considered, the comparison of observables that are calculable in perturbative QCD with experimental results improved if next-to-leading order (NLO) QCD computations were used. This fact establishes perturbative QCD as a systematic framework to describe the physics of hard hadronic collisions. It also suggests that, ideally, the theoretical toolkit for the LHC should contain next-to-leading computation for a large variety of processes.

Improvements at NLO

The spectacular success of perturbative QCD:

In all the cases considered, the comparison of observables that are calculable in perturbative QCD with experimental results improved if next-to-leading order (NLO) QCD computations were used. This fact establishes perturbative QCD as a systematic framework to describe the physics of hard hadronic collisions. It also suggests that, ideally, the theoretical toolkit for the LHC should contain next-to-leading computation for a large variety of processes.

Improvements at NLO

- ☑ Reduced theoretical uncertainties due to more meaningful scale dependence and more precisely predicted rates and shapes
- ☑ Data at HERA,LEP,Tevatron and LHC fully validate the improvements of the agreement between theory and experiment if NLO corrections are included
- ☑ Smaller uncertainties in extrapolating measured background cross-sections into signal regions
- ☑ Better estimate for PDF uncertainties

The spectacular success of perturbative QCD:

In all the cases considered, the comparison of observables that are calculable in perturbative QCD with experimental results improved if next-to-leading order (NLO) QCD computations were used. This fact establishes perturbative QCD as a systematic framework to describe the physics of hard hadronic collisions. It also suggests that, ideally, the theoretical toolkit for the LHC should contain next-to-leading computation for a large variety of processes.

Improvements at NLO

- ☑ Reduced theoretical uncertainties due to more meaningful scale dependence and more precisely predicted rates and shapes

- ☑ Data at HERA,LEP,Tevatron and LHC fully validate the improvements of the agreement between theory and experiment if NLO corrections are included

- ☑ Smaller uncertainties in extrapolating measured background cross-sections into signal regions

- ☑ Better estimate for PDF uncertainties

Further improvements at NNLO

The spectacular success of perturbative QCD:

In all the cases considered, the comparison of observables that are calculable in perturbative QCD with experimental results improved if next-to-leading order (NLO) QCD computations were used. This fact establishes perturbative QCD as a systematic framework to describe the physics of hard hadronic collisions. It also suggests that, ideally, the theoretical toolkit for the LHC should contain next-to-leading computation for a large variety of processes.

Improvements at NLO

- ☑ Reduced theoretical uncertainties due to more meaningful scale dependence and more precisely predicted rates and shapes

- ☑ Data at HERA,LEP,Tevatron and LHC fully validate the improvements of the agreement between theory and experiment if NLO corrections are included

- ☑ Smaller uncertainties in extrapolating measured background cross-sections into signal regions

- ☑ Better estimate for PDF uncertainties

Further improvements at NNLO

Further improvements at NNNLO

Hard parton cross-section at NLO

Hard parton cross-section at NLO



Hard parton cross-section at NLO





Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method





Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\mathbf{\Phi}_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\mathbf{\Phi}_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\mathbf{\Phi}_{\mathbf{n-1}}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual Real

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual Real

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual Real

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

$$\mathbf{d}\hat{\sigma}_{\mathbf{a_1 a_2}}^{\mathrm{hard}} = \mathbf{d}\sigma_{\mathbf{a_1 a_2}}^{\mathrm{born}} + \{\mathbf{d}\sigma_{\mathbf{a_1 a_2}}^{\mathrm{virt}} + \mathbf{d}\sigma_{\mathbf{a_1 a_2}}^{\mathrm{soft}} + \mathbf{d}\sigma_{\mathbf{a_1 a_2}}^{(\mathrm{coll,initial})} + \mathbf{d}\sigma_{\mathbf{a_1 a_2}}^{(\mathrm{coll,final})}\}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual Real

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

$$\mathbf{d}\hat{\sigma}^{\mathrm{hard}}_{\mathbf{a_1 a_2}} = \mathbf{d}\sigma^{\mathrm{born}}_{\mathbf{a_1 a_2}} + \{\mathbf{d}\sigma^{\mathrm{virt}}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{\mathrm{soft}}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{(\mathrm{coll,initial})}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{(\mathrm{coll,final})}_{\mathbf{a_1 a_2}}\}$$

$$+ \mathbf{d}\sigma^{(\mathrm{coll,counter})}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{(\mathrm{real,subtracted})}_{\mathbf{a_1 a_2}}$$

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual Real

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

$$\mathbf{d}\hat{\sigma}_{\mathbf{a_1a_2}}^{\mathrm{hard}} = \mathbf{d}\sigma_{\mathbf{a_1a_2}}^{\mathrm{born}} + \left\{\mathbf{d}\sigma_{\mathbf{a_1a_2}}^{\mathrm{virt}} + \mathbf{d}\sigma_{\mathbf{a_1a_2}}^{\mathrm{soft}} + \mathbf{d}\sigma_{\mathbf{a_1a_2}}^{(\mathrm{coll,initial})} + \mathbf{d}\sigma_{\mathbf{a_1a_2}}^{(\mathrm{coll,final})}\right.$$

$$\left. + \mathbf{d}\sigma_{\mathbf{a_1a_2}}^{(\mathrm{coll,counter})}\right\} + \mathbf{d}\sigma_{\mathbf{a_1a_2}}^{(\mathrm{real,subtracted})}$$

Finite

Finite

Hard parton cross-section at NLO

Soft and collinear singularities of virtual and real contributions are cancelled by subtraction method



..times tree amplitude



squared amplitude

Born Virtual Real

$$\mathbf{d}\sigma_{\mathbf{n}}^{(\mathbf{1})} \approx |\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-2}} + \mathbf{2}\mathrm{Re}(\mathbf{M}_{\mathbf{n}}^{(\mathbf{0})\dagger}\mathbf{M}_{\mathbf{n}}^{(\mathbf{1})})\mathbf{d}\Phi_{\mathbf{n-2}} + |\mathbf{M}_{\mathbf{n+1}}^{(\mathbf{0})}|^{\mathbf{2}}\mathbf{d}\Phi_{\mathbf{n-1}}$$

$$\mathbf{d}\hat{\sigma}^{\mathrm{hard}}_{\mathbf{a_1 a_2}} = \mathbf{d}\sigma^{\mathrm{born}}_{\mathbf{a_1 a_2}} + \Big\{\mathbf{d}\sigma^{\mathrm{virt}}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{\mathrm{soft}}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{(\mathrm{coll,initial})}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{(\mathrm{coll,final})}_{\mathbf{a_1 a_2}} + \mathbf{d}\sigma^{(\mathrm{coll,counter})}_{\mathbf{a_1 a_2}}\Big\} + \mathbf{d}\sigma^{(\mathrm{real,subtracted})}_{\mathbf{a_1 a_2}}$$

Finite

Finite

Calculating higher order corrections with many legs and loop is hard:

Calculating higher order corrections with many legs and loop is hard:

- ☑ Traditional Feynman-diagram approach with Passarino-Veltman reduction has factorial growth with the number of the legs

- ☑ Major improvements in the last 15 years

  recursion relations, double box loop integral, IBP identities and Lorenz invariance identities for loop integrals, Laporta algorithm, understanding the structure of soft and collinear singularities, unitarity method, OPP reduction....

- ☑ NNLO and multi-leg NLO spectacular progress

- ☑ NNLO evolution of patron densities (Moch, Vermaseren,Vogt,2005)

- ☑ Subtraction methods for the real contributions (diploe, FKS)

- ☑ Embedding into shower Monte Carlo programs

Engineering progress for NLO:

- Application of unitarity to the calculation of one loop diagrams.
- Improvements in traditional techniques for Passarino-Veltman reduction.
- Generation of one loop integrand by parametric fit, (in dimensions D, OPP).
- Automatic procedures for generation of graphs and the consequent integrands.
- Automatic procedures for the generation of counter terms to implement the real virtual subtraction.
- Tabulation and numerical implementation of all integrals.

Jet production at Tevatron and the LHC

$p\bar{p} \rightarrow$ jet + X Tevatron

$pp \rightarrow$ jet + X LHC





- NLO QCD predictions (Ellis, ZK, Soper 1990)

5-loop NNLO calculation, 2-loop NNLO, 1-loop NLO multi-leg

Hadronic Z and tau decays, Baikov, Chetyrkin,Kuhn (2008)

$$R(s) = 1 + a_s + 1.4097 a_s^2 - 12.76703 a_s^3 - 80.0075 a_s^4 ..$$

$$\alpha_s(M_z)^{\mathrm{NNNLO}} = 0.1190 \pm 0.0026_{\mathrm{exp}}$$

5-loop NNLO calculation, 2-loop NNLO, 1-loop NLO multi-leg

Hadronic Z and tau decays, Baikov, Chetyrkin,Kuhn (2008)

$$R(s) = 1 + a_s + 1.4097 a_s^2 - 12.76703 a_s^3 - 80.0075 a_s^4 ..$$

$$\alpha_s(M_z)^{\mathrm{NNNLO}} = 0.1190 \pm 0.0026_{\mathrm{exp}}$$

Drell - Yan production
Anastasiou, Dixon, Melnikov
Petriello (2004)



5-loop NNLO calculation, 2-loop NNLO, 1-loop NLO multi-leg

Hadronic Z and tau decays, Baikov, Chetyrkin,Kuhn (2008)

$$R(s) = 1 + a_s + 1.4097 a_s^2 - 12.76703 a_s^3 - 80.0075 a_s^4 ..$$

$$\alpha_s(M_z)^{\mathrm{NNNLO}} = 0.1190 \pm 0.0026_{\mathrm{exp}}$$

Drell - Yan production
Anastasiou, Dixon, Melnikov
Petriello (2004)



$Z/\gamma^* + 4\mathrm{jet}$ production
BlackHat (2011)



What is available?



What is available?



What is available?



Higgs search by CMS





Figure 7: Distribution of $m_{\ell\ell}$ for the zero-jet eμ category in the H $\rightarrow$ WW search at 8 TeV. The signal expected from a Higgs boson with a mass $m_{\mathrm{H}} = 125$ GeV is shown added to the background.

New physics search, complex final states

New physics search, complex final states

SUSY searches

New physics search, complex final states

SUSY searches



New physics search, complex final states

SUSY searches



Typical event signature in CMS at LHC:



n leptons + n jets + missing E_T

New physics search, complex final states

SUSY searches



Typical event signature in CMS at LHC:



* $p + p \rightarrow 1, 2, \ldots 7$ jets, inclusive
* $p + p \rightarrow W/Z + 1, 2, \ldots 5$ jets,
* $p + p \rightarrow t + \bar{t} + 1,\ 2,\ 3$ jets

n leptons + n jets + missing E_T

Schematic view of hard processes at the LHC

The NLO revolution

G. Salam, ICHEP 2010



2009: NLO W+3j [Rocket: Ellis, Melnikov & Zanderighi] [unitarity]
2009: NLO W+3j [BlackHat: Berger et al] [unitarity]
2009: NLO $t\bar{t}b\bar{b}$ [Bredenstein et al] [traditional]
2009: NLO $t\bar{t}b\bar{b}$ [HELAC-NLO: Bevilacqua et al] [unitarity]
2009: NLO $q\bar{q} \to b\bar{b}b\bar{b}$ [Golem: Binoth et al] [traditional]
2010: NLO $t\bar{t}jj$ [HELAC-NLO: Bevilacqua et al] [unitarity]
2010: NLO Z+3j [BlackHat: Berger et al] [unitarity]

Les Houches Experimenters' Wish List

2010

process wanted at NLO	background to
1. $pp \to VV +$ jet	$t\bar{t}H$, new physics Dittmaier, Kallweit, Uwer; Campbell, Ellis, Zanderighi
2. $pp \to H + 2$ jets	H in VBF BCDEGMRSW; Campbell, Ellis, Williams Campbell, Ellis, Zanderighi; Ciccolini, Denner Dittmaier
3. $pp \to t\bar{t}b\bar{b}$	$t\bar{t}H$ Bredenstein, Denner Dittmaier, Pozzorini; Bevilacqua, Czakon, Papadopoulos, Pittau, Worek
4. $pp \to t\bar{t} + 2$ jets	$t\bar{t}H$ Bevilacqua, Czakon, Papadopoulos, Worek
5. $pp \to VVb\bar{b}$	VBF $\to H \to VV$, $t\bar{t}H$, new physics
6. $pp \to VV + 2$ jets	VBF $\to H \to VV$ VBF: Bozzi, Jäger, Oleari, Zeppenfeld
7. $pp \to V + 3$ jets	new physics Berger Bern, Dixon, Febres Cordero, Forde, Gleisberg, Ita, Kosower, Maitre; Ellis, Melnikov, Zanderighi
8. $pp \to VVV$	SUSY trilepton Lazopoulos, Melnikov, Petriello; Hankele, Zeppenfeld; Binoth, Ossola, Papadopoulos, Pittau
9. $pp \to b\bar{b}b\bar{b}$	Higgs, new physics GOLEM

Feynman diagram methods

now joined by

on-shell methods

table courtesy of C. Berger

N=4 sYM Lagrangian

The field content of the model:
one gauge field $\mathbf{A_\mu}$, four Majorana fermions $\psi_\mathbf{i}$, three real scalars ,$\mathbf{X_p}$ and three real pseudo-scalars $\mathbf{Y_q}$. All fields are in the adjoint representation.

$$\mathcal{L} = \mathrm{tr}\Big\{ -\frac{\mathbf{1}}{\mathbf{2}}\mathbf{F}_{\mu\nu}\mathbf{F}^{\mu\nu} + \bar{\psi}_\mathbf{i}\not{\mathbf{D}}\psi_\mathbf{i} + \mathbf{D}^\mu\mathbf{X_p}\mathbf{D}_\mu\mathbf{X_p} + \mathbf{D}^\mu\mathbf{Y_q}\mathbf{D}_\mu\mathbf{Y_q}$$

$$- \mathbf{ig}\bar{\psi}_\mathbf{i}\alpha^\mathbf{p}_\mathbf{ij}[\mathbf{X_p},\psi_\mathbf{j}] + \mathbf{g}\bar{\psi}_\mathbf{i}\gamma_\mathbf{5}\beta^\mathbf{q}_\mathbf{ij}[\mathbf{Y_q},\psi_\mathbf{j}]$$

$$+ \frac{\mathbf{g^2}}{\mathbf{2}}\Big([\mathbf{X_l},\mathbf{X_k}][\mathbf{X_l},\mathbf{X_k}] + [\mathbf{Y_l},\mathbf{Y_k}][\mathbf{Y_l},\mathbf{Y_k}] + \mathbf{2}[\mathbf{X_l},\mathbf{Y_k}][\mathbf{X_l},\mathbf{Y_k}] \Big) \Big\},$$

$$\alpha^\mathbf{1} = \begin{pmatrix} i\sigma_2 & 0 \\ 0 & i\sigma_2 \end{pmatrix}, \quad \alpha^\mathbf{2} = \begin{pmatrix} 0 & -\sigma_1 \\ \sigma_1 & 0 \end{pmatrix}, \quad \alpha^\mathbf{3} = \begin{pmatrix} 0 & \sigma_3 \\ -\sigma_3 & 0 \end{pmatrix},$$

$$\beta^\mathbf{1} = \begin{pmatrix} -i\sigma_2 & 0 \\ 0 & i\sigma_2 \end{pmatrix}, \quad \beta^\mathbf{2} = \begin{pmatrix} 0 & -i\sigma_2 \\ -i\sigma_2 & 0 \end{pmatrix}, \quad \beta^\mathbf{3} = \begin{pmatrix} 0 & \sigma_0 \\ -\sigma_0 & 0 \end{pmatrix}$$

$$\beta(\mathbf{g}) = \mu^\mathbf{2}\frac{\partial}{\partial\mu^\mathbf{2}} = -\frac{\mathbf{g^3}}{\mathbf{16\pi^2}}\left(\frac{\mathbf{11}}{\mathbf{3}} - \frac{\mathbf{2}}{\mathbf{3}}\mathbf{n_f} - \frac{\mathbf{1}}{\mathbf{6}}\mathbf{n_s}\right)\mathbf{C_A}$$

$$\beta(\mathbf{g}) = \mu^\mathbf{2}\frac{\partial}{\partial\mu^\mathbf{2}} = -\frac{\mathbf{g^3}}{\mathbf{16\pi^2}}\left(\frac{\mathbf{11}}{\mathbf{3}} - \frac{\mathbf{2}}{\mathbf{3}}\mathbf{4} - \frac{\mathbf{1}}{\mathbf{6}}\mathbf{6}\right)\mathbf{C_A} = \mathbf{0}$$

Beta function vanishes to all orders in PT.

N=4 sYM Lagrangian

- The scale invariance of the classical Lagrange density remains a symmetry at the quantum level.

- Nc limit, ‘t Hooft coupling $\lambda = N_c g^2$, relation to string theory?

- It has a much reacher set of symmetries:
  superconformal symmetry, integrability (hidden symmetry)

- Local gauge invariant composite operators (observables) with calculable anonalous dimensions

- AdS/CFT relates it to D=2 integrability, string theory in AdS, strong and weak coupling

On-shell methods of calculating amplitudes in QCD and N=4 sYM at large N

Helicity method
unitarity cuts (BDDK), OPP, cuts in D-dim.
soft and collinear limits
Anomalous dimensions at NNLO
Regge limit
Automated NLO codes for phenomenology (MCFM, BH, Sherpa, Powheg)

Complexification of external moment with twistors
CSW and BCFW recursion relations
BDS Ansatz; AdS/CFT
Amplitudes and Wilson loops
Dual conformal invariance,
Yangian symmetry of tree amplitudes,
Integrability, Bethe Ansatz
Relation to N=8 supergratvity

QCD N=4 sYM

Planar N=4 sYM and QCD in perturbation theory

- Tree-level amplitudes calculated in N=4 SYM could be carried over to QCD since all possible spins appear
- Generalized unitarity was invented in the simple case of

  N=4 SYM

  Any 4D gauge theory can be viewed as N=4 sYM with some particles

  and interactions added and removed
- Regge-limit and transcendentality in N=4 sYM and QCD
- New tree level recursion relations BCFW
- Dual conformal invariance, Yangian symmetry for tree amplitudes,

  2- loop results for 5, 6 point amplitudes

Parma International School of Theoretical Physics

September 3 - September 8, 2012

Parma, Italy

Scattering Amplitudes in QCD, Supersymmetric Gauge Theories and Supergravity

One-loop calculations in QCD and N=4 supersymmetric gauge theories

OUTLINE

Lecture 1: QCD and review its main features.

Lecture 2: One loop tensor integrals and their reduction

Lecture 3: Unitarity method and amplitudes

Lecture 4: Analytic and numerical computations

Lecture 5: Different implementations, Outlook

One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts.
R.Keith Ellis,,Zoltan Kunszt, Kirill Melnikov, Giulia Zanderighi,
to appear in Phys. Rep. arXiv:1105.4319 [hep-ph]

One loop calculation with traditional methods



One loop calculation with traditional methods



$\mathcal{N}(1)$ =1 one loop scalar, otherwise one loop tensor integral

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\mathbf{r \geq 2N - 4}$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \rightarrow \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \rightarrow \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \quad , \; \mathbf{d(l)=l^2-\mu^2}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \rightarrow \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \quad , \ \mathbf{d(l)=l^2-\mu^2}$$

$$\mathbf{I_4} \rightarrow \int \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \frac{\mathbf{l_\mu l_\nu l_\rho l_\delta}}{[\mathbf{d(l)}]^4} = \frac{(\mathbf{g_{\mu\nu}g_{\rho\delta}+g_{\mu\rho}g_{\nu\delta}+g_{\mu\delta}g_{\nu\rho}})}{\mathbf{D(D+2)}} \int \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \frac{\mathbf{l^4}}{[\mathbf{d(l)}]^4}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu} g_{\rho\delta} + g_{\mu\rho} g_{\nu\delta} + g_{\mu\delta} g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = \; (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s + D/2)\Gamma(m - s - D/2)}{\Gamma(D/2)\Gamma(m)},$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu} g_{\rho\delta} + g_{\mu\rho} g_{\nu\delta} + g_{\mu\delta} g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\mathbf{\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = \; (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s+D/2)\Gamma(m-s-D/2)}{\Gamma(D/2)\Gamma(m)},}$$

$$\mathbf{\Omega_D = 2\pi^{D/2}/\Gamma(D/2)}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu} g_{\rho\delta} + g_{\mu\rho} g_{\nu\delta} + g_{\mu\delta} g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\mathbf{\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = \; (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s+D/2)\Gamma(m-s-D/2)}{\Gamma(D/2)\Gamma(m)},}$$

$$\mathbf{\Omega_D = 2\pi^{D/2}/\Gamma(D/2)}$$

Finite result for $\quad \mathbf{D \neq 4\,, m = 4\,, s = 2}$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu}g_{\rho\delta} + g_{\mu\rho}g_{\nu\delta} + g_{\mu\delta}g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\mathbf{\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s+D/2)\Gamma(m-s-D/2)}{\Gamma(D/2)\Gamma(m)},}$$

$$\mathbf{\Omega_D = 2\pi^{D/2}/\Gamma(D/2)}$$

IR divergent if sufficient number of propagators can go to mass-shell. Soft and collinear singularities.

Finite result for $\quad \mathbf{D \neq 4\,, m = 4\,, s = 2}$

Integral basis in the limit $\mathrm{D} \to 4$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j}} \mathbf{I_{4;j}} + \mathbf{c_{3;j}} \mathbf{I_{3;j}} + \mathbf{c_{2;j}} \mathbf{I_{2;j}} + \mathbf{c_{1;j}} \mathbf{I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D}-4).$$

Only one-,two-,three-,four-point scalar integrals occur.

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j}} \mathbf{I_{4;j}} + \mathbf{c_{3;j}} \mathbf{I_{3;j}} + \mathbf{c_{2;j}} \mathbf{I_{2;j}} + \mathbf{c_{1;j}} \mathbf{I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D}-4).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j}} \mathbf{I_{4;j}} + \mathbf{c_{3;j}} \mathbf{I_{3;j}} + \mathbf{c_{2;j}} \mathbf{I_{2;j}} + \mathbf{c_{1;j}} \mathbf{I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D}-4).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j}} + \mathbf{c_{3;j} I_{3;j}} + \mathbf{c_{2;j} I_{2;j}} + \mathbf{c_{1;j} I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D-4}).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1}},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j}} + \mathbf{c_{3;j} I_{3;j}} + \mathbf{c_{2;j} I_{2;j}} + \mathbf{c_{1;j} I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D-4}).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1}},$$

$$\mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j} + \mathcal{R} + \mathcal{O}(D-4)}.$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1}},$$

$$\mathbf{d_i = (l + q_{i-1})^2 - m_i^2 + i\varepsilon} \qquad \mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j} + \mathcal{R} + \mathcal{O}(D-4)}.$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1}},$$

$$\mathbf{d_i = (l + q_{i-1})^2 - m_i^2 + i\varepsilon} \qquad \mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{q_n \equiv \sum_{i=1}^{n} p_i}, \quad \mathbf{q_0 = 0}$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j} + \mathcal{R} + \mathcal{O}(D-4)}.$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3 d_4},$$

$$\mathbf{d_i = (l + q_{i-1})^2 - m_i^2 + i\varepsilon} \qquad \mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{q_n \equiv \sum_{i=1}^{n} p_i}, \quad \mathbf{q_0 = 0}$$

$$\mathbf{r_\Gamma = \Gamma^2(1-\epsilon)\Gamma(1+\epsilon)/\Gamma(1-2\epsilon)}$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D-4}).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l + q_{i-1}})^2 - \mathbf{m_i^2} + \mathbf{i}\varepsilon \qquad \mathbf{s_{ij}} = (\mathbf{p_i + p_j})^2 ,$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2}},$$

$$\mathbf{q_n} \equiv \sum_{\mathbf{i=1}}^{\mathbf{n}} \mathbf{p_i}\,, \quad \mathbf{q_0 = 0}$$

$$\mathbf{r_\Gamma} = \mathbf{\Gamma^2}(1-\epsilon)\mathbf{\Gamma}(1+\epsilon)/\mathbf{\Gamma}(1-\mathbf{2}\epsilon)$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3}},$$



$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3 d_4}} ,$$

Only the constant terms and some of the terms depending on $(\mathbf{l} \cdot \mathbf{n}_\epsilon)^2$ give non-vanishing integrals

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}\mathbf{i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

$$\mathcal{R} = -\sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \frac{\tilde{\mathbf{d}}^{(\mathbf{4})}_{\mathbf{i_1 i_2 i_3 i_4}}}{\mathbf{6}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \frac{\tilde{\mathbf{c}}^{(\mathbf{7})}_{\mathbf{i_1 i_2 i_3}}}{\mathbf{2}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \left[\frac{\mathbf{m^2_{i_1}} + \mathbf{m^2_{i_2}}}{\mathbf{2}} - \frac{(\mathbf{q_{i_1}} - \mathbf{q_{i_2}})^{\mathbf{2}}}{\mathbf{6}} \right] \tilde{\mathbf{b}}^{(\mathbf{9})}_{\mathbf{i_1 i_2}}.$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

Reduction with the method of form factor expansions (Passarino, Veltman, 1979):

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

Reduction with the method of form factor expansions (Passarino, Veltman, 1979):

Examples:

$$\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}\,, \quad \mathbf{C^{\mu\nu} = g^{\mu\nu} C_{00} + \sum_{i,j=1}^{2} p_i^\mu p_j^\nu C_{ij}}\,, \quad \mathbf{C_{12} = C_{21}}$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

Reduction with the method of form factor expansions (Passarino, Veltman, 1979):

Examples:

$$\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}\,, \quad \mathbf{C^{\mu\nu} = g^{\mu\nu} C_{00} + \sum_{i,j=1}^{2} p_i^\mu p_j^\nu C_{ij}}\,, \quad \mathbf{C_{12} = C_{21}}$$

rank 1 and 2 tensor three point integrals

Reduction of $\mathbf{C}^\mu = \mathbf{p}_1^\mu \mathbf{C_1} + \mathbf{p}_2^\mu \mathbf{C_2}$,

Reduction of $\mathbf{C}^\mu = \mathbf{p_1^\mu C_1 + p_2^\mu C_2}\,,$

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\langle \mathbf{l \cdot p_j} \rangle = \int \frac{\mathbf{d^D l}}{\mathbf{i\pi^{D/2}}} \frac{\mathbf{l \cdot p_j}}{\mathbf{d_1 d_2 d_3}}, \quad \text{for } \mathbf{j = 1, 2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

Reduction of $\mathbf{C}^\mu = \mathbf{p}_1^\mu \mathbf{C_1} + \mathbf{p}_2^\mu \mathbf{C_2}$,

$$\mathbf{G_2} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\langle \mathbf{l} \cdot \mathbf{p_j} \rangle = \int \frac{\mathbf{d^D l}}{\mathbf{i}\pi^{\mathbf{D/2}}} \frac{\mathbf{l} \cdot \mathbf{p_j}}{\mathbf{d_1 d_2 d_3}}, \quad \text{for } \mathbf{j} = \mathbf{1}, \mathbf{2}$$

$$\mathbf{R_1^{[c]}} = \frac{\mathbf{1}}{\mathbf{2}} (\mathbf{f_1 C_0(1,2,3)} + \mathbf{B_0(1,3)} - \mathbf{B_0(2,3)}),$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

$$\mathbf{R_2^{[c]}} = \frac{\mathbf{1}}{\mathbf{2}} (\mathbf{f_2 C_0(1,2,3)} + \mathbf{B_0(1,2)} - \mathbf{B_0(1,3)})$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\langle \mathbf{l \cdot p_j} \rangle = \int \frac{\mathbf{d^D l}}{\mathbf{i\pi^{D/2}}} \frac{\mathbf{l \cdot p_j}}{\mathbf{d_1 d_2 d_3}}, \quad \text{for } \mathbf{j = 1, 2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l \cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1), \quad f_1 = m_2^2 - m_1^2 - p_1^2},$$

$$\mathbf{l \cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2), \quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1 \cdot p_2}\,.$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2, m_2, m_3) = \int \frac{d^D l}{i\pi^{D/2}}\frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\,\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}}\begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2,m_2,m_3) = \int \frac{d^D l}{i\pi^{D/2}}\,\frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l \cdot p_j \rangle = \int \frac{d^D l}{i\pi^{D/2}} \frac{l \cdot p_j}{d_1 d_2 d_3}, \quad \text{for } j = 1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l \cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1), \quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l \cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2), \quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1 \cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2, m_2, m_3) = \int \frac{d^D l}{i\pi^{D/2}} \frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}}\begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2,m_2,m_3) = \int \frac{d^D l}{i\pi^{D/2}}\frac{1}{(l^2-m_2^2)((l+p_2)^2-m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{p_1}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_1 C_{11} + p_1{\cdot}p_2 C_{12} + C_{00}) + p_2^\nu(p_1{\cdot}p_1 C_{12} + p_1{\cdot}p_2 C_{22})},$$
$$\mathbf{p_2}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_2 C_{11} + p_2{\cdot}p_2 C_{12}) + p_2^\nu(p_1{\cdot}p_2 C_{12} + p_2{\cdot}p_2 C_{22} + C_{00})}\,.$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}}\begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2,m_2,m_3) = \int \frac{d^D l}{i\pi^{D/2}}\frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{p_1}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_1 C_{11} + p_1{\cdot}p_2 C_{12} + C_{00}) + p_2^\nu(p_1{\cdot}p_1 C_{12} + p_1{\cdot}p_2 C_{22})},$$

$$\mathbf{p_2}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_2 C_{11} + p_2{\cdot}p_2 C_{12}) + p_2^\nu(p_1{\cdot}p_2 C_{12} + p_2{\cdot}p_2 C_{22} + C_{00})}\,.$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{C_\mu^\nu = B_0(2,3) + m_1^2 C_0(1,2,3) = DC_{00} + p_1^2 C_{11} + 2p_1{\cdot}p_2 C_{12} + p_2^2 C_{22}}$$

trace with $\mathbf{g^{\mu\nu}}$

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

Reduction chains for Passarino-Veltman procedure.

For more then 4 particles it is cumbersome

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

For more then 4 particles it is cumbersome

- The number of Feynman diagrams grows dramatically with the number of external particle (factorial).
- The number of terms generated during the reduction of tensor integrals grows rapidly with the number of external particles and with the rank of the integral.
- In cases with degenerate kinematics, the reduction procedure may lead to numerical instabilities.

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

For more then 4 particles it is cumbersome

- The number of Feynman diagrams grows dramatically with the number of external particle (factorial).
- The number of terms generated during the reduction of tensor integrals grows rapidly with the number of external particles and with the rank of the integral.
- In cases with degenerate kinematics, the reduction procedure may lead to numerical instabilities.

Well established technique, applied in many cases.

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

For more then 4 particles it is cumbersome

- The number of Feynman diagrams grows dramatically with the number of external particle (factorial).
- The number of terms generated during the reduction of tensor integrals grows rapidly with the number of external particles and with the rank of the integral.
- In cases with degenerate kinematics, the reduction procedure may lead to numerical instabilities.

Well established technique, applied in many cases.

For 5-,6-, and more particles external momenta not linearly independent (additional input), Denner, Dittmaier 2006

Singular regions in PV reduction:

Singular regions in PV reduction:

Example: rank 1 triangle

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$

but the integral is well defined in this limit

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ and necessarily $\mathbf{R_1^{[c]} = 0}$ and $\mathbf{R_2^{[c]} = 0}$

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ and necessarily $\mathbf{R_1^{[c]} = 0}$ and $\mathbf{R_2^{[c]} = 0}$

Exceptional points: vanishing Gram determinants, numerical instabilities
two and three point scalar integrals become linearly dependent
also higher powers of Gram-determinants

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ and necessarily $\mathbf{R_1^{[c]} = 0}$ and $\mathbf{R_2^{[c]} = 0}$

Exceptional points: vanishing Gram determinants, numerical instabilities
two and three point scalar integrals become linearly dependent
also higher powers of Gram-determinants

Exercise: Investigate the collinear behaviour of reduction using form factors with

using $\mathbf{p_1^\mu}$ and $\mathbf{\tilde{p}_2^\mu = p_2^\mu - \frac{p_1 \cdot p_2}{p_1^2} p_1^\mu}$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l \cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left[((\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2} \right] = \frac{\mathbf{1}}{\mathbf{2}} \left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2} \right]$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^{\lambda}\epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^{\lambda}\epsilon^{\mathbf{q_1}\mathbf{q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda\mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^{\lambda}\epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^{\lambda}\epsilon^{\mathbf{q_1}\mathbf{q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda\mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^{\lambda}} = \frac{\epsilon^{\lambda\mathbf{q_2}}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_2^{\lambda}} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^{\lambda}\epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^{\lambda}\epsilon^{\mathbf{q_1}\mathbf{q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda\mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^{\lambda}} = \frac{\epsilon^{\lambda\mathbf{q_2}}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_2^{\lambda}} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

$$\mathbf{l}^{\lambda} = (\mathbf{l}\cdot\mathbf{q_1})\mathbf{v_1^{\lambda}} + (\mathbf{l}\cdot\mathbf{q_2})\mathbf{v_2^{\lambda}}.$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^\lambda \epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^\lambda \epsilon^{\mathbf{q_1 q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda \mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^\lambda} = \frac{\epsilon^{\lambda \mathbf{q_2}}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_2^\lambda} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

$$\mathbf{l}^\lambda = (\mathbf{l}\cdot\mathbf{q_1})\mathbf{v_1^\lambda} + (\mathbf{l}\cdot\mathbf{q_2})\mathbf{v_2^\lambda}.$$

Can we embed this a coordinate into higher dimensional space time?

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^\lambda \epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^\lambda \epsilon^{\mathbf{q_1 q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda \mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^\lambda} = \frac{\epsilon^{\lambda \mathbf{q_2}}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_2^\lambda} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

$$\mathbf{l}^\lambda = (\mathbf{l}\cdot\mathbf{q_1})\mathbf{v_1^\lambda} + (\mathbf{l}\cdot\mathbf{q_2})\mathbf{v_2^\lambda}.$$

Can we embed this a coordinate into higher dimensional space time?

Yes.

Generalized Kronecker-symbols:

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \qquad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu = \frac{\epsilon_{q_1 q_2}\epsilon^{\mu q_2}}{\epsilon_{q_1 q_2}\epsilon^{q_1 q_2}}}, \quad \mathbf{v_2^\mu = \frac{\epsilon_{q_1 q_2}\epsilon^{q_1 \mu}}{\epsilon_{q_1 q_2}\epsilon^{q_1 q_2}}}, \quad \epsilon^{\mu_1\mu_2}\epsilon_{\nu_1\nu_2} = \delta^{\mu_1}_{\nu_1}\delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2}\delta^{\mu_2}_{\nu_1} = \det|\delta^\mu_\nu| \equiv \delta^{\mu_1\mu_2}_{\nu_1\nu_2},$$

$$\mathbf{v_1^\mu = \frac{\delta^{\mu q_2}_{q_1 q_2}}{\Delta_2}}, \quad \mathbf{v_2^\mu = \frac{\delta^{q_1 \mu}_{q_1 q_2}}{\Delta_2}}, \quad \mathbf{\Delta_2 = \delta^{q_1 q_2}_{q_1 q_2} = q_1^2 q_2^2 - (q_1 q_2)^2} \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \quad ,$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \ ,$$

$$\mathbf{\Delta}(\mathbf{k_1}, \mathbf{k_2}, \cdots, \mathbf{k_R}) = \delta^{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}_{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^{\mathbf{2}} \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \ ,$$

$$\mathbf{\Delta}(\mathbf{k_1}, \mathbf{k_2}, \cdots, \mathbf{k_R}) = \delta^{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}_{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}$$

$$\mathbf{v_i^\mu}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1} \ldots \mathbf{k_{i-1}} \mu \mathbf{k_{i+1}} \ldots \mathbf{k_{D_P}}}_{\mathbf{k_1} \ldots \mathbf{k_{i-1}} \mathbf{k_i} \mathbf{k_{i+1}} \ldots \mathbf{k_{D_P}}}}{\mathbf{\Delta}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}})} \ ,$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^{\mathbf{2}} \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \ ,$$

$$\mathbf{\Delta}(\mathbf{k_1}, \mathbf{k_2}, \cdots, \mathbf{k_R}) = \delta^{\mathbf{k_1 k_2 \cdots k_R}}_{\mathbf{k_1 k_2 \cdots k_R}}$$

$$\mathbf{v_i^\mu}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1 \ldots k_{i-1} \mu k_{i+1} \ldots k_{D_P}}}_{\mathbf{k_1 \ldots k_{i-1} k_i k_{i+1} \ldots k_{D_P}}}}{\mathbf{\Delta}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}})} \ , \qquad \mathbf{v_i \cdot k_j} = \delta_{\mathbf{ij}} \, , \quad \text{for} \quad \mathbf{j \leq D_P}$$

Physical and transverse space

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum

$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1}, \mathbf{p_2}, \ldots, \mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1}, \mathbf{p_2}, \ldots, \mathbf{p_N}; \mathbf{l})}{\mathbf{d_1 d_2 \cdots d_R}}$$



Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical space: vector space spanned by the inflow momenta. D-dimensional vectors can be decomposed to physical space and transverse space components:

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical space: vector space spanned by the inflow momenta. D-dimensional vectors can be decomposed to physical space and transverse space components:

$$\mathbf{D} = \mathbf{D_P} + \mathbf{D_T},\quad \mathbf{D_P} = \min(\mathbf{D},\mathbf{R}-\mathbf{1}),\quad \mathbf{D_T} = \max(\mathbf{0},\mathbf{D}-\mathbf{R}+\mathbf{1})\,,\qquad \mathbf{l} = \mathbf{l_P} + \mathbf{l_T}$$

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}}-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical space: vector space spanned by the inflow momenta. D-dimensional vectors can be decomposed to physical space and transverse space components:

$$\mathbf{D} = \mathbf{D_P} + \mathbf{D_T},\quad \mathbf{D_P} = \min(\mathbf{D},\mathbf{R}-\mathbf{1}),\quad \mathbf{D_T} = \max(\mathbf{0},\mathbf{D}-\mathbf{R}+\mathbf{1})\,,\qquad \mathbf{l} = \mathbf{l_P} + \mathbf{l_T}$$

If R>D, the transverse space is zero dimensional.

Use NV (dual) coordinates in the physical space:

Use NV (dual) coordinates in the physical space:

$$l^\mu = \sum_{i=1}^{D_P} (l \cdot k_i)\, v_i^\mu + \sum_{i=1}^{D_T} (l \cdot n_i)\; n_i^\mu \;\; .$$

$$g^{\mu\nu} = \sum_{i=1}^{D_P} k_i^\mu v_i^\nu + \sum_{i=1}^{D_T} n_i^\mu n_i^\nu$$

Use NV (dual) coordinates in the physical space:

$$\mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{l}\cdot\mathbf{k_i})\,\mathbf{v_i^{\mu}} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\;\mathbf{n_i^{\mu}}\;.$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} \mathbf{k_i^{\mu}}\mathbf{v_i^{\nu}} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} \mathbf{n_i^{\mu}}\mathbf{n_i^{\nu}}$$

$$\mathbf{n_r}\cdot\mathbf{n_s} = \delta_{\mathbf{rs}}\,,\;\mathbf{k_i}\cdot\mathbf{n_r} = \mathbf{0}\,,\;\mathbf{v_i}\cdot\mathbf{n_r} = \mathbf{0},\qquad \mathbf{w}^{\mu\nu} = \sum_{\mathbf{r=1}}^{\mathbf{D_T}} \mathbf{n_r^{\mu}}\mathbf{n_r^{\nu}},$$

Use NV (dual) coordinates in the physical space:

$$\mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{l}\cdot\mathbf{k_i})\,\mathbf{v_i}^{\mu} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\;\mathbf{n_i}^{\mu}\;.$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} \mathbf{k_i}^{\mu}\mathbf{v_i}^{\nu} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} \mathbf{n_i}^{\mu}\mathbf{n_i}^{\nu}$$

$$\mathbf{n_r}\cdot\mathbf{n_s} = \delta_{\mathbf{rs}}\,,\;\mathbf{k_i}\cdot\mathbf{n_r} = \mathbf{0}\,,\;\mathbf{v_i}\cdot\mathbf{n_r} = \mathbf{0},\qquad \mathbf{w}^{\mu\nu} = \sum_{\mathbf{r=1}}^{\mathbf{D_T}} \mathbf{n_r}^{\mu}\mathbf{n_r}^{\nu},$$

$\mathbf{w}_{\mu\nu}$ is the projector operator to transverse space

$$\mathbf{w}_{\mu}{}^{\nu}(\mathbf{k_1},\ldots,\mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1}\cdots\mathbf{k_{D_P}}\,\nu}_{\mathbf{k_1}\ldots\mathbf{k_{D_P}}\,\mu}}{\mathbf{\Delta}(\mathbf{k_1},\ldots,\mathbf{k_{D_P}})}$$

$$\mathbf{w}_{\mu}{}^{\mu} = \mathbf{D_T}\,,\;\mathbf{k_i^{\mu}}\,\mathbf{w}_{\mu\nu} = \mathbf{0}\,,\;\mathbf{w}^{\mu}{}_{\alpha}\mathbf{w}^{\alpha\nu} = \mathbf{w}^{\mu\nu}.$$

Use NV (dual) coordinates in the physical space:

$$\mathbf{l}^\mu = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{l}\cdot\mathbf{k_i})\,\mathbf{v_i}^\mu + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\;\mathbf{n_i}^\mu \;.$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} \mathbf{k_i}^\mu \mathbf{v_i}^\nu + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} \mathbf{n_i}^\mu \mathbf{n_i}^\nu$$

$$\mathbf{n_r}\cdot\mathbf{n_s} = \delta_{\mathbf{rs}}\,,\ \mathbf{k_i}\cdot\mathbf{n_r} = \mathbf{0}\,,\ \mathbf{v_i}\cdot\mathbf{n_r} = \mathbf{0},\qquad \mathbf{w}^{\mu\nu} = \sum_{\mathbf{r=1}}^{\mathbf{D_T}} \mathbf{n_r}^\mu \mathbf{n_r}^\nu,$$

$\mathbf{w}_{\mu\nu}$ is the projector operator to transverse space

$$\mathbf{w}_\mu{}^\nu(\mathbf{k_1},\ldots,\mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1}\cdots\mathbf{k_{D_P}}\nu}_{\mathbf{k_1}\ldots\mathbf{k_{D_P}}\mu}}{\mathbf{\Delta}(\mathbf{k_1},\ldots,\mathbf{k_{D_P}})}$$

$$\mathbf{w}_\mu{}^\mu = \mathbf{D_T}\,,\ \mathbf{k_i}^\mu\,\mathbf{w}_{\mu\nu} = \mathbf{0}\,,\ \mathbf{w}^\mu{}_\alpha \mathbf{w}^{\alpha\nu} = \mathbf{w}^{\mu\nu}.$$

The dual coordinates of the loop momentum vector provide us the reduction at the integrand level:

$$\mathbf{l}\cdot\mathbf{k_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[\mathbf{d_i} - \mathbf{d_{i-1}} - \left(\mathbf{q_i^2} - \mathbf{m_i^2}\right) + \left(\mathbf{q_{i-1}^2} - \mathbf{m_{i-1}^2}\right)\right]$$

$$\mathbf{l}^\mu = \mathbf{V_R}^\mu + \frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{d_i} - \mathbf{d_{i-1}})\,\mathbf{v_i}^\mu + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\,\mathbf{n_i}^\mu \;,$$

$$\mathbf{V_R}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i=1}}^{\mathbf{D_P}} \Big((\mathbf{q_i^2} - \mathbf{m_i^2}) - (\mathbf{q_{i-1}^2} - \mathbf{m_{i-1}^2})\Big)\,\mathbf{v_i}^\mu$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\, \mathcal{I}_{\mathbf{3}} \quad , \quad \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\, \mathcal{I}_{\mathbf{3}} \;\;,\;\; \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2},\, i \in [0,1,2]$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\,\mathcal{I}_{\mathbf{3}} \quad , \quad \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}}, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]\,, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]$$

$$\mathbf{l}^{\mu} = \mathbf{v_1^{\mu}}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^{\mu}}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^{\mu}} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \;\; \mathbf{v_2^{\mu}} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1},\mathbf{2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\mu}}\,\mathbf{q_i^{\nu}} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\nu}}\,\mathbf{q_i^{\mu}}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\,\mathcal{I}_{\mathbf{3}} \quad , \quad \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}}, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]\,, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]$$

$$\mathbf{l}^{\mu} = \mathbf{v_1^{\mu}}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^{\mu}}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^{\mu}} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \;\; \mathbf{v_2^{\mu}} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1},\mathbf{2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\mu}}\mathbf{q_i^{\nu}} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\nu}}\,\mathbf{q_i^{\mu}}$$

$$\mathbf{2}(\mathbf{d_0} + \mathbf{m_0^2}) = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1},\mathbf{2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w})$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;= \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;= \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;\; = \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;= \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

$$\frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}} = \frac{\mathbf{1}}{(\mathbf{4m_0^2 - w^2})} \left\{ \frac{\mathbf{2(l\cdot v_1) - (w\cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l\cdot v_2) - (w\cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4 + (2l - w)\cdot(v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{\mathbf{d_0 d_1 d_2}} = \frac{1}{(4\mathbf{m_0^2} - \mathbf{w^2})} \left\{ \frac{\mathbf{2(l \cdot v_1)} - \mathbf{(w \cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l \cdot v_2)} - \mathbf{(w \cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4} + \mathbf{(2l - w)} \cdot \mathbf{(v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{\mathbf{d_0 d_1 d_2}} = \frac{1}{(4\mathbf{m_0^2} - \mathbf{w^2})} \left\{ \frac{\mathbf{2(l \cdot v_1) - (w \cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l \cdot v_2) - (w \cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4 + (2l - w) \cdot (v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

In the right hand side the terms proportional to $\mathbf{l}_\mu$ depend only on $\mathbf{l}_\mathbf{T}^\mu$ and integrate to zero.

The inflow momenta for the three bubble denominators are different

$$\mathbf{q_2 \cdot v_1 = 0}\,,\ \mathbf{q_1 \cdot v_2 = 0}\,,\ \mathbf{(q_2 - q_1) \cdot (v_1 + v_2) = 0}$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{d_0 d_1 d_2} = \frac{1}{(4m_0^2 - w^2)} \left\{ \frac{2(l \cdot v_1) - (w \cdot v_1)}{d_0 d_2} + \frac{2(l \cdot v_2) - (w \cdot v_2)}{d_0 d_1} - \frac{4 + (2l - w) \cdot (v_1 + v_2)}{d_1 d_2} \right\}.$$

In the right hand side the terms proportional to l_μ depend only on l_T^μ and integrate to zero.

The inflow momenta for the three bubble denominators are different

$$q_2 \cdot v_1 = 0\,,\ q_1 \cdot v_2 = 0\,,\ (q_2 - q_1) \cdot (v_1 + v_2) = 0$$

$$I_{3;012} = \frac{(-1)}{(4m_0^2 - w^2)} \{(w \cdot v_1) I_{2;02} + (w \cdot v_2) I_{2;01} + (4 - w \cdot (v_1 + v_2)) I_{2;12}\}$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{\mathbf{d_0 d_1 d_2}} = \frac{1}{(\mathbf{4m_0^2 - w^2})} \left\{ \frac{\mathbf{2(l \cdot v_1) - (w \cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l \cdot v_2) - (w \cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4 + (2l - w) \cdot (v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

In the right hand side the terms proportional to $\mathbf{l}_\mu$ depend only on $\mathbf{l}_\mathbf{T}^\mu$ and integrate to zero.

The inflow momenta for the three bubble denominators are different

$$\mathbf{q_2 \cdot v_1 = 0\,,\ q_1 \cdot v_2 = 0\,,\ (q_2 - q_1) \cdot (v_1 + v_2) = 0}$$

$$\mathbf{I_{3;012} = \frac{(-1)}{(4m_0^2 - w^2)} \{(w \cdot v_1) I_{2;02} + (w \cdot v_2) I_{2;01} + (4 - w \cdot (v_1 + v_2)) I_{2;12}\}}$$

Exercise:

Any N-point scalar one-loop integrand function, for N > D, where D is the dimensionality of space-time, can be written as a linear combination of the D-point scalar and vector integrand functions.

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^{\mathbf{2}}}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2 - m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l + k})^{\mathbf{2}} - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^{\mathbf{2}} = \mathbf{1}$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^{\mathbf{2}}}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2 - m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l + k})^{\mathbf{2}} - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^{\mathbf{2}} = \mathbf{1}$$

Our aim is to reduce it using VN method.

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^{\mathbf{2}}}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1 = l^2 - m_1^2}\,,\ \mathbf{d_2 = (l + k)^2 - m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^{\mathbf{2}} = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})}, \qquad (\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})},$$

$$(\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

$$\mathbf{l}^\mu = (\mathbf{l} \cdot \mathbf{n})\mathbf{n}^\mu + (\mathbf{l} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}^\mu + \mathbf{n}^\mu_\epsilon (\mathbf{l} \cdot \mathbf{n}_\epsilon)$$

$$\mathbf{l}^2_{(\mathbf{2})} = \mathbf{d_1} + \mathbf{m_1^2} - \mu^2, \quad \mathbf{2\, l \cdot k} = \mathbf{d_2} - \mathbf{d_1} - \mathbf{r_1^2},$$

$$\mu^2 = (\mathbf{n}_\epsilon \cdot \mathbf{l})^2$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})},$$

$$(\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

$$\mathbf{l}^\mu = (\mathbf{l} \cdot \mathbf{n})\mathbf{n}^\mu + (\mathbf{l} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}^\mu + \mathbf{n}^\mu_\epsilon (\mathbf{l} \cdot \mathbf{n}_\epsilon)$$

$$\mathbf{l}^2_{(\mathbf{2})} = \mathbf{d_1} + \mathbf{m_1^2} - \mu^2, \quad \mathbf{2\, l \cdot k} = \mathbf{d_2} - \mathbf{d_1} - \mathbf{r_1^2},$$

$$\mu^2 = (\mathbf{n}_\epsilon \cdot \mathbf{l})^2$$

Result of the reduction:

$$\frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} = -\frac{(\lambda^2 + \mu^2)}{\mathbf{d_1 d_2}} + \frac{\mathbf{1}}{\mathbf{4k^2}} \left[\frac{\mathbf{r_1^2} - \mathbf{2\, l \cdot k}}{\mathbf{d_1}} + \frac{\mathbf{r_2^2} + \mathbf{2\, l \cdot k} + \mathbf{2k^2}}{\mathbf{d_2}} \right].$$

$$\mathbf{r_1^2} = \mathbf{k^2} + \mathbf{m_1^2} - \mathbf{m_2^2}, \quad \mathbf{r_2^2} = \mathbf{k^2} + \mathbf{m_2^2} - \mathbf{m_1^2},$$

$$\lambda^2 = \frac{\mathbf{k^4} - \mathbf{2k^2}(\mathbf{m_1^2} + \mathbf{m_2^2}) + (\mathbf{m_1^2} - \mathbf{m_2^2})^2}{\mathbf{4k^2}}$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})},$$

$$(\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

$$\mathbf{l}^\mu = (\mathbf{l} \cdot \mathbf{n})\mathbf{n}^\mu + (\mathbf{l} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}^\mu + \mathbf{n}^\mu_\epsilon (\mathbf{l} \cdot \mathbf{n}_\epsilon)$$

$$\mathbf{l}^2_{(\mathbf{2})} = \mathbf{d_1} + \mathbf{m_1^2} - \mu^2, \quad \mathbf{2\, l \cdot k} = \mathbf{d_2} - \mathbf{d_1} - \mathbf{r_1^2},$$

$$\mu^2 = (\mathbf{n}_\epsilon \cdot \mathbf{l})^2$$

Result of the reduction:

$$\frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} = -\frac{(\lambda^2 + \mu^2)}{\mathbf{d_1 d_2}} + \frac{\mathbf{1}}{\mathbf{4k^2}} \left[\frac{\mathbf{r_1^2} - \mathbf{2\, l \cdot k}}{\mathbf{d_1}} + \frac{\mathbf{r_2^2} + \mathbf{2\, l \cdot k} + \mathbf{2k^2}}{\mathbf{d_2}} \right].$$

$$\mathbf{r_1^2} = \mathbf{k^2} + \mathbf{m_1^2} - \mathbf{m_2^2}, \quad \mathbf{r_2^2} = \mathbf{k^2} + \mathbf{m_2^2} - \mathbf{m_1^2},$$

$$\lambda^2 = \frac{\mathbf{k^4} - \mathbf{2k^2}(\mathbf{m_1^2} + \mathbf{m_2^2}) + (\mathbf{m_1^2} - \mathbf{m_2^2})^2}{\mathbf{4k^2}}$$

It follows the parametrization:

$$\frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} = \frac{\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}} \cdot \mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} + \frac{\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n} \cdot \mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}} \cdot \mathbf{l})}{\mathbf{d_1}} + \frac{\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n} \cdot \mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}} \cdot \mathbf{l})}{\mathbf{d_2}}.$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^{\mathbf{2}} = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^{\mathbf{2}}\right] + \left[\mathbf{a}_{1,0} + \mathbf{a}_{1,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{1,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a}_{2,0} + \mathbf{a}_{2,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{2,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a}_{1,0} + \mathbf{a}_{1,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{1,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a}_{2,0} + \mathbf{a}_{2,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{2,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1} + \mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1} + \mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1}+\mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

$$\frac{(\hat{\mathbf{n}}\cdot\mathbf{l_1})^\mathbf{2}}{\mathbf{d_2(l_1)}} - \frac{\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l_1})^\mathbf{2}}{\mathbf{d_2(l_1)}} = \mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \frac{\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})}{\mathbf{d_2(l_1)}}\mathbf{d_1(l_1)}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1}+\mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

$$\frac{(\hat{\mathbf{n}}\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} - \frac{\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} = \mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \frac{\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})}{\mathbf{d_2(l_1)}}\mathbf{d_1(l_1)}$$

$$\mathbf{l_1} = \gamma_\mathbf{1}\mathbf{n} + \gamma_\mathbf{2}\hat{\mathbf{n}}\,, \quad \gamma_\mathbf{1}^\mathbf{2} + \gamma_\mathbf{2}^\mathbf{2} = \mathbf{m_1^2}$$

Choose $\gamma_\mathbf{1} = \mathbf{0}, \gamma_\mathbf{2} = \pm\mathbf{m_1}$

Choose $\gamma_\mathbf{2} = \mathbf{0}, \gamma_\mathbf{1} = \mathbf{m_1}$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1}+\mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

$$\frac{(\hat{\mathbf{n}}\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} - \frac{\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} = \mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \frac{\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})}{\mathbf{d_2(l_1)}}\mathbf{d_1(l_1)}$$

$$\mathbf{l_1} = \gamma_\mathbf{1}\mathbf{n} + \gamma_\mathbf{2}\hat{\mathbf{n}}\,, \quad \gamma_\mathbf{1}^\mathbf{2} + \gamma_\mathbf{2}^\mathbf{2} = \mathbf{m_1^2}$$

$$\mathbf{a_{1,0}} \pm \mathbf{a_{1,2}}\mathbf{m_1} = \frac{\mathbf{m_1^2}+\lambda^\mathbf{2}}{\mathbf{r_1^2}} = \frac{\mathbf{r_1^2}}{\mathbf{4k^2}}$$

Choose $\gamma_\mathbf{1} = \mathbf{0}, \gamma_\mathbf{2} = \pm\mathbf{m_1}$

$$\mathbf{a_{1,2}} = \mathbf{0}\,, \mathbf{a_{1,0}} = \mathbf{r_1^2}/(\mathbf{4k^2})\,,$$

Choose $\gamma_\mathbf{2} = \mathbf{0}, \gamma_\mathbf{1} = \mathbf{m_1}$

$$\mathbf{a_{1,1}} = -(\mathbf{4k^2})^{\mathbf{-1/2}}$$

OPP reduction *(Ossola, Papadopoulos, Pittau)*

First we made direct NV reduction of the loop integrand.

Next we have pointed out a generic parametric form of the loop integrand and fitted the parameters with the help of on-shell values of the loop momenta solving first double cut and single cut conditions (iteratively).

The loop integration can be easily carried out:

$$\mathbf{I(k, m_1^2, m_2^2)} = \int \frac{\mathbf{d^{2-2\epsilon}}}{\mathbf{i(2\pi)^{2-2\epsilon}}} \mathcal{I}(\mathbf{k, m_1, m_2}) = \mathbf{b_0\, I_2(k^2, m_1^2, m_2^2) + a_{1,0}\, I_1(m_1^2) + a_{2,0}\, I_1(m_2^2) + \frac{b_2}{4\pi}}$$

Exercise:

Calculate the photon self-energy in D=2 QED (Scwinger-model) using NV reduction at the integrand level.

OPP reduction: general case

OPP reduction: general case

✦ **Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.**

OPP reduction: general case

- Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.

- "Integrand of a diagram has fully ordered external legs:
  Ordered amplitudes given as sums of diagrams.
  N different l-dependent scalar propagators. Momentum inflow to the loop.

  This gives unique prescription of the integrand function as a function of the loop momentum modulo overall shift.

OPP reduction: general case

✦ Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.

✦ "Integrand of a diagram has fully ordered external legs:
Ordered amplitudes given as sums of diagrams.
N different l-dependent scalar propagators. Momentum inflow to the loop.

This gives unique prescription of the integrand function as a function of the loop momentum modulo overall shift.



OPP reduction: general case

✦ Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.

✦ "Integrand of a diagram has fully ordered external legs:
Ordered amplitudes given as sums of diagrams.
N different l-dependent scalar propagators. Momentum inflow to the loop.

This gives unique prescription of the integrand function as a function of the loop momentum modulo overall shift.

✦ For 4D external kinematics, the integrand of any one-loop Feynman amplitude with arbitrary number of external legs can always be written in linear combination of penta, quadru-,triple-,double- and single-pole terms



Ordered amplitudes have well defined integrand

$$\mathcal{I}_N(p_1,p_2,\ldots,p_N,l)=\frac{\mathcal{N}(p_1,p_2,\ldots,p_N;l)}{d_1 d_2\cdots d_N}$$



$$\mathcal{I}_N(p_1,p_2,\ldots,p_N,l)$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms





The number of terms with k denominators is $\binom{\mathbf{N}}{\mathbf{k}}$







The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d^D l}}{(2\pi)^{\mathrm{D}}} \frac{\mathrm{Num(l)}}{\prod_{\mathrm{i}} \mathrm{d_i(l)}} = \int \frac{\mathrm{d^D l}}{(2\pi)^{\mathrm{D}}} \frac{1}{\prod_{\mathrm{i}} \mathrm{d_i(l)}} \times \Bigg\{ \sum_{\mathrm{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathrm{e}}_{\mathrm{i_1,i_2,i_3,i_4,i_5}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2,i_3,i_4,i_5}]} \mathrm{d_j(l)}$$

$$+ \sum_{\mathrm{i_1,i_2,i_3,i_4}} \tilde{\mathrm{d}}_{\mathrm{i_1,i_2,i_3,i_4}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2,i_3,i_4}]} \mathrm{d_j(l)}$$

$$+ \sum_{\mathrm{i_1,i_2,i_3}} \tilde{\mathrm{c}}_{\mathrm{i_1,i_2,i_3}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2,i_3}]} \mathrm{d_j(l)} + \sum_{\mathrm{i_1,i_2}} \tilde{\mathrm{b}}_{\mathrm{i_1,i_2}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2}]} \mathrm{d_j(l)} + \sum_{\mathrm{i_1}} \tilde{\mathrm{a}}_{\mathrm{i_1}}(\mathrm{l}) \prod_{\mathrm{j} \neq \mathrm{i_1}} \mathrm{d_j(l)} \Bigg\}.$$

$$\mathbf{d_i(l) = (l + q_i)^2 - m_i^2 \, , i = 0, \ldots, 4 \, , \; q_0 = 0}$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathbf{1}}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} \times \Bigg\{ \sum_{\mathbf{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathbf{e}}_{\mathbf{i_1,i_2,i_3,i_4,i_5}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4,i_5]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3,i_4}} \tilde{\mathbf{d}}_{\mathbf{i_1,i_2,i_3,i_4}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3}} \tilde{\mathbf{c}}_{\mathbf{i_1,i_2,i_3}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1,i_2}} \tilde{\mathbf{b}}_{\mathbf{i_1,i_2}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}_{\mathbf{i_1}}(\mathbf{l}) \prod_{\mathbf{j \neq i_1}} \mathbf{d_j}(\mathbf{l}) \Bigg\}.$$

$$\mathbf{d_i}(\mathbf{l}) = (\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}\,, \mathbf{i} = \mathbf{0}, \ldots, \mathbf{4}\,, \;\; \mathbf{q_0} = \mathbf{0}$$

$$\mathrm{Num}(\mathbf{l}) = \mathbf{N_5}(\mathbf{l}) = \prod_{\mathbf{i=1}}^{\mathbf{5}} \mathbf{u_i} \cdot \mathbf{l}, \qquad \mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{4}} (\mathbf{l} \cdot \mathbf{q_i}) \mathbf{v_i^{\mu}} + (\mathbf{l} \cdot \mathbf{n}_{\epsilon}) \mathbf{n}_{\epsilon}^{\mu} \quad \mathbf{l} \cdot \mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left(\mathbf{d_i} - \mathbf{d_0} - (\mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}) \right)$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathbf{1}}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} \times \Bigg\{ \sum_{\mathbf{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathbf{e}}_{\mathbf{i_1,i_2,i_3,i_4,i_5}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4,i_5]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3,i_4}} \tilde{\mathbf{d}}_{\mathbf{i_1,i_2,i_3,i_4}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3}} \tilde{\mathbf{c}}_{\mathbf{i_1,i_2,i_3}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1,i_2}} \tilde{\mathbf{b}}_{\mathbf{i_1,i_2}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}_{\mathbf{i_1}}(\mathbf{l}) \prod_{\mathbf{j \neq i_1}} \mathbf{d_j}(\mathbf{l}) \Bigg\}.$$

$$\mathbf{d_i}(\mathbf{l}) = (\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}\,, \mathbf{i} = \mathbf{0}, \ldots, \mathbf{4}\,, \;\; \mathbf{q_0} = \mathbf{0}$$

$$\mathrm{Num}(\mathbf{l}) = \mathbf{N_5}(\mathbf{l}) = \prod_{\mathbf{i=1}}^{\mathbf{5}} \mathbf{u_i} \cdot \mathbf{l}, \qquad \mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{4}} (\mathbf{l} \cdot \mathbf{q_i}) \mathbf{v_i^{\mu}} + (\mathbf{l} \cdot \mathbf{n}_{\epsilon}) \mathbf{n}_{\epsilon}^{\mu} \quad \mathbf{l} \cdot \mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left(\mathbf{d_i} - \mathbf{d_0} - (\mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}) \right)$$

$$\mathbf{N_5}(\mathbf{l}) = \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{u_5} \cdot \mathbf{l}) = \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{d_j} - \mathbf{d_0}) \; - \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{q_j^2} - \mathbf{m_j^2} + \mathbf{m_0^2})$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathbf{1}}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} \times \Bigg\{ \sum_{\mathbf{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathbf{e}}_{\mathbf{i_1,i_2,i_3,i_4,i_5}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4,i_5]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3,i_4}} \tilde{\mathbf{d}}_{\mathbf{i_1,i_2,i_3,i_4}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3}} \tilde{\mathbf{c}}_{\mathbf{i_1,i_2,i_3}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1,i_2}} \tilde{\mathbf{b}}_{\mathbf{i_1,i_2}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}_{\mathbf{i_1}}(\mathbf{l}) \prod_{\mathbf{j \neq i_1}} \mathbf{d_j}(\mathbf{l}) \Bigg\}.$$

$$\mathbf{d_i}(\mathbf{l}) = (\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}\,, \mathbf{i} = \mathbf{0}, \ldots, \mathbf{4}\,, \;\; \mathbf{q_0} = \mathbf{0}$$

$$\mathrm{Num}(\mathbf{l}) = \mathbf{N_5}(\mathbf{l}) = \prod_{\mathbf{i=1}}^{\mathbf{5}} \mathbf{u_i} \cdot \mathbf{l}, \qquad \mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{4}} (\mathbf{l} \cdot \mathbf{q_i}) \mathbf{v_i^{\mu}} + (\mathbf{l} \cdot \mathbf{n}_{\epsilon}) \mathbf{n}_{\epsilon}^{\mu} \quad \mathbf{l} \cdot \mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left(\mathbf{d_i} - \mathbf{d_0} - (\mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}) \right)$$

$$\mathbf{N_5}(\mathbf{l}) = \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{u_5} \cdot \mathbf{l}) = \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{d_j} - \mathbf{d_0}) \; - \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{q_j^2} - \mathbf{m_j^2} + \mathbf{m_0^2})$$

$$\tilde{\mathbf{e}}_{\mathbf{01234}}(\mathbf{l}) = \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{01234}}$$

Parameter counting

The numerators are simple polynomials of the loop momentum components of the corresponding trivial space.

For a given 5,-4-,3-,2-,1 denominator 1, 5, 10, 10, 5 parameters;

Pentagon (rank five): $\mathbf{D_P = 4\,, D_T = 0 + 1\,, l_T^2 = (ln_\epsilon)^2} = \text{constant terms} + \mathcal{O}(\mathbf{d_i})$

Box (rank four): $\mathbf{D_P = 3\,, D_T = 1 + 1\,, l_T^2 = (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Triangle (rank three): $\mathbf{D_P = 2\,, D_T = 3 + 1\,, l_T^2 = (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Bubble (rank two): $\mathbf{D_P = 1\,, D_T = 3 + 1\,, l_T^2 = (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Tadpole (rank one): $\mathbf{D_P = 0\,, D_T = 4 + 1\,, l_T^2 = (ln_1)^2 + (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Parameter counting

The numerators are simple polynomials of the loop momentum components of the corresponding trivial space.

For a given 5,-4-,3-,2-,1 denominator 1, 5, 10, 10, 5 parameters; $l_T^2 = l^2 - l_P^2\,, \quad l_P^\mu = (lq_i)v_i^\mu$

Pentagon (rank five): $\mathbf{D_P = 4\,, D_T = 0+1\,, l_T^2 = (ln_\epsilon)^2} = \text{constant terms} + \mathcal{O}(\mathbf{d_i})$

Box (rank four): $\mathbf{D_P = 3\,, D_T = 1+1\,, l_T^2 = (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Triangle (rank three): $\mathbf{D_P = 2\,, D_T = 3+1\,, l_T^2 = (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Bubble (rank two): $\mathbf{D_P = 1\,, D_T = 3+1\,, l_T^2 = (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Tadpole (rank one): $\mathbf{D_P = 0\,, D_T = 4+1\,, l_T^2 = (ln_1)^2 + (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Parameter counting

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

Parameter counting

$$\tilde{\mathbf{e}}_{\mathbf{01234}}(\mathbf{l}) = \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{01234}}$$

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathbf{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

Parameter counting

$$\tilde{\mathbf{e}}_{\mathbf{01234}}(\mathbf{l}) = \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{01234}}$$

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathbf{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\begin{aligned}\tilde{\mathbf{c}}_{\mathbf{012}}(\mathbf{l}) = \tilde{\mathbf{c}}_{\mathbf{0}} &+ \tilde{\mathbf{c}}_{\mathbf{1}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}}) + \tilde{\mathbf{c}}_{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{c}}_{\mathbf{3}}((\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}})^{\mathbf{2}} - (\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}})^{\mathbf{2}}) + \tilde{\mathbf{c}}_{\mathbf{4}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}})(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{c}}_{\mathbf{5}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}})^{\mathbf{3}} \\ &+ \tilde{\mathbf{c}}_{\mathbf{6}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}})^{\mathbf{3}} + \tilde{\mathbf{c}}_{\mathbf{7}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{c}}_{\mathbf{8}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}}) + \tilde{\mathbf{c}}_{\mathbf{9}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}})\end{aligned}$$

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

$$\tilde{d}_{0123}(l) = \tilde{d}_0 + \tilde{d}_1(l\cdot n_4) + \tilde{d}_2(l\cdot n_\epsilon)^2 + \tilde{d}_3(l\cdot n_\epsilon)^2(l\cdot n_4) + \tilde{d}_4(l\cdot n_\epsilon)^4,$$

$$\begin{aligned}\tilde{c}_{012}(l) = \tilde{c}_0 &+ \tilde{c}_1(l\cdot n_3) + \tilde{c}_2(l\cdot n_4) + \tilde{c}_3((l\cdot n_3)^2 - (l\cdot n_4)^2) + \tilde{c}_4(l\cdot n_3)(l\cdot n_4) + \tilde{c}_5(l\cdot n_3)^3 \\ &+ \tilde{c}_6(l\cdot n_4)^3 + \tilde{c}_7(l\cdot n_\epsilon)^2 + \tilde{c}_8(l\cdot n_\epsilon)^2(l\cdot n_3) + \tilde{c}_9(l\cdot n_\epsilon)^2(l\cdot n_4)\end{aligned}$$

$$\begin{aligned}\tilde{b}_{01}(l) = \tilde{b}_0 &+ \tilde{b}_1(l\cdot n_2) + \tilde{b}_2(l\cdot n_3) + \tilde{b}_3(l\cdot n_4) + \tilde{b}_4((l\cdot n_2)^2 - (l\cdot n_4)^2) + \tilde{b}_5((l\cdot n_3)^2 - (l\cdot n_4)^2) \\ &+ \tilde{b}_6(l\cdot n_2)(l\cdot n_3) + \tilde{b}_7(l\cdot n_3)(l\cdot n_4) + \tilde{b}_8(l\cdot n_2)(l\cdot n_4) + \tilde{b}_9(l\cdot n_\epsilon)^2,\end{aligned}$$

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

$$\tilde{d}_{0123}(l) = \tilde{d}_0 + \tilde{d}_1(l\cdot n_4) + \tilde{d}_2(l\cdot n_\epsilon)^2 + \tilde{d}_3(l\cdot n_\epsilon)^2(l\cdot n_4) + \tilde{d}_4(l\cdot n_\epsilon)^4,$$

$$\begin{aligned}\tilde{c}_{012}(l) = \tilde{c}_0 &+ \tilde{c}_1(l\cdot n_3) + \tilde{c}_2(l\cdot n_4) + \tilde{c}_3((l\cdot n_3)^2 - (l\cdot n_4)^2) + \tilde{c}_4(l\cdot n_3)(l\cdot n_4) + \tilde{c}_5(l\cdot n_3)^3 \\ &+ \tilde{c}_6(l\cdot n_4)^3 + \tilde{c}_7(l\cdot n_\epsilon)^2 + \tilde{c}_8(l\cdot n_\epsilon)^2(l\cdot n_3) + \tilde{c}_9(l\cdot n_\epsilon)^2(l\cdot n_4)\end{aligned}$$

$$\begin{aligned}\tilde{b}_{01}(l) = \tilde{b}_0 &+ \tilde{b}_1(l\cdot n_2) + \tilde{b}_2(l\cdot n_3) + \tilde{b}_3(l\cdot n_4) + \tilde{b}_4((l\cdot n_2)^2 - (l\cdot n_4)^2) + \tilde{b}_5((l\cdot n_3)^2 - (l\cdot n_4)^2) \\ &+ \tilde{b}_6(l\cdot n_2)(l\cdot n_3) + \tilde{b}_7(l\cdot n_3)(l\cdot n_4) + \tilde{b}_8(l\cdot n_2)(l\cdot n_4) + \tilde{b}_9(l\cdot n_\epsilon)^2,\end{aligned}$$

$$\tilde{a}_i(l) = \tilde{a}_0 + \tilde{a}_1(l\cdot n_1) + \tilde{a}_2(l\cdot n_2) + \tilde{a}_3(l\cdot n_3) + \tilde{a}_4(l\cdot n_4)$$

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

$$\tilde{d}_{0123}(l) = \tilde{d}_0 + \tilde{d}_1(l\cdot n_4) + \tilde{d}_2(l\cdot n_\epsilon)^2 + \tilde{d}_3(l\cdot n_\epsilon)^2(l\cdot n_4) + \tilde{d}_4(l\cdot n_\epsilon)^4,$$

$$\begin{aligned}\tilde{c}_{012}(l) = \tilde{c}_0 &+ \tilde{c}_1(l\cdot n_3) + \tilde{c}_2(l\cdot n_4) + \tilde{c}_3((l\cdot n_3)^2 - (l\cdot n_4)^2) + \tilde{c}_4(l\cdot n_3)(l\cdot n_4) + \tilde{c}_5(l\cdot n_3)^3 \\ &+ \tilde{c}_6(l\cdot n_4)^3 + \tilde{c}_7(l\cdot n_\epsilon)^2 + \tilde{c}_8(l\cdot n_\epsilon)^2(l\cdot n_3) + \tilde{c}_9(l\cdot n_\epsilon)^2(l\cdot n_4)\end{aligned}$$

$$\begin{aligned}\tilde{b}_{01}(l) = \tilde{b}_0 &+ \tilde{b}_1(l\cdot n_2) + \tilde{b}_2(l\cdot n_3) + \tilde{b}_3(l\cdot n_4) + \tilde{b}_4((l\cdot n_2)^2 - (l\cdot n_4)^2) + \tilde{b}_5((l\cdot n_3)^2 - (l\cdot n_4)^2) \\ &+ \tilde{b}_6(l\cdot n_2)(l\cdot n_3) + \tilde{b}_7(l\cdot n_3)(l\cdot n_4) + \tilde{b}_8(l\cdot n_2)(l\cdot n_4) + \tilde{b}_9(l\cdot n_\epsilon)^2,\end{aligned}$$

$$\tilde{a}_i(l) = \tilde{a}_0 + \tilde{a}_1(l\cdot n_1) + \tilde{a}_2(l\cdot n_2) + \tilde{a}_3(l\cdot n_3) + \tilde{a}_4(l\cdot n_4)$$

The coefficients $\tilde{a}_0, \ldots, \tilde{e}_{01234}$ are independent from the loop momenta and in all numerator functions we can replace $(l\cdot n_i)$ with $(l_T\cdot n_i)$

Note that the integration over the transverse space is trivial

$$\int d^{D_1}l_\perp \delta(l_\perp^2 - \mu_0^2)\,(l_\perp^\mu, l_\perp^\mu l_\perp^\nu) = \int d^{D_1}l_\perp \delta(l_\perp^2 - \mu_0^2)\left(0,\ \frac{g_\perp^{\mu\nu}}{D_1}\, l_\perp^2\right), \qquad D_1 = D - 1$$

Only the constant terms and some of the terms depending on $(\mathbf{l} \cdot \mathbf{n}_\epsilon)^2$ give non-vanishing integrals

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}\mathbf{i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

$$\mathcal{R} = -\sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \frac{\tilde{\mathbf{d}}^{(\mathbf{4})}_{\mathbf{i_1 i_2 i_3 i_4}}}{\mathbf{6}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \frac{\tilde{\mathbf{c}}^{(\mathbf{7})}_{\mathbf{i_1 i_2 i_3}}}{\mathbf{2}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \left[\frac{\mathbf{m^2_{i_1}} + \mathbf{m^2_{i_2}}}{\mathbf{2}} - \frac{(\mathbf{q_{i_1}} - \mathbf{q_{i_2}})^{\mathbf{2}}}{\mathbf{6}} \right] \tilde{\mathbf{b}}^{(\mathbf{9})}_{\mathbf{i_1 i_2}}.$$

Example: Projecting out individual (quadrupole) coefficients in D-dimension

$$\mathcal{I}_N(p_1,p_2,\ldots,p_N,l) = \frac{\mathcal{N}(p_1,p_2,\ldots,p_N;l)}{d_1 d_2 \cdots d_N} =$$

$$= \sum_{i_1\le i_2\le i_3\le i_4\le i_5\le n} \frac{\bar{e}_{i_1 i_2 i_3 i_4 i_5}(l)}{d_{i_1} d_{i_2} d_{i_3} d_{i_4} d_{i_5}} +$$

$$\sum_{1\le i_1<i_2<i_3<i_4\le N} \frac{\bar{d}_{i_1 i_2 i_3 i_4}(l)}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} + \sum_{1\le i_1<i_2<i_3\le N} \frac{\bar{c}_{i_1 i_2 i_3}(l)}{d_{i_1} d_{i_2} d_{i_3}} + \sum_{1\le i_1<i_2\le N} \frac{\bar{b}_{i_1 i_2}(l)}{d_{i_1} d_{i_2}} + \sum_{1\le i_1\le N} \frac{\bar{a}_{i_1}(l)}{d_{i_1}}$$

Denote:

$$\mathbf{RR(l) = Residuum_{0123}(Integrand)} \\ \mathbf{- \text{ pentagon contributions } = \tilde{d}_{0123}(l)}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathrm{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathrm{n}_\epsilon)^2 + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathrm{n}_\epsilon)^2(\mathrm{l}\cdot\mathrm{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathrm{n}_\epsilon)^4,$$

$$\mathrm{l}^\mu = \mathrm{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathrm{n}_4^\mu + \sin\phi\ \mathrm{n}_\epsilon^\mu), \qquad \mathrm{V}^\mu = -\frac{1}{2}\sum_{\mathrm{i}}^{3} \mathrm{v}_\mathrm{i}^\mu \left(\mathrm{q}_\mathrm{i}^2 - \mathrm{m}_\mathrm{i}^2 + \mathrm{m}_0^2\right),$$

- Choose $\sin\phi = 0, \cos\phi = \pm 1$ denote $\mathrm{l}_\pm^\mu = \mathrm{V}^\mu \pm \mathrm{l}_\perp \mathrm{n}_4^\mu,$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

- Choose $\mathrm{l}_{\epsilon}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

- Choose $\mathrm{l}_{\epsilon}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\mathrm{l}_{\perp}^{\mathbf{2}} = \frac{\mathrm{Num}(\mathrm{l}_{\epsilon}) - \tilde{\mathbf{d}}_{\mathbf{0}}}{\mathrm{l}_{\perp}^{\mathbf{2}}}.$$

Comments on the rational part:

The origin is UV divergent tensor integrals. Reduction requires regularization.
After reduction: D-dimensional finite tensor integrals. In the limiting case D=4
they provide finite constants independent from the kinematics.

It may happen that the numerator has manifest dependence on (D-4) coming from polarization sum. These terms can only contribute if it appears in a term leading to UV divergent integrals. There are only few UV divergent one-loop Feynman diagrams even for large number of external particles (OPP).

Sophisticated recursion relations (BDK) as an option in Black Hat

Comment on N=4 sYM amplitudes

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

- N=1 sYM scattering amplitudes have no rational parts

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

- N=1 sYM scattering amplitudes have no rational parts

BDDK theorem (1994) for one loop amplitudes:

The maximum number of loop momentum in the numerator of Feynman-diagrams is reduced by one for N=1 and by four for N=4 .

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

- N=1 sYM scattering amplitudes have no rational parts

BDDK theorem (1994) for one loop amplitudes:

The maximum number of loop momentum in the numerator of Feynman-diagrams is reduced by one for N=1 and by four for N=4 .

Excercise (not easy):

Find proper parametrization for the bubble numerator $\tilde{\mathrm{b}}_{\mathbf{01}}(\mathrm{l})$ in case of light-like inflow momentum.

Lecture 3: Unitarity method and amplitudes

Constraints from Unitarity: $M^\dagger - M = -iM^\dagger M$



$$-i\,\mathrm{Disc}\;A_4(1,2,3,4)\Big|_{s-\mathrm{cut}} = \int \frac{d^4p}{(2\pi)^4}\, 2\pi\delta^{(+)}(\ell_1^2 - m^2)\, 2\pi\delta^{(+)}(\ell_2^2 - m^2)$$
$$\times\, A_4^{\mathrm{tree}}(-\ell_1, 1, 2, \ell_2)\, A_4^{\mathrm{tree}}(-\ell_2, 3, 4, \ell_1)\,,$$

Imaginary part of NLO an amplitude is calculated from tree amplitudes.

- Non-linear relation, iterative in the coupling.
- Iterative in amplitudes. Building blocks are amplitudes and not Feynman diagrams
- Manifestly gauge invariant.

Unitarity and Cutkosky rule:

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s = (p_1 + p_2)^2, t = (p_1 + p_3)^2,}$

$$\mathbf{s = 4E^2_{cm}, \quad t = -\frac{(s - 4m^2)}{2}\,(1 - \cos\theta_{cms})}$$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity:

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,}$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^* \langle n|T|p_1,p_2\rangle}$$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed θ_{cms} ?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^* \langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed θ_{cms} ?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,}$ $\qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

More external particles?

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

More external particles?

In perturbation theory simple recipe to calculate discontinuities across branch cuts of multi-leg, multi-loop Feynman diagrams:

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

More external particles?

In perturbation theory simple recipe to calculate discontinuities across branch cuts of multi-leg, multi-loop Feynman diagrams:

Cutkosky rule: $\mathbf{1/(p^2-m^2+i\delta)\to(-2\pi i)\delta(p^2-m^2)\theta(p_0)}$

Cutkosky rule in case of the bubble integral:

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2}, \qquad D_1 = l^2 - m^2 + i0, \quad D_2 = (p-l)^2 - m^2 + i0$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2}, \qquad D_1 = l^2 - m^2 + i0, \quad D_2 = (p-l)^2 - m^2 + i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s = p^2$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2}, \qquad D_1 = l^2 - m^2 + i0, \quad D_2 = (p-l)^2 - m^2 + i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s = p^2$

$$\epsilon_l = \sqrt{\vec{l}^2 + m^2}$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$$D_1:\qquad a_1)\ l_0=\epsilon_l-i0,\qquad b_1)\ l_0=-\epsilon_l+i0,$$

$$D_2:\qquad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$$D_1:\qquad a_1)\ l_0=\epsilon_l-i0,\qquad\qquad b_1)\ l_0=-\epsilon_l+i0,$$

$$D_2:\qquad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$$



Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$$D_1:\qquad a_1)\ l_0=\epsilon_l-i0,\qquad\qquad b_1)\ l_0=-\epsilon_l+i0,$$

$$D_2:\qquad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$$

No discontinuity in p_0 if the poles a_2,b_2 are far from the contour



Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$D_1:\quad a_1)\ l_0=\epsilon_l-i0,\qquad b_1)\ l_0=-\epsilon_l+i0,$

$D_2:\quad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$

No discontinuity in p_0 if the poles a_2,b_2 are far from the contour

The distance of the poles a_1,b_2 can vanish if $p_0>2m$ when these poles can pinch the contour.



One can avoid pinching the contour by moving the first pole above the contour.
To compensate the difference for the integral we add an integral over a closed small circle around the first pole

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

$$\mathcal{I}^{(2)} = \int \frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\left[\int_{C'}\frac{\mathrm{d}l_0}{(2\pi)}\ldots + \oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\ldots\right]$$

$$\oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)} = \int\frac{\mathrm{d}l_0}{(2\pi)}\frac{(-2\pi i)\delta_+(l^2-m^2)}{((l_0-p_0)^2-\epsilon_l^2)}$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \mathcal{I}_2(p^2+i\delta)-\mathcal{I}_2(p^2-i\delta) = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\times\left[\frac{1}{(l_0-p_0-i\delta)^2-\epsilon_l^2}-\mathrm{c.c}\right]$$

$$= \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\left[\frac{1}{(l-p)^2-m^2+i\delta}-\mathrm{c.c.}\right]$$

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

$$\mathcal{I}^{(2)} = \int \frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\left[\int_{C'}\frac{\mathrm{d}l_0}{(2\pi)}\ldots + \oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\ldots\right]$$

$$\oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)} = \int\frac{\mathrm{d}l_0}{(2\pi)}\frac{(-2\pi i)\delta_+(l^2-m^2)}{((l_0-p_0)^2-\epsilon_l^2)}$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \mathcal{I}_2(p^2+i\delta)-\mathcal{I}_2(p^2-i\delta) = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\times\left[\frac{1}{(l_0-p_0-i\delta)^2-\epsilon_l^2}-\mathrm{c.c}\right]$$

$$= \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\left[\frac{1}{(l-p)^2-m^2+i\delta}-\mathrm{c.c.}\right]$$

$$\frac{1}{x+i\delta} = \mathrm{P}\left[\frac{1}{x}\right] - i\pi\delta(x)$$

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

$$\mathcal{I}^{(2)} = \int \frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\left[\int_{C'}\frac{\mathrm{d}l_0}{(2\pi)}\ldots + \oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\ldots\right]$$

$$\oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)} = \int\frac{\mathrm{d}l_0}{(2\pi)}\frac{(-2\pi i)\delta_+(l^2-m^2)}{((l_0-p_0)^2-\epsilon_l^2)}$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \mathcal{I}_2(p^2+i\delta)-\mathcal{I}_2(p^2-i\delta) = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\times\left[\frac{1}{(l_0-p_0-i\delta)^2-\epsilon_l^2}-\mathrm{c.c}\right]$$

$$= \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\left[\frac{1}{(l-p)^2-m^2+i\delta}-\mathrm{c.c.}\right]$$

$$\frac{1}{x+i\delta} = \mathrm{P}\left[\frac{1}{x}\right] - i\pi\delta(x)$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\;(-2\pi i)\delta_+((l-p)^2-m^2).$$

Cutkosky rule, generalizations from the case of the bubble integral:

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2} = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \int \mathrm{d}\alpha_1 \mathrm{d}\alpha_2 \frac{\delta(1 - \sum \alpha_i)}{D^2}, \qquad D = \alpha_1 D_1 + \alpha_2 D_2$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2} = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \int \mathrm{d}\alpha_1 \mathrm{d}\alpha_2 \frac{\delta(1 - \sum \alpha_i)}{D^2}, \qquad D = \alpha_1 D_1 + \alpha_2 D_2$$

Necessary condition for singularity: $D_1(l^*) = 0\,,\ D_2(l^*) = 0$ and so $D(l^*) = 0$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

$$\mathcal{I}(\{p_i\},\{m_i\})=\int\prod_{k=1}^{L}\frac{d^4l_k}{(2\pi)^4}\frac{\mathcal{N}(\{l_i\}\{p_i\})}{\prod_{i=1}^{N}D_i}\qquad D_i=q_i^2-m_i^2,i\in N$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

$$\mathcal{I}(\{p_i\},\{m_i\})=\int\prod_{k=1}^{L}\frac{d^4l_k}{(2\pi)^4}\,\frac{\mathcal{N}(\{l_i\}\,\{p_i\})}{\prod_{i=1}^{N}D_i}\qquad D_i=q_i^2-m_i^2,i\in N$$

$$D_i=q_i^2-m_i^2=0,\quad (i=1,2,...,r\leq 4L)$$

$$\sum_{i\subset\text{loop j}}\alpha_i^{(j)}q_i=0\quad\text{for every loop.}$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

$$\mathcal{I}(\{p_i\},\{m_i\})=\int\prod_{k=1}^{L}\frac{d^4l_k}{(2\pi)^4}\frac{\mathcal{N}(\{l_i\}\{p_i\})}{\prod_{i=1}^{N}D_i}\qquad D_i=q_i^2-m_i^2, i\in N$$

$$D_i=q_i^2-m_i^2=0,\quad (i=1,2,...,r\leq 4L)$$

$$\sum_{i\subset\text{loop j}}\alpha_i^{(j)}q_i=0\quad\text{for every loop.}$$

Excercise: Work out the Landau equations and solve them for the self energy and two different internal mass and find the location of the branch cut.

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

$$\text{Disc}\,[\mathcal{I}] = \int \prod_{k=1}^{L} \frac{d^4 l_k}{(2\pi)^4} \; \frac{\mathcal{N}(\{l_i\}\,\{p_i\}) \prod_{i=1}^{r} (-2\pi i)\delta^{(+)}(q_i^2 - m_i^2)}{\prod_{j=r+1}^{N} (q_j^2 - m_j^2)}$$

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

$$\mathrm{Disc}\,[\mathcal{I}] = \int \prod_{k=1}^{L} \frac{d^4 l_k}{(2\pi)^4} \; \frac{\mathcal{N}(\{l_i\}\,\{p_i\}) \prod_{i=1}^{r} (-2\pi i)\delta^{(+)}(q_i^2 - m_i^2)}{\prod_{j=r+1}^{N} (q_j^2 - m_j^2)}$$

Unitarity: Non-linear relations between scattering amplitudes.
It can be used to compute the discontinuities of scattering amplitudes at a given order in PT in terms of amplitudes at lower order.

It is built into perturbation theory in even for external off-shell lines and external lines with on-shell complex momenta.
If the Landau equations have solutions, the Cutkosky formula provide the correct singularities.

The diagrammatic version of the Cutkosky formula also can be applied to scattering amplitudes, leading to generalized unitarity relations. E.g. triple cut.

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

$$\text{Disc}\,[\mathcal{I}] = \int \prod_{k=1}^{L} \frac{d^4 l_k}{(2\pi)^4} \; \frac{\mathcal{N}(\{l_i\}\,\{p_i\}) \prod_{i=1}^{r} (-2\pi i)\delta^{(+)}(q_i^2 - m_i^2)}{\prod_{j=r+1}^{N} (q_j^2 - m_j^2)}$$

Unitarity: Non-linear relations between scattering amplitudes.
It can be used to compute the discontinuities of scattering amplitudes at a given order in PT in terms of amplitudes at lower order.

It is built into perturbation theory in even for external off-shell lines and external lines with on-shell complex momenta.
If the Landau equations have solutions, the Cutkosky formula provide the correct singularities.

The diagrammatic version of the Cutkosky formula also can be applied to scattering amplitudes, leading to generalized unitarity relations. E.g. triple cut.

$$\text{Disc}_{l \to l_{n_1 n_2 n_3}} \left[A_n^{1-\text{loop}}\right] = \sum_{\text{states}} \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \prod_{i=1}^{3} (-2\pi i)\delta_+(l_i^2 - m_i^2)$$

$$\times A_{n_1}^{\text{tree}}(-l_3, l_1) A_{n_2}^{\text{tree}}(-l_1, l_2) A_{n_3}^{\text{tree}}(-l_2, l_3),$$

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:



Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.



Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})} + \frac{\mathbf{l}_\mu \mathbf{b}_\nu + \mathbf{b}_\mu \mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}} = \sum_{\mathbf{i=1}}^{\mathbf{D_s-2}} \mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}} \mathbf{l}_\mu \mathbf{\Gamma}^\mu + \mathbf{m} = \sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}} \mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})} + \frac{\mathbf{l}_\mu \mathbf{b}_\nu + \mathbf{b}_\mu \mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}} = \sum_{\mathbf{i=1}}^{\mathbf{D_s-2}} \mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}} \mathbf{l}_\mu \mathbf{\Gamma}^\mu + \mathbf{m} = \sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}} \mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1 n_2 n_3 n_4}}} \left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right] = \sum_{\mathrm{states}}$$

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})}+\frac{\mathbf{l}_\mu\mathbf{b}_\nu+\mathbf{b}_\mu\mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}}=\sum_{\mathbf{i=1}}^{\mathbf{D_s-2}}\mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}}\mathbf{l}_\mu\mathbf{\Gamma}^\mu+\mathbf{m}\;=\sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}}\mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1}}\mathbf{n_2}\mathbf{n_3}\mathbf{n_4}}\;\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right]=\sum_{\mathrm{states}}\;\mathcal{A}_{\mathbf{n_1}}^{\mathrm{tree}}(-\mathbf{l_4},\mathbf{l_1})\mathcal{A}_{\mathbf{n_2}}^{\mathrm{tree}}(-\mathbf{l_1},\mathbf{l_2})\mathcal{A}_{\mathbf{n_3}}^{\mathrm{tree}}(-\mathbf{l_2},\mathbf{l_3})\mathcal{A}_{\mathbf{n_4}}^{\mathrm{tree}}(-\mathbf{l_3},\mathbf{l_4}),$$

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})}+\frac{\mathbf{l}_\mu\mathbf{b}_\nu+\mathbf{b}_\mu\mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}}=\sum_{\mathbf{i=1}}^{\mathbf{D_s-2}}\mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}}\mathbf{l}_\mu\mathbf{\Gamma}^\mu+\mathbf{m}\;=\sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}}\mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1}}\mathbf{n_2}\mathbf{n_3}\mathbf{n_4}}\;\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right]=\sum_{\mathrm{states}}\;\mathcal{A}_{\mathbf{n_1}}^{\mathrm{tree}}(-\mathbf{l_4},\mathbf{l_1})\mathcal{A}_{\mathbf{n_2}}^{\mathrm{tree}}(-\mathbf{l_1},\mathbf{l_2})\mathcal{A}_{\mathbf{n_3}}^{\mathrm{tree}}(-\mathbf{l_2},\mathbf{l_3})\mathcal{A}_{\mathbf{n_4}}^{\mathrm{tree}}(-\mathbf{l_3},\mathbf{l_4}),$$

Key point: two independent expressions for the same cut amplitude

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})} + \frac{\mathbf{l}_\mu \mathbf{b}_\nu + \mathbf{b}_\mu \mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}} = \sum_{\mathbf{i=1}}^{\mathbf{D_s-2}} \mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}} \mathbf{l}_\mu \mathbf{\Gamma}^\mu + \mathbf{m} = \sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}} \mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1 n_2 n_3 n_4}}}\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right] = \sum_{\mathrm{states}} \mathcal{A}_{\mathbf{n_1}}^{\mathrm{tree}}(-\mathbf{l_4},\mathbf{l_1})\mathcal{A}_{\mathbf{n_2}}^{\mathrm{tree}}(-\mathbf{l_1},\mathbf{l_2})\mathcal{A}_{\mathbf{n_3}}^{\mathrm{tree}}(-\mathbf{l_2},\mathbf{l_3})\mathcal{A}_{\mathbf{n_4}}^{\mathrm{tree}}(-\mathbf{l_3},\mathbf{l_4}),$$

Key point: two independent expressions for the same cut amplitude

$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}*=\mathbf{l}_{\mathbf{n_1 n_2 n_3 n_4}}}\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right] = \sum_{\mathbf{n_5}} \frac{\tilde{\mathbf{e}}_{\mathbf{n_1 n_2 n_3 n_4 n_5}}(\mathbf{l}^*)}{\mathbf{d}_{\mathbf{n_5}}(\mathbf{l}^*)} + \tilde{\mathbf{d}}_{\mathbf{n_1 n_2 n_3 n_4}}(\mathbf{l}^*)$$

Projecting out individual quadrupole coefficients in D-dimension:

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^{\mathbf{4}},$$

$$\mathrm{l}^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathbf{n}_4^\mu + \sin\phi\ \mathbf{n}_\epsilon^\mu), \qquad \mathbf{V}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_\mathbf{i}^\mu \left(\mathbf{q}_\mathbf{i}^\mathbf{2} - \mathbf{m}_\mathbf{i}^\mathbf{2} + \mathbf{m}_\mathbf{0}^\mathbf{2}\right),$$

$$\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2 + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^4,$$

$$\mathrm{l}^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathbf{n}_4^\mu + \sin\phi\ \mathbf{n}_\epsilon^\mu), \qquad \mathbf{V}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_\mathbf{i}^\mu \left(\mathbf{q}_\mathbf{i}^\mathbf{2} - \mathbf{m}_\mathbf{i}^\mathbf{2} + \mathbf{m}_\mathbf{0}^\mathbf{2}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_\pm^\mu = \mathbf{V}^\mu \pm \mathrm{l}_\perp \mathbf{n}_4^\mu,$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_\mathbf{0} = \frac{\mathrm{RR}(\mathrm{l}_+) + \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_\mathbf{1} = \frac{\mathrm{RR}(\mathrm{l}_+) - \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}\mathrm{l}_\perp}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

- Choose $\mathrm{l}_{\epsilon}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2 + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^4,$$

$$\mathrm{l}^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathbf{n}_4^\mu + \sin\phi\ \mathbf{n}_\epsilon^\mu), \qquad \mathbf{V}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_\mathbf{i}^\mu \left(\mathbf{q}_\mathbf{i}^\mathbf{2} - \mathbf{m}_\mathbf{i}^\mathbf{2} + \mathbf{m}_\mathbf{0}^\mathbf{2}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_\pm^\mu = \mathbf{V}^\mu \pm \mathrm{l}_\perp \mathbf{n}_4^\mu,$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_\mathbf{0} = \frac{\mathrm{RR}(\mathrm{l}_+) + \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_\mathbf{1} = \frac{\mathrm{RR}(\mathrm{l}_+) - \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}\mathrm{l}_\perp}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_\pm = \mathbf{V} \pm \mathrm{l}_\perp(\mathbf{n}_4 + \mathbf{n}_\epsilon)/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_\mathbf{2} + \tilde{\mathbf{d}}_\mathbf{4}\frac{\mathrm{l}_\perp^\mathbf{2}}{\mathbf{2}} = \mathrm{l}_\perp^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_+) + \mathrm{RR}(\tilde{\mathrm{l}}_-) - \mathbf{2}\,\tilde{\mathbf{d}}_\mathbf{0}\right), \tilde{\mathbf{d}}_\mathbf{3} = \sqrt{\mathbf{2}}\,\mathrm{l}_\perp^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_+) - \mathrm{RR}(\tilde{\mathrm{l}}_-) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_\mathbf{1}\mathrm{l}_\perp\right).$$

- Choose $\mathrm{l}_\epsilon^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp$

$$\tilde{\mathbf{d}}_\mathbf{2} + \tilde{\mathbf{d}}_\mathbf{4}\mathrm{l}_\perp^\mathbf{2} = \frac{\mathrm{Num}(\mathrm{l}_\epsilon) - \tilde{\mathbf{d}}_\mathbf{0}}{\mathrm{l}_\perp^\mathbf{2}}.$$

The parameters are fixed by linear algebraic equations

$$\mathrm{Res}_{ij\cdots k}\left[F(l)\right] \equiv \left[d_i(l)d_j(l)\cdots d_k(l)F(l)\right]\Big]_{l=l_{ij\cdots k}} .$$

$$\bar{d}_{ijkl}(l) = \mathrm{Res}_{ijkl}\left(\mathcal{A}_N(l)\right) \qquad d_i=d_j=d_k=d_l=0 \qquad \text{two solutions}$$

$$\bar{c}_{ijk}(l) = \mathrm{Res}_{ijk}\left(\mathcal{A}_N(l) - \sum_{l\neq i,j,k}\frac{\bar{d}_{ijkl}(l)}{d_i d_j d_k d_l}\right) \qquad d_i=d_j=d_k=0 \qquad \text{infinite \# of solutions}$$

$$\bar{b}_{ij}(l) = \mathrm{Res}_{ij}\left(\mathcal{A}_N(l) - \sum_{k\neq i,j}\frac{\bar{c}_{ijk}(l)}{d_i d_j d_k} - \frac{1}{2!}\sum_{k,l\neq i,j}\frac{\bar{d}_{ijkl}(l)}{d_i d_j d_k d_l}\right) \qquad d_i=d_j=0$$

infinite # of solutions

unitarity: the residues factorize into the products of tree amplitudes

we fully reconstruct the integrand in terms of product of tree amplitudes

no Feynman diagrams

Unitarity method: one-loop amplitudes from tree amplitudes + scalar integral functions

- Decompose the amplitude in terms of basic set of scalar integral functions and read out the coefficients using unitarity cuts (‘98) (BDK)

- Consider the integrand, the amplitude is parametric integral over the loop momentum OPP(‘06) (EGK (’07))

- Use generalized cuts, read out the coefficients in terms of tree amplitudes at cut-momenta (complex) BCF/BDK(’05)

- Rational part is obtained by carrying out the algorithm in two different integer D>4 dimensions GKM (08) (see also Badger,BDK, OPP)

Unitarity method: one-loop amplitudes from tree amplitudes + scalar integral functions

- Decompose the amplitude in terms of basic set of scalar integral functions and read out the coefficients using unitarity cuts (‘98) (BDK)

- Consider the integrand, the amplitude is parametric integral over the loop momentum OPP(‘06) (EGK (’07))

- Use generalized cuts, read out the coefficients in terms of tree amplitudes at cut-momenta (complex) BCF/BDK(’05)

- Rational part is obtained by carrying out the algorithm in two different integer D>4 dimensions GKM (08) (see also Badger,BDK, OPP)

Bern, Dixon, Kosower (BDK) ; Ellis, Giele, ZK, Melnikov (EGKM); Britto, Cachazo, Feng (BCF)

Managing color of amplitudes

Unitarity: instead of Feynman diagrams amplitudes
Tree and one loop Feynman diagrams: color and space time part

$$\mathcal{A}_{\mathbf{n}}^{\mathrm{tree}}(\{\mathbf{k}\},\{\mathbf{h}\},\{\mathbf{a}\}),\ \mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}(\{\mathbf{k}\},\{\mathbf{h}\},\{\mathbf{a}\})$$

➡ Color decomposition of amplitudes with the help of a basis color space
(T-based, F-based, color flow based}

$$[T^a,T^b]=-F^a_{bc}T^c\,,\qquad \mathrm{Tr}(T^aT^b)=\delta_{ab}.$$

$$[F^a,F^b]=-F^a_{bc}F^c\,,\quad F^a_{bc}=-i\sqrt{2}f^{abc}\,,\quad \mathrm{Tr}(F^aF^b)=2N_c\delta_{ab}.$$
$$\left(F^{a_1}\right)_{a_2a_3}=-\frac{1}{2N_c}\mathrm{Tr}\left([F^{a_1},F^{a_2}]F^{a_3}\right)=-\mathrm{Tr}\left([T^{a_1},T^{a_2}]T^{a_3}\right).$$

$$(T^a)^{\bar{j}_1}_{i_1}(T^a)^{\bar{j}_2}_{i_2}=(\delta)^{\bar{j}_2}_{i_1}(\delta)^{\bar{j}_1}_{i_2}-\frac{1}{N_c}(\delta)^{\bar{j}_1}_{i_1}(\delta)^{\bar{j}_2}_{i_2}$$

$$\begin{aligned}\left(F^{a_2}F^{a_3}\ldots F^{a_{(n-2)}}F^{a_{(n-1)}}\right)_{a_1a_n}&=\frac{1}{2N_c}\mathrm{Tr}\left([[\ldots[[F^{a_1},F^{a_2}],F^{a_3}],\ldots,F^{a_{n-2}}]\,[F^{a_{n-1}},F^{a_n}]\right)\\&=\mathrm{Tr}\left([[\ldots[[T^{a_1},T^{a_2}],T^{a_3}],\ldots,T^{a_{n-2}}]\,[T^{a_{n-1}},T^{a_n}]\right).\end{aligned}$$

Color ordered n-gluon tree sub-amplitudes

$$\mathcal{A}_{\mathbf{n}}^{\text{tree}} = \frac{\mathbf{g_s^{n-2}}}{\mathbf{2N_c}} \sum_{\sigma \in \mathbf{S_n/Z_n}} \mathrm{Tr}\left(\mathbf{F^{a_{\sigma(1)}} F^{a_{\sigma(2)}} F^{a_{\sigma(3)}} \dots F^{a_{\sigma(n)}}}\right) \mathbf{A}_{\mathbf{n},\sigma}^{\text{tree}},$$

(n-1)! color ordered sub-amplitudes

$$\mathcal{A}_{\mathbf{n}}^{\text{tree}} = \mathbf{g_s^{n-2}} \sum_{\sigma \in \mathbf{S_n/Z_n}} \mathrm{Tr}\left(\mathbf{T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} T^{a_{\sigma(3)}} \dots T^{a_{\sigma(n)}}}\right) \mathbf{A}_{\mathbf{n},\sigma}^{\text{tree}}$$

$$\mathbf{A}_{\mathbf{n},\sigma}^{\text{tree}} = \mathbf{m_n}\left(\mathbf{g_{\sigma(1)}, g_{\sigma(2)}, g_{\sigma(3)}, \dots, g_{\sigma(n)}}\right)$$

Some properties of sub-amplitudes:

$$\mathbf{m_n}(\mathbf{g_1, g_2, g_3, \dots, g_n}) = \mathbf{m_n}(\mathbf{g_2, g_3, \dots, g_n, g_1})$$ (cyclic identity)

$$\mathbf{m_n}(\mathbf{g_1, g_2, g_3, \dots, g_{n-1}, g_n}) = (-1)^{\mathbf{n}} \mathbf{m_n}(\mathbf{g_n, g_{n-1}, \dots, g_2, g_1}),$$ (reflection identity)

$$\mathbf{m_n}(\mathbf{1}, \underline{\mathbf{2}, \dots \mathbf{n_1}}, \overline{\mathbf{n_1+1}, \dots, \mathbf{n}}) \equiv \sum_{\sigma(\mathbf{n})} \mathbf{m_n}(\mathbf{g_1, g_{\sigma(2)}, g_{\sigma(3)}, \dots, g_{\sigma(n)}}) = \mathbf{0}$$ (Abelian identity)

~ Kleiss-Kuijf relations

$$\mathcal{A}_{\mathbf{n}}^{\text{tree}}(\mathbf{1,2,3,\dots,n}) = \mathbf{g_s^{n-2}} \sum_{\sigma = \mathcal{P}(\mathbf{2,3,\dots,n-1})} \left(\mathbf{F^{a_{\sigma(2)}} \dots F^{a_{\sigma(n-1)}}}\right)_{\mathbf{a_1 a_n}} \mathbf{m_n}(\mathbf{g_1, g_{\sigma(2)}, g_{\sigma(3)}, \dots, g_{\sigma(n-1)}, g_n}).$$

(n-2)! color ordered sub-amplitudes (see also BCJ relations)

Unitary color basis: on each pole of the tree amplitude the color factor of a given colorless amplitude also factorizes

three gluon vertex



$$i\frac{g}{\sqrt{2}}\left((k_1-k_2)_{\mu_3}g_{\mu_1\mu_2}+(k_2-k_3)_{\mu_1}g_{\mu_2\mu_3}+(k_3-k_1)_{\mu_2}g_{\mu_3\mu_1}\right.$$

four gluon vertex



$$i\frac{g^2}{2}(2g_{\mu_1\mu_3}g_{\mu_2\mu_4}-g_{\mu_1\mu_4}g_{\mu_2\mu_3}-g_{\mu_1\mu_2}g_{\mu_3\mu_4})$$

gluon-quark-antiquark vertex



sign depends on orientation

Color stripped Feynman rules for color ordered sub-amplitudes

three gluon vertex



$$i\frac{g}{\sqrt{2}}\left((k_1-k_2)_{\mu_3}g_{\mu_1\mu_2}+(k_2-k_3)_{\mu_1}g_{\mu_2\mu_3}+(k_3-k_1)_{\mu_2}g_{\mu_3\mu_1}\right)$$

four gluon vertex



$$i\frac{g^2}{2}(2g_{\mu_1\mu_3}g_{\mu_2\mu_4}-g_{\mu_1\mu_4}g_{\mu_2\mu_3}-g_{\mu_1\mu_2}g_{\mu_3\mu_4})$$

gluon-quark-antiquark vertex



sign depends on orientation

Color ordered n-gluon one-loop sub-amplitudes

$$\mathcal{A}_{\mathbf{n}}^{1-\text{loop}} = \mathbf{g}_{\mathbf{s}}^{\mathbf{n}} \mathbf{c}_{\mathbf{\Gamma}} \sum_{\sigma \in \mathbf{S}_{\mathbf{n-1}}} \mathbf{A}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g}_{\sigma(\mathbf{2})}, \ldots, \mathbf{g}_{\sigma(\mathbf{n})}),$$

$$\mathbf{c}_{\mathbf{\Gamma}} = \frac{\mathbf{1}}{(\mathbf{4\pi})^{\mathbf{2-\epsilon}}} \frac{\mathbf{\Gamma(1+\epsilon)\Gamma^2(1-\epsilon)}}{\mathbf{\Gamma(1-2\epsilon)}}$$

Consider a double cut between external line k and k+1 and n and 1

$$\begin{aligned}\text{Im}_{(\mathbf{k},\mathbf{n})}\left[\mathbf{A}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_n})\right] &= \{\mathbf{a_1 a_2 \cdots a_k}\}_{\mathbf{vu}} \{\mathbf{a_{k+1} \cdots a_n}\}_{\mathbf{uv}} \\ &\quad \times \mathbf{m_{k+2}}(\mathbf{g_v}, \mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_k}, \mathbf{g_u}) \mathbf{m_{n-k+2}}(\mathbf{g_u}, \mathbf{g_{k+1}}, \ldots, \mathbf{g_n}, \mathbf{g_v}) \\ &= \{\mathbf{a_1 a_2 \cdots a_n}\} \, \text{Im}_{(\mathbf{k},\mathbf{n})}\left[\mathbf{m}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_n})\right]\end{aligned}$$

$$\mathcal{A}_{\mathbf{n}}^{1-\text{loop}} = \mathbf{g}_{\mathbf{s}}^{\mathbf{n}} \mathbf{c}_{\mathbf{\Gamma}} \sum_{\mathcal{P}(\mathbf{2},\cdots,\mathbf{n})/\mathcal{R}} \text{Tr}(\mathbf{F}^{\mathbf{a_1}}, \ldots, \mathbf{F}^{\mathbf{a_n}}) \mathbf{m}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_n})$$

one loop color order sub-amplitude

A reflection transformation is factored out. The cyclic property and reflection symmetry remain valid. The number of independent one-loop amplitudes is (n-1)!/2.

Color ordered n-gluon one-loop sub-amplitudes for large N_c

Decomposition in T-basis:

$$\mathcal{A}_{n}^{1-\text{loop}} = g_s^n N_c c_\Gamma \sum_{\mathcal{P}(2,\cdots,n)/\mathcal{R}} \text{Tr}(T^{a_1},\ldots,T^{a_n}) m_{1,n}^{(1)}(g_1,g_2,\ldots,g_n)$$

$$+g_s^n c_\Gamma \sum_{r=2}^{[n/2]+1} \left(\sum_{\mathcal{P}(2,\cdots,n)/\mathcal{Z}_{r-1}\times\mathcal{Z}_{n-r+1}} \text{Tr}\left(T^{a_1},\ldots,T^{a_{r-1}}\right) \text{Tr}\left(T^{a_r},\ldots,T^{a_n}\right) m_{2,n}^{(1)}\left(g_1,g_2,\ldots,g_n\right) \right)$$

- The single-trace color structures have an explicit factor of Nc out.
- They dominate in the large Nc limit.
- The planar L-loop color decomposition formula remains the same .
- The decomposition remains the same also for the N=4sYM theory.
- T-based color decomposition is preferred.

Two particle unitarity gives color decomposition of a quark-loop to n-gluon amplitudes

$$\mathcal{A}_{n;n_f}^{1-\text{loop}} = g_s^n c_\Gamma n_f \sum_{\sigma\in S_{n-1}} \text{Tr}(T^{a_1} T^{a_{\sigma(2)}} \ldots T^{a_{\sigma n}})\, m_{n;n_f}^{(1)}\left(g_1, g_{\sigma(2)},\ldots,g_{\sigma(n)}\right),$$

$\bar{q}q + (n-2)g$ amplitudes and fully ordered primitive amplitudes

Color factors of Feynman diagrams $\quad \left(\mathbf{T}^{\mathbf{b_1}} \dots \mathbf{T}^{\mathbf{b_k}} \dots\right)_{\mathbf{j\bar{\imath}}} \times \left(\mathbf{F}^{\mathbf{a_1}} \dots \mathbf{F}^{\mathbf{a_r}}\right)_{\mathbf{b_1 a_{r+1}}} \dots \left(\mathbf{F}^{\mathbf{a_p}} \dots \mathbf{F}^{\mathbf{a_t - 1}}\right)_{\mathbf{b_k a_t}} \dots$

$$\mathcal{A}_{\mathbf{n}}^{\mathrm{tree}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \mathbf{g_3}, \dots, \mathbf{g_n}) = \mathbf{g_s^{n-2}} \sum_{\sigma \in \mathbf{S_{n-2}}} \left(\mathbf{T}^{\mathbf{a_{\sigma(3)}}} \mathbf{T}^{\mathbf{a_{\sigma(4)}}} \dots \mathbf{T}^{\mathbf{a_{\sigma(n)}}}\right)_{\mathbf{i_2 \bar{\imath}_1}} \mathbf{m_n}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \mathbf{g_{\sigma(3)}}, \dots, \mathbf{g_{\sigma(n)}}).$$

(n-2)! colorless color ordered tree sub-amplitudes

the quark labels do not participate in the permutation sum

Decomposition in mixed basis such that the quark is also in the permutation sum

$$\mathcal{A}_{\mathbf{n}}^{\mathrm{tree}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \mathbf{g_3}, \dots, \mathbf{g_n}) = \mathbf{g_s^{n-2}} (-\mathbf{1})^{\mathbf{n}} \sum_{\mathbf{k=3}}^{\mathbf{n}} \sum_{\mathcal{P}(\mathbf{4},\dots,\mathbf{n})} \left(\mathbf{T^y}\, \mathbf{T}^{\mathbf{a_{k+1}}} .. \mathbf{T}^{\mathbf{a_n}}\right)_{\mathbf{i_2 \bar{\imath}_1}} \mathrm{Tr}\left(\mathbf{F}^{\mathbf{a_4}} \dots \mathbf{F}^{\mathbf{a_k}}\right)_{\mathbf{a_3 y}}$$
$$\times \tilde{\mathbf{m}}_{\mathbf{n}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{g_n}, \dots, \mathbf{g_{k+1}}, \mathbf{q_2}, \mathbf{g_k}, \dots, \mathbf{g_3})$$

Color ordered tree “left primitive amplitudes”

$$\tilde{\mathbf{m}}_{\mathbf{n}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{g_n}, .., \mathbf{g_{(k+1)}}, \mathbf{q_2}, \mathbf{g_k}, .., \mathbf{g_3}) = (-\mathbf{1})^{\mathbf{k}} \mathbf{m_n}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \underline{\mathbf{k}, ..., \mathbf{3}}, \overline{(\mathbf{k+1}), .., \mathbf{n}})$$

Excercise: derive this relation using commutator identities

When anti-quark is in the permutation sum: “right primitive amplitude”

This mixed basis is “unitary”

$\bar{\mathbf{q}}\mathbf{q}+(\mathbf{n}-\mathbf{2})\mathbf{g}$ colorless one-loop fully ordered left primitive amplitude



$$\mathcal{A}_{\mathbf{n}}^{1-\text{loop}}(\bar{\mathbf{q}}_{\mathbf{1}},\mathbf{q}_{\mathbf{2}},\mathbf{g}_{\mathbf{3}},..,\mathbf{g}_{\mathbf{n}})=\mathbf{g}_{\mathbf{s}}^{\mathbf{n}}\mathbf{c}_{\Gamma}\sum_{\mathbf{p=2}}^{\mathbf{n}}\sum_{\sigma\in\mathbf{S}_{\mathbf{n-2}}}(\mathbf{T}^{\mathbf{x_2}}\mathbf{T}^{\mathbf{a}_{\sigma_3}}..\mathbf{T}^{\mathbf{a}_{\sigma_p}}\mathbf{T}^{\mathbf{x_1}})_{\mathbf{i_2\bar{\imath}_1}}(\mathbf{F}^{\mathbf{a}_{\sigma_{p+1}}}..\mathbf{F}^{\mathbf{a}_{\sigma_n}})_{\mathbf{x_1x_2}}$$
$$\times(-\mathbf{1})^{\mathbf{n}}\tilde{\mathbf{m}}_{\mathbf{n}}^{(\mathbf{1})}(\bar{\mathbf{q}}_{\mathbf{1}},\mathbf{g}_{\sigma(\mathbf{p})},..,\mathbf{g}_{\sigma(\mathbf{3})},\mathbf{q}_{\mathbf{2}},\mathbf{g}_{\sigma(\mathbf{n})},..,\mathbf{g}_{\sigma(\mathbf{p+1})})$$

Color ordered amplitudes for n-gluons and primitive amplitudes
for $\bar{\mathbf{q}}\mathbf{q}+(\mathbf{n}-\mathbf{2})\mathbf{g}$ can be calculated using colorless Feynman rules
Berends-Giele recursion relations

Comment: Leading color is good approximations for gluons only

Amplitudes with multiple quarks

Colorless ordered amplitudes, primitive amplitudes, parent diagrams

$$\mathcal{B}^{\rm tree}(\bar{\mathbf{q}}_1,\mathbf{q}_2,\bar{\mathbf{Q}}_3,\mathbf{Q}_4,\mathbf{g}_5)=\mathbf{g}_s^3\Big[(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_1}\delta_{\mathbf{i}_2\bar{\imath}_3}\mathbf{B}^{\rm tree}_{\mathbf{5};1}+\frac{\mathbf{1}}{\mathbf{N_c}}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_1}\delta_{\mathbf{i}_4\bar{\imath}_3}\mathbf{B}^{\rm tree}_{\mathbf{5};2}$$
$$+(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_3}\delta_{\mathbf{i}_4\bar{\imath}_1}\mathbf{B}^{\rm tree}_{\mathbf{5};3}+\frac{\mathbf{1}}{\mathbf{N_c}}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_3}\delta_{\mathbf{i}_2\bar{\imath}_1}\mathbf{B}^{\rm tree}_{\mathbf{5};4}\Big],$$

$$\mathcal{B}^{1-\rm loop}(\bar{\mathbf{q}}_1,\mathbf{q}_2,\bar{\mathbf{Q}}_3,\mathbf{Q}_4,\mathbf{g}_5)=\mathbf{g}_s^5\Big[\mathbf{N_c}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_1}\delta_{\mathbf{i}_2\bar{\imath}_3}\mathbf{B}_{\mathbf{5};1}+(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_1}\delta_{\mathbf{i}_4\bar{\imath}_3}\mathbf{B}_{\mathbf{5};2}+\mathbf{N_c}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_3}\delta_{\mathbf{i}_4\bar{\imath}_1}\mathbf{B}_{\mathbf{5};3}$$
$$+(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_3}\delta_{\mathbf{i}_2\bar{\imath}_1}\mathbf{B}_{\mathbf{5};4}\Big]$$

$$\mathbf{B}_{\mathbf{5};\mathbf{i}}=\mathbf{B}^{[\mathbf{1}]}_{\mathbf{5};\mathbf{i}}+\frac{\mathbf{n_f}}{\mathbf{N_c}}\mathbf{B}^{[\mathbf{1/2}]}_{\mathbf{5};\mathbf{i}},\qquad \mathbf{i}=\mathbf{1},\mathbf{2},\mathbf{3},\mathbf{4},$$

$$\mathbf{B}^{[1/2]}_{5;1}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{g}_5,\mathbf{Q}_4,\bar{\mathbf{Q}}_3,\mathbf{q}_2),\quad \mathbf{B}^{[1/2]}_{5;2}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{Q}_4,\bar{\mathbf{Q}}_3,\mathbf{q}_2,\mathbf{g}_5),$$
$$\mathbf{B}^{[1/2]}_{5;3}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{Q}_4,\bar{\mathbf{Q}}_3,\mathbf{g}_5,\mathbf{q}_2),\quad \mathbf{B}^{[1/2]}_{5;4}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{Q}_4,\mathbf{g}_5,\bar{\mathbf{Q}}_3,\mathbf{q}_2).$$



Amplitudes with multiple quarks

Colorless ordered amplitudes, primitive amplitudes, parent diagrams

More and more complicated.....

End of color management

Singular behavior of one loop primitive amplitudes

n-gluon:

$$\mathbf{m_n^{(1)}(g_1,g_2,\ldots g_n)} = -\sum_{\mathbf{i=1}}^{\mathbf{n}} \left[\frac{\mathbf{1}}{\epsilon^{\mathbf{2}}} + \frac{\mathbf{1}}{\epsilon}\left(\frac{\mathbf{11}}{\mathbf{3n}} + \mathbf{L_{i,i+1}}\right)\right] \mathbf{m_n^{(0)}(g_1,g_2,\ldots g_n)},$$

$$\mathbf{L_{kn}} = \ln\left(\mu^{\mathbf{2}}/(-\mathbf{s_{kn}} - \mathbf{i0})\right)$$

$\mathbf{\bar{q}q + (n-2)g}$:

$$\mathbf{\tilde{m}_n^{(1)}(\bar{q}_n, g_{k+1}, \ldots, g_{n-1}, q_2, g_3, \ldots g_k)} =$$

$$\mathbf{\tilde{m}_n(\bar{q}_n, g_{k+1}, \ldots, g_{n-1}, q_2, g_3, \ldots g_k)} \left[-\frac{\mathbf{k}}{\epsilon^{\mathbf{2}}} - \frac{\mathbf{1}}{\epsilon}\left(\frac{\mathbf{3}}{\mathbf{2}} + \sum_{\mathbf{i=1}}^{\mathbf{k-1}} \mathbf{L_{i\,i+1}} + \mathbf{L_{kn}}\right)\right]$$

Color is eliminated. Important for testing the calculations.
Primitive tree amplitude is calculated with Berends-Giele recursion relations based on color stripped Feynman rules or BCFW recursion relations

Parma International School of Theoretical Physics

September 3 - September 8, 2012

Parma, Italy

Scattering Amplitudes in QCD, Supersymmetric Gauge Theories and Supergravity

One-loop calculations in QCD and N=4 supersymmetric gauge theories

OUTLINE

Lecture 1: QCD and review its main features.

Lecture 2: One loop tensor integrals and their reduction

Lecture 3: Unitarity method and amplitudes

Lecture 4: Analytic and numerical computations

Lecture 5: Different implementations, Outlook

One-loop calculations in quantum field theory: from Feynman diagrams to unitarity cuts.
R.Keith Ellis,,Zoltan Kunszt, Kirill Melnikov, Giulia Zanderighi,
to appear in Phys. Rep. arXiv:1105.4319 [hep-ph]

One loop calculation with traditional methods



One loop calculation with traditional methods



$\mathcal{N}(1)$ =1 one loop scalar, otherwise one loop tensor integral

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\mathbf{r \geq 2N - 4}$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \rightarrow \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \rightarrow \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \quad , \; \mathbf{d(l)=l^2-\mu^2}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N} \sim \int \frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \frac{\mathcal{N}(\mathbf{l})}{((\mathbf{l}+\mathbf{q_0})^2-\mathbf{m_1^2})((\mathbf{l}+\mathbf{q_1})^2-\mathbf{m_2^2})....((\mathbf{l}+\mathbf{q_{N-1}})^2-\mathbf{m_N^2})}$$

UV divergent if $\quad \mathbf{r \geq 2N-4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\frac{\mathrm{d}^4\mathbf{l}}{(2\pi)^4} \rightarrow \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \quad , \ \mathbf{d(l)=l^2-\mu^2}$$

$$\mathbf{I_4} \rightarrow \int \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \frac{\mathbf{l_\mu l_\nu l_\rho l_\delta}}{[\mathbf{d(l)}]^4} = \frac{(\mathbf{g_{\mu\nu}g_{\rho\delta}+g_{\mu\rho}g_{\nu\delta}+g_{\mu\delta}g_{\nu\rho}})}{\mathbf{D(D+2)}} \int \frac{\mathrm{d}^\mathbf{D}\mathbf{l}}{(2\pi)^\mathbf{D}} \frac{\mathbf{l^4}}{[\mathbf{d(l)}]^4}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu} g_{\rho\delta} + g_{\mu\rho} g_{\nu\delta} + g_{\mu\delta} g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = \; (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s + D/2)\Gamma(m - s - D/2)}{\Gamma(D/2)\Gamma(m)},$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu} g_{\rho\delta} + g_{\mu\rho} g_{\nu\delta} + g_{\mu\delta} g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\mathbf{\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = \; (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s+D/2)\Gamma(m-s-D/2)}{\Gamma(D/2)\Gamma(m)},}$$

$$\mathbf{\Omega_D = 2\pi^{D/2}/\Gamma(D/2)}$$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu} g_{\rho\delta} + g_{\mu\rho} g_{\nu\delta} + g_{\mu\delta} g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\mathbf{\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = \; (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s+D/2)\Gamma(m-s-D/2)}{\Gamma(D/2)\Gamma(m)},}$$

$$\mathbf{\Omega_D = 2\pi^{D/2}/\Gamma(D/2)}$$

Finite result for $\quad \mathbf{D \neq 4\,, m = 4\,, s = 2}$

One loop calculation with traditional methods



$\mathcal{N}(\mathbf{l})$ =1 one loop scalar, otherwise one loop tensor integral

$\mathcal{N}(\mathbf{l})$ is polynomial of degree r in l $\quad \mathbf{r \leq N}$

$$\mathbf{I_N \sim \int \frac{d^4 l}{(2\pi)^4} \frac{\mathcal{N}(l)}{((l+q_0)^2 - m_1^2)((l+q_1)^2 - m_2^2)....((l+q_{N-1})^2 - m_N^2)}}$$

UV divergent if $\quad \mathbf{r \geq 2N - 4}$

Only one-, two-, three- and four-point one-loop integrals can be UV-divergent.

$$\mathbf{\frac{d^4 l}{(2\pi)^4} \rightarrow \frac{d^D l}{(2\pi)^D} \quad , \; d(l) = l^2 - \mu^2}$$

$$\mathbf{I_4 \rightarrow \int \frac{d^D l}{(2\pi)^D} \frac{l_\mu l_\nu l_\rho l_\delta}{[d(l)]^4} = \frac{(g_{\mu\nu}g_{\rho\delta} + g_{\mu\rho}g_{\nu\delta} + g_{\mu\delta}g_{\nu\rho})}{D(D+2)} \int \frac{d^D l}{(2\pi)^D} \frac{l^4}{[d(l)]^4}}$$

$$\mathbf{\int \frac{d^D l}{i\pi^{D/2}} \frac{(l^2)^s}{\left(l^2 - \mu^2\right)^m} = \frac{\Omega_D}{\pi^{D/2}} (-1)^{s-m} \mu^{D+2s-2m} \int_0^\infty dx \frac{x^{2s+D-1}}{(1+x^2)^m} = (-1)^{r-s} \mu^{D+2s-2m} \frac{\Gamma(s+D/2)\Gamma(m-s-D/2)}{\Gamma(D/2)\Gamma(m)},}$$

$$\mathbf{\Omega_D = 2\pi^{D/2}/\Gamma(D/2)}$$

IR divergent if sufficient number of propagators can go to mass-shell. Soft and collinear singularities.

Finite result for $\quad \mathbf{D \neq 4\,, m = 4\,, s = 2}$

Integral basis in the limit $\mathrm{D} \to 4$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j}} \mathbf{I_{4;j}} + \mathbf{c_{3;j}} \mathbf{I_{3;j}} + \mathbf{c_{2;j}} \mathbf{I_{2;j}} + \mathbf{c_{1;j}} \mathbf{I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D}-4).$$

Only one-,two-,three-,four-point scalar integrals occur.

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j}} \mathbf{I_{4;j}} + \mathbf{c_{3;j}} \mathbf{I_{3;j}} + \mathbf{c_{2;j}} \mathbf{I_{2;j}} + \mathbf{c_{1;j}} \mathbf{I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D}-4).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j}} \mathbf{I_{4;j}} + \mathbf{c_{3;j}} \mathbf{I_{3;j}} + \mathbf{c_{2;j}} \mathbf{I_{2;j}} + \mathbf{c_{1;j}} \mathbf{I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D}-4).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j}} + \mathbf{c_{3;j} I_{3;j}} + \mathbf{c_{2;j} I_{2;j}} + \mathbf{c_{1;j} I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D-4}).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1}},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j}} + \mathbf{c_{3;j} I_{3;j}} + \mathbf{c_{2;j} I_{2;j}} + \mathbf{c_{1;j} I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D-4}).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1}},$$

$$\mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{\mathbf{4-D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j} + \mathcal{R} + \mathcal{O}(D-4)}.$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1}},$$

$$\mathbf{d_i = (l + q_{i-1})^2 - m_i^2 + i\varepsilon} \qquad \mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j} + \mathcal{R} + \mathcal{O}(D-4)}.$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1}},$$

$$\mathbf{d_i = (l + q_{i-1})^2 - m_i^2 + i\varepsilon} \qquad \mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{q_n \equiv \sum_{i=1}^{n} p_i}, \quad \mathbf{q_0 = 0}$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2}},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3}},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2) = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3 d_4}},$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j} + \mathcal{R} + \mathcal{O}(D-4)}.$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}$, $\mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1},$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2},$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3},$$

$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{4-D}}{i\pi^{\frac{D}{2}} r_\Gamma} \int \frac{d^D l}{d_1 d_2 d_3 d_4},$$

$$\mathbf{d_i = (l + q_{i-1})^2 - m_i^2 + i\varepsilon} \qquad \mathbf{s_{ij} = (p_i + p_j)^2},$$

$$\mathbf{q_n \equiv \sum_{i=1}^{n} p_i}, \quad \mathbf{q_0 = 0}$$

$$\mathbf{r_\Gamma = \Gamma^2(1-\epsilon)\Gamma(1+\epsilon)/\Gamma(1-2\epsilon)}$$

Integral basis in the limit $\mathrm{D} \to 4$

Any one-loop integral can be written as linear combination of scalar one-loop integrals and rational terms:

$$\mathbf{I_N} = \sum_{\mathbf{j}} \mathbf{c_{4;j} I_{4;j} + c_{3;j} I_{3;j} + c_{2;j} I_{2;j} + c_{1;j} I_{1;j}} + \mathcal{R} + \mathcal{O}(\mathbf{D-4}).$$

Only one-,two-,three-,four-point scalar integrals occur.

The problem of the analytic calculation of one-loop scalar integrals $\mathbf{I_{r;j}}, \quad \mathrm{r} = 1, \ldots, 4$ and of their numerical evaluation is solved (QCDLoop, OneLOop).

The problem of one-loop calculation is reduced to the determination of the coefficient $\mathbf{c_{n;j}}$ and $\mathcal{R}$

$$\mathbf{I_1(m_1^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l + q_{i-1}})^2 - \mathbf{m_i^2} + \mathbf{i}\varepsilon \qquad \mathbf{s_{ij}} = (\mathbf{p_i + p_j})^2 ,$$

$$\mathbf{I_2(p_1^2; m_1^2, m_2^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2}},$$

$$\mathbf{q_n} \equiv \sum_{\mathbf{i=1}}^{\mathbf{n}} \mathbf{p_i}\,, \quad \mathbf{q_0 = 0}$$

$$\mathbf{r_\Gamma} = \mathbf{\Gamma^2}(1-\epsilon)\mathbf{\Gamma}(1+\epsilon)/\mathbf{\Gamma}(1-\mathbf{2}\epsilon)$$

$$\mathbf{I_3(p_1^2, p_2^2, p_3^2; m_1^2, m_2^2, m_3^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3}},$$



$$\mathbf{I_4(p_1^2, p_2^2, p_3^2, p_4^2; s_{12}, s_{23}; m_1^2, m_2^2, m_3^2, m_4^2)} = \frac{\mu^{4-\mathrm{D}}}{\mathbf{i\pi^{\frac{D}{2}} r_\Gamma}} \int \frac{\mathbf{d^D l}}{\mathbf{d_1 d_2 d_3 d_4}} ,$$

Only the constant terms and some of the terms depending on $(\mathbf{l} \cdot \mathbf{n}_\epsilon)^2$ give non-vanishing integrals

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}\mathbf{i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

$$\mathcal{R} = -\sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \frac{\tilde{\mathbf{d}}^{(\mathbf{4})}_{\mathbf{i_1 i_2 i_3 i_4}}}{\mathbf{6}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \frac{\tilde{\mathbf{c}}^{(\mathbf{7})}_{\mathbf{i_1 i_2 i_3}}}{\mathbf{2}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \left[\frac{\mathbf{m^2_{i_1}} + \mathbf{m^2_{i_2}}}{\mathbf{2}} - \frac{(\mathbf{q_{i_1}} - \mathbf{q_{i_2}})^{\mathbf{2}}}{\mathbf{6}} \right] \tilde{\mathbf{b}}^{(\mathbf{9})}_{\mathbf{i_1 i_2}}.$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

Reduction with the method of form factor expansions (Passarino, Veltman, 1979):

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

Reduction with the method of form factor expansions (Passarino, Veltman, 1979):

Examples:

$$\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}\,, \quad \mathbf{C^{\mu\nu} = g^{\mu\nu} C_{00} + \sum_{i,j=1}^{2} p_i^\mu p_j^\nu C_{ij}}\,, \quad \mathbf{C_{12} = C_{21}}$$

PV reduction of one-loop tensor integrals (coming from individual Feynman diagrams):

$$\mathbf{A_0(m_1)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1}}{\mathbf{d_1}},$$

$$\mathbf{d_i} = (\mathbf{l} + \sum_{\mathbf{k=1}}^{\mathbf{i-1}} \mathbf{p_k})^{\mathbf{2}} - \mathbf{m_i^2}$$

$$\mathbf{B_0; B^\mu; B^{\mu\nu}(p_1, m_1, m_2)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu}}{\mathbf{d_1 d_2}},$$

$$\mathbf{C_0; C^\mu; C^{\mu\nu}; C^{\mu\nu\alpha}(p_1, p_2, m_1, m_2, m_3)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha}}{\mathbf{d_1 d_2 d_3}},$$

$$\mathbf{D_0; D^\mu; D^{\mu\nu}; D^{\mu\nu\alpha}; D^{\mu\nu\alpha\beta}(p_1, p_2, p_3, m_1, m_2, m_3, m_4)} = \frac{\mathbf{1}}{\mathbf{i\pi^{D/2}}} \int \mathbf{d^D l}\, \frac{\mathbf{1; l^\mu; l^\mu l^\nu; l^\mu l^\nu l^\alpha; l^\mu l^\nu l^\alpha l^\beta}}{\mathbf{d_1 d_2 d_3 d_4}}$$

Reduction with the method of form factor expansions (Passarino, Veltman, 1979):

Examples:

$$\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}\,, \quad \mathbf{C^{\mu\nu} = g^{\mu\nu} C_{00} + \sum_{i,j=1}^{2} p_i^\mu p_j^\nu C_{ij}}\,, \quad \mathbf{C_{12} = C_{21}}$$

rank 1 and 2 tensor three point integrals

Reduction of $\mathbf{C}^\mu = \mathbf{p}_1^\mu \mathbf{C_1} + \mathbf{p}_2^\mu \mathbf{C_2}$,

Reduction of $\mathbf{C}^\mu = \mathbf{p_1^\mu C_1 + p_2^\mu C_2}\,,$

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\langle \mathbf{l \cdot p_j} \rangle = \int \frac{\mathbf{d^D l}}{\mathbf{i\pi^{D/2}}} \frac{\mathbf{l \cdot p_j}}{\mathbf{d_1 d_2 d_3}}, \quad \text{for } \mathbf{j = 1, 2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

Reduction of $\mathbf{C}^\mu = \mathbf{p}_1^\mu \mathbf{C_1} + \mathbf{p}_2^\mu \mathbf{C_2}$,

$$\mathbf{G_2} \begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\langle \mathbf{l} \cdot \mathbf{p_j} \rangle = \int \frac{\mathbf{d^D l}}{\mathbf{i}\pi^{\mathbf{D/2}}} \frac{\mathbf{l} \cdot \mathbf{p_j}}{\mathbf{d_1 d_2 d_3}}, \quad \text{for } \mathbf{j} = \mathbf{1}, \mathbf{2}$$

$$\mathbf{R_1^{[c]}} = \frac{\mathbf{1}}{\mathbf{2}} (\mathbf{f_1 C_0(1,2,3)} + \mathbf{B_0(1,3)} - \mathbf{B_0(2,3)}),$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

$$\mathbf{R_2^{[c]}} = \frac{\mathbf{1}}{\mathbf{2}} (\mathbf{f_2 C_0(1,2,3)} + \mathbf{B_0(1,2)} - \mathbf{B_0(1,3)})$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\langle \mathbf{l \cdot p_j} \rangle = \int \frac{\mathbf{d^D l}}{\mathbf{i\pi^{D/2}}} \frac{\mathbf{l \cdot p_j}}{\mathbf{d_1 d_2 d_3}}, \quad \text{for } \mathbf{j = 1, 2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l \cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1), \quad f_1 = m_2^2 - m_1^2 - p_1^2},$$

$$\mathbf{l \cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2), \quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1 \cdot p_2}\,.$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2, m_2, m_3) = \int \frac{d^D l}{i\pi^{D/2}}\frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\,\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}}\begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2,m_2,m_3) = \int \frac{d^D l}{i\pi^{D/2}}\,\frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l \cdot p_1 \rangle \\ \langle l \cdot p_2 \rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l \cdot p_j \rangle = \int \frac{d^D l}{i\pi^{D/2}} \frac{l \cdot p_j}{d_1 d_2 d_3}, \quad \text{for } j = 1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1 \cdot p_1 & p_1 \cdot p_2 \\ p_1 \cdot p_2 & p_2 \cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l \cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1), \quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l \cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2), \quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1 \cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2, m_2, m_3) = \int \frac{d^D l}{i\pi^{D/2}} \frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}}\begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2,m_2,m_3) = \int \frac{d^D l}{i\pi^{D/2}}\frac{1}{(l^2-m_2^2)((l+p_2)^2-m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{p_1}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_1 C_{11} + p_1{\cdot}p_2 C_{12} + C_{00}) + p_2^\nu(p_1{\cdot}p_1 C_{12} + p_1{\cdot}p_2 C_{22})},$$
$$\mathbf{p_2}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_2 C_{11} + p_2{\cdot}p_2 C_{12}) + p_2^\nu(p_1{\cdot}p_2 C_{12} + p_2{\cdot}p_2 C_{22} + C_{00})}\,.$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{G_2}\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \begin{pmatrix} \langle l\cdot p_1\rangle \\ \langle l\cdot p_2\rangle \end{pmatrix} = \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix},$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{\langle l\cdot p_j\rangle = \int \frac{d^D l}{i\pi^{D/2}}\frac{l\cdot p_j}{d_1 d_2 d_3},\quad \text{for } j=1,2}$$

$$\mathbf{G_2} = \begin{pmatrix} p_1\cdot p_1 & p_1\cdot p_2 \\ p_1\cdot p_2 & p_2\cdot p_2 \end{pmatrix},$$

$$\mathbf{R_1^{[c]} = \frac{1}{2}(f_1 C_0(1,2,3) + B_0(1,3) - B_0(2,3))},$$

$$\mathbf{R_2^{[c]} = \frac{1}{2}(f_2 C_0(1,2,3) + B_0(1,2) - B_0(1,3))}$$

cancel terms in numerator with denominator factors

$$\mathbf{l\cdot p_1 = \frac{1}{2}(f_1 + d_2 - d_1),\quad f_1 = m_2^2 - m_1^2 - p_1^2,}$$

$$\mathbf{l\cdot p_2 = \frac{1}{2}(f_2 + d_3 - d_2),\quad f_2 = m_3^2 - m_2^2 - p_2^2 - 2p_1\cdot p_2}\,.$$

Solve for $\mathbf{C_1}$ and $\mathbf{C_2}$

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}}\begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{B_0(2,3) \equiv B_0(p_2,m_2,m_3) = \int \frac{d^D l}{i\pi^{D/2}}\frac{1}{(l^2 - m_2^2)((l+p_2)^2 - m_3^2)}}$$

Reduction of $\mathbf{C^\mu = p_1^\mu C_1 + p_2^\mu C_2}$,

$$\mathbf{p_1}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_1 C_{11} + p_1{\cdot}p_2 C_{12} + C_{00}) + p_2^\nu(p_1{\cdot}p_1 C_{12} + p_1{\cdot}p_2 C_{22})},$$

$$\mathbf{p_2}_\mu \mathbf{C^{\mu\nu}} = \mathbf{p_1^\nu(p_1{\cdot}p_2 C_{11} + p_2{\cdot}p_2 C_{12}) + p_2^\nu(p_1{\cdot}p_2 C_{12} + p_2{\cdot}p_2 C_{22} + C_{00})}\,.$$

trace with $\mathbf{p_1}$ and $\mathbf{p_2}$

$$\mathbf{C_\mu^\nu = B_0(2,3) + m_1^2 C_0(1,2,3) = DC_{00} + p_1^2 C_{11} + 2p_1{\cdot}p_2 C_{12} + p_2^2 C_{22}}$$

trace with $\mathbf{g^{\mu\nu}}$

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

Reduction chains for Passarino-Veltman procedure.

For more then 4 particles it is cumbersome

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

For more then 4 particles it is cumbersome

- The number of Feynman diagrams grows dramatically with the number of external particle (factorial).
- The number of terms generated during the reduction of tensor integrals grows rapidly with the number of external particles and with the rank of the integral.
- In cases with degenerate kinematics, the reduction procedure may lead to numerical instabilities.

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

For more then 4 particles it is cumbersome

- The number of Feynman diagrams grows dramatically with the number of external particle (factorial).
- The number of terms generated during the reduction of tensor integrals grows rapidly with the number of external particles and with the rank of the integral.
- In cases with degenerate kinematics, the reduction procedure may lead to numerical instabilities.

Well established technique, applied in many cases.

Reduction chains for Passarino-Veltman procedure.

$\mathbf{D_{ijkl}}$	$\rightarrow$	$D_{00ij}, D_{ijk}, C_{ijk}, C_{ij}, C_i, C_0$
$\mathbf{D_{00ij}}$	$\rightarrow$	$D_{ijk}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{0000}}$	$\rightarrow$	D_{00i}, D_{00}, C_{00}
$\mathbf{D_{ijk}}$	$\rightarrow$	$D_{00i}, D_{ij}, C_{ij}, C_i$
$\mathbf{D_{00i}}$	$\rightarrow$	D_{ij}, D_i, C_i, C_0
$\mathbf{D_{ij}}$	$\rightarrow$	D_{00}, D_i, C_i, C_0
$\mathbf{D_{00}}$	$\rightarrow$	D_i, D_0, C_0
$\mathbf{D_i}$	$\rightarrow$	D_0, C_0
$\mathbf{C_{ijk}}$	$\rightarrow$	$C_{00i}, C_{ij}, B_{ij}, B_i$
$\mathbf{C_{00i}}$	$\rightarrow$	C_{ii}, C_i, B_i, B_0
$\mathbf{C_{ij}}$	$\rightarrow$	C_{00}, C_i, B_i, B_0
$\mathbf{C_{00}}$	$\rightarrow$	C_i, C_0, B_0
$\mathbf{C_i}$	$\rightarrow$	C_0, B_0
$\mathbf{B_{ii}}$	$\rightarrow$	B_{00}, B_i, A_0
$\mathbf{B_{00}}$	$\rightarrow$	B_i, B_0, A_0
$\mathbf{B_i}$	$\rightarrow$	B_0, A_0

For more then 4 particles it is cumbersome

- The number of Feynman diagrams grows dramatically with the number of external particle (factorial).
- The number of terms generated during the reduction of tensor integrals grows rapidly with the number of external particles and with the rank of the integral.
- In cases with degenerate kinematics, the reduction procedure may lead to numerical instabilities.

Well established technique, applied in many cases.

For 5-,6-, and more particles external momenta not linearly independent (additional input), Denner, Dittmaier 2006

Singular regions in PV reduction:

Singular regions in PV reduction:

Example: rank 1 triangle

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$

but the integral is well defined in this limit

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ and necessarily $\mathbf{R_1^{[c]} = 0}$ and $\mathbf{R_2^{[c]} = 0}$

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ and necessarily $\mathbf{R_1^{[c]} = 0}$ and $\mathbf{R_2^{[c]} = 0}$

Exceptional points: vanishing Gram determinants, numerical instabilities
two and three point scalar integrals become linearly dependent
also higher powers of Gram-determinants

Singular regions in PV reduction:

Example: rank 1 triangle

$$\begin{pmatrix} C_1 \\ C_2 \end{pmatrix} = \mathbf{G_2^{-1}} \begin{pmatrix} R_1^{[c]} \\ R_2^{[c]} \end{pmatrix}$$

$$\mathbf{G_2^{-1}} = \frac{\begin{pmatrix} \mathbf{p_2 \cdot p_2} & \mathbf{-p_1 \cdot p_2} \\ \mathbf{-p_1 \cdot p_2} & \mathbf{p_1 \cdot p_1} \end{pmatrix}}{\mathbf{\Delta_2(p_1, p_2)}},$$

$$\mathbf{\Delta_2(p_1, p_2)} = \det[\mathbf{G_2}] = \mathbf{p_1^2 p_2^2} - \mathbf{(p_1 \cdot p_2)^2},$$

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ but the integral is well defined in this limit

We assume in the PV expansion that $\mathbf{p_1, p_2}$ are linearly independent

if $\mathbf{p_1 || p_2}$ with $\mathbf{p_1^2 \neq 0}$ then $\mathbf{\Delta_2 = 0}$ and necessarily $\mathbf{R_1^{[c]} = 0}$ and $\mathbf{R_2^{[c]} = 0}$

Exceptional points: vanishing Gram determinants, numerical instabilities
two and three point scalar integrals become linearly dependent
also higher powers of Gram-determinants

Exercise: Investigate the collinear behaviour of reduction using form factors with

using $\mathbf{p_1^\mu}$ and $\mathbf{\tilde{p}_2^\mu = p_2^\mu - \frac{p_1 \cdot p_2}{p_1^2} p_1^\mu}$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l \cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left[((\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2} \right] = \frac{\mathbf{1}}{\mathbf{2}} \left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2} \right]$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^{\lambda}\epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^{\lambda}\epsilon^{\mathbf{q_1}\mathbf{q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda\mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^{\lambda}\epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^{\lambda}\epsilon^{\mathbf{q_1}\mathbf{q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda\mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^{\lambda}} = \frac{\epsilon^{\lambda\mathbf{q_2}}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_2^{\lambda}} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^{\lambda}\epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^{\lambda}\epsilon^{\mathbf{q_1}\mathbf{q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda\mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^{\lambda}} = \frac{\epsilon^{\lambda\mathbf{q_2}}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_2^{\lambda}} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1}\mathbf{q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

$$\mathbf{l}^{\lambda} = (\mathbf{l}\cdot\mathbf{q_1})\mathbf{v_1^{\lambda}} + (\mathbf{l}\cdot\mathbf{q_2})\mathbf{v_2^{\lambda}}.$$

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^\lambda \epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^\lambda \epsilon^{\mathbf{q_1 q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda \mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^\lambda} = \frac{\epsilon^{\lambda \mathbf{q_2}}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_2^\lambda} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

$$\mathbf{l}^\lambda = (\mathbf{l}\cdot\mathbf{q_1})\mathbf{v_1^\lambda} + (\mathbf{l}\cdot\mathbf{q_2})\mathbf{v_2^\lambda}.$$

Can we embed this a coordinate into higher dimensional space time?

van Nerveen-Vermaseren reduction of one-loop tensor integrals

Allows for decomposition at the integrand level.

PV reduction was achieved by canceling numerator terms with denominator factors:

$$\mathbf{l\cdot q_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[((\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}) - (\mathbf{l^2} - \mathbf{m_0^2}) + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right] = \frac{\mathbf{1}}{\mathbf{2}}\left[(\mathbf{d_i} - \mathbf{d_0} + \mathbf{m_i^2} - \mathbf{m_0^2} - \mathbf{q_i^2}\right]$$

Simplest example is in D=2 space-time:

$$\mathbf{l}^\lambda \epsilon^{\mu_1\mu_2} = \mathbf{l}^{\mu_1}\epsilon^{\lambda\mu_2} + \mathbf{l}^{\mu_2}\epsilon^{\mu_1\lambda}. \qquad \text{Shouten-identity}$$

$$\mathbf{l}^\lambda \epsilon^{\mathbf{q_1 q_2}} = (\mathbf{l}\cdot\mathbf{q_1})\epsilon^{\lambda \mathbf{q_2}} + (\mathbf{l}\cdot\mathbf{q_2})\epsilon^{\mathbf{q_1}\lambda},$$

$$\mathbf{v_1^\lambda} = \frac{\epsilon^{\lambda \mathbf{q_2}}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_2^\lambda} = \frac{\epsilon^{\mathbf{q_1}\lambda}}{\epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_i}\cdot\mathbf{q_j} = \delta_{\mathbf{ij}},$$

$$\mathbf{l}^\lambda = (\mathbf{l}\cdot\mathbf{q_1})\mathbf{v_1^\lambda} + (\mathbf{l}\cdot\mathbf{q_2})\mathbf{v_2^\lambda}.$$

Can we embed this a coordinate into higher dimensional space time?

Yes.

Generalized Kronecker-symbols:

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \qquad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \qquad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu = \frac{\epsilon_{q_1 q_2}\epsilon^{\mu q_2}}{\epsilon_{q_1 q_2}\epsilon^{q_1 q_2}}}, \quad \mathbf{v_2^\mu = \frac{\epsilon_{q_1 q_2}\epsilon^{q_1 \mu}}{\epsilon_{q_1 q_2}\epsilon^{q_1 q_2}}}, \quad \epsilon^{\mu_1\mu_2}\epsilon_{\nu_1\nu_2} = \delta^{\mu_1}_{\nu_1}\delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2}\delta^{\mu_2}_{\nu_1} = \det|\delta^\mu_\nu| \equiv \delta^{\mu_1\mu_2}_{\nu_1\nu_2},$$

$$\mathbf{v_1^\mu = \frac{\delta^{\mu q_2}_{q_1 q_2}}{\Delta_2}}, \quad \mathbf{v_2^\mu = \frac{\delta^{q_1 \mu}_{q_1 q_2}}{\Delta_2}}, \quad \mathbf{\Delta_2 = \delta^{q_1 q_2}_{q_1 q_2} = q_1^2 q_2^2 - (q_1 q_2)^2} \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \quad ,$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^2 \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \ ,$$

$$\mathbf{\Delta}(\mathbf{k_1}, \mathbf{k_2}, \cdots, \mathbf{k_R}) = \delta^{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}_{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^{\mathbf{2}} \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \ ,$$

$$\mathbf{\Delta}(\mathbf{k_1}, \mathbf{k_2}, \cdots, \mathbf{k_R}) = \delta^{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}_{\mathbf{k_1 k_2} \cdots \mathbf{k_R}}$$

$$\mathbf{v_i^\mu}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1} \ldots \mathbf{k_{i-1}} \mu \mathbf{k_{i+1}} \ldots \mathbf{k_{D_P}}}_{\mathbf{k_1} \ldots \mathbf{k_{i-1}} \mathbf{k_i} \mathbf{k_{i+1}} \ldots \mathbf{k_{D_P}}}}{\mathbf{\Delta}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}})} \ ,$$

Generalized Kronecker-symbols:

$$\mathbf{v_1^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mu \mathbf{q_2}}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \mathbf{v_2^\mu} = \frac{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1} \mu}}{\epsilon_{\mathbf{q_1 q_2}} \epsilon^{\mathbf{q_1 q_2}}}, \quad \epsilon^{\mu_1 \mu_2} \epsilon_{\nu_1 \nu_2} = \delta^{\mu_1}_{\nu_1} \delta^{\mu_2}_{\nu_2} - \delta^{\mu_1}_{\nu_2} \delta^{\mu_2}_{\nu_1} = \det |\delta^\mu_\nu| \equiv \delta^{\mu_1 \mu_2}_{\nu_1 \nu_2},$$

$$\mathbf{v_1^\mu} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1} \mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \quad \mathbf{\Delta_2} = \delta^{\mathbf{q_1 q_2}}_{\mathbf{q_1 q_2}} = \mathbf{q_1^2 q_2^2} - (\mathbf{q_1 q_2})^{\mathbf{2}} \quad , \quad \mu = 1, \ldots, D$$

In Kronecker-deltas we can have Lorenz-vector indices in any space and Shouten-identies also valid in any space

$$\delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} = \begin{vmatrix} \delta^{\mu_1}_{\nu_1} & \delta^{\mu_1}_{\nu_2} & \cdots & \delta^{\mu_1}_{\nu_R} \\ \delta^{\mu_2}_{\nu_1} & \delta^{\mu_2}_{\nu_2} & \cdots & \delta^{\mu_2}_{\nu_R} \\ \vdots & \vdots & & \vdots \\ \delta^{\mu_R}_{\nu_1} & \delta^{\mu_R}_{\nu_2} & \cdots & \delta^{\mu_R}_{\nu_R} \end{vmatrix}$$

$$\delta^{\mathbf{k} \mu_2 \cdots \mu_R}_{\nu_1 \mathbf{k} \cdots \nu_R} \equiv \delta^{\mu_1 \mu_2 \cdots \mu_R}_{\nu_1 \nu_2 \cdots \nu_R} \mathbf{k}_{\mu_1} \mathbf{k}^{\nu_2} \ ,$$

$$\mathbf{\Delta}(\mathbf{k_1}, \mathbf{k_2}, \cdots, \mathbf{k_R}) = \delta^{\mathbf{k_1 k_2 \cdots k_R}}_{\mathbf{k_1 k_2 \cdots k_R}}$$

$$\mathbf{v_i^\mu}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1 \ldots k_{i-1} \mu k_{i+1} \ldots k_{D_P}}}_{\mathbf{k_1 \ldots k_{i-1} k_i k_{i+1} \ldots k_{D_P}}}}{\mathbf{\Delta}(\mathbf{k_1}, \ldots, \mathbf{k_{D_P}})} \ , \qquad \mathbf{v_i \cdot k_j} = \delta_{\mathbf{ij}} \, , \quad \text{for} \quad \mathbf{j \leq D_P}$$

Physical and transverse space

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum

$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1}, \mathbf{p_2}, \ldots, \mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1}, \mathbf{p_2}, \ldots, \mathbf{p_N}; \mathbf{l})}{\mathbf{d_1 d_2 \cdots d_R}}$$



Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical space: vector space spanned by the inflow momenta. D-dimensional vectors can be decomposed to physical space and transverse space components:

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical space: vector space spanned by the inflow momenta. D-dimensional vectors can be decomposed to physical space and transverse space components:

$$\mathbf{D} = \mathbf{D_P} + \mathbf{D_T},\quad \mathbf{D_P} = \min(\mathbf{D},\mathbf{R}-\mathbf{1}),\quad \mathbf{D_T} = \max(\mathbf{0},\mathbf{D}-\mathbf{R}+\mathbf{1})\,,\qquad \mathbf{l} = \mathbf{l_P} + \mathbf{l_T}$$

Physical and transverse space

QFT in D-dimension, N-particle scattering amplitude, consider one one-loop Feynman-diagram with R loop-momentum dependent propagator. The integrand is a rational function of the loop momentum



$$\mathcal{I}_{\mathbf{N}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N}|\mathbf{l}) = \frac{\mathcal{N}_{\mathcal{I}}(\mathbf{p_1},\mathbf{p_2},\ldots,\mathbf{p_N};\mathbf{l})}{\mathbf{d_1 d_2}\cdots\mathbf{d_R}}$$

The amplitude has R inflow momenta $\mathbf{k_1},\ldots,\mathbf{k_R}$.

$$\mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}}-\mathbf{m_i^2}\,,\quad \mathbf{k_i} = \mathbf{q_i}-\mathbf{q_{i-1}},\quad \mathbf{k_i} = \sum_{\mathbf{j=1}}^{\mathbf{N}} \alpha_{\mathbf{ij}}\mathbf{p_j}\,,\qquad \sum_{\mathbf{i=1}}^{\mathbf{R}} \mathbf{k_i} = \mathbf{0},\ \alpha_{\mathbf{ij}} = \mathbf{0},\mathbf{1}$$

Physical space: vector space spanned by the inflow momenta. D-dimensional vectors can be decomposed to physical space and transverse space components:

$$\mathbf{D} = \mathbf{D_P} + \mathbf{D_T},\quad \mathbf{D_P} = \min(\mathbf{D},\mathbf{R}-\mathbf{1}),\quad \mathbf{D_T} = \max(\mathbf{0},\mathbf{D}-\mathbf{R}+\mathbf{1})\,,\qquad \mathbf{l} = \mathbf{l_P} + \mathbf{l_T}$$

If R>D, the transverse space is zero dimensional.

Use NV (dual) coordinates in the physical space:

Use NV (dual) coordinates in the physical space:

$$l^\mu = \sum_{i=1}^{D_P} (l \cdot k_i)\, v_i^\mu + \sum_{i=1}^{D_T} (l \cdot n_i)\; n_i^\mu \;\; .$$

$$g^{\mu\nu} = \sum_{i=1}^{D_P} k_i^\mu v_i^\nu + \sum_{i=1}^{D_T} n_i^\mu n_i^\nu$$

Use NV (dual) coordinates in the physical space:

$$\mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{l}\cdot\mathbf{k_i})\,\mathbf{v_i^{\mu}} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\;\mathbf{n_i^{\mu}}\;.$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} \mathbf{k_i^{\mu}}\mathbf{v_i^{\nu}} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} \mathbf{n_i^{\mu}}\mathbf{n_i^{\nu}}$$

$$\mathbf{n_r}\cdot\mathbf{n_s} = \delta_{\mathbf{rs}}\,,\;\mathbf{k_i}\cdot\mathbf{n_r} = \mathbf{0}\,,\;\mathbf{v_i}\cdot\mathbf{n_r} = \mathbf{0},\qquad \mathbf{w}^{\mu\nu} = \sum_{\mathbf{r=1}}^{\mathbf{D_T}} \mathbf{n_r^{\mu}}\mathbf{n_r^{\nu}},$$

Use NV (dual) coordinates in the physical space:

$$\mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{l}\cdot\mathbf{k_i})\,\mathbf{v_i}^{\mu} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\;\mathbf{n_i}^{\mu}\;.$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} \mathbf{k_i}^{\mu}\mathbf{v_i}^{\nu} + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} \mathbf{n_i}^{\mu}\mathbf{n_i}^{\nu}$$

$$\mathbf{n_r}\cdot\mathbf{n_s} = \delta_{\mathbf{rs}}\,,\;\mathbf{k_i}\cdot\mathbf{n_r} = \mathbf{0}\,,\;\mathbf{v_i}\cdot\mathbf{n_r} = \mathbf{0},\qquad \mathbf{w}^{\mu\nu} = \sum_{\mathbf{r=1}}^{\mathbf{D_T}} \mathbf{n_r}^{\mu}\mathbf{n_r}^{\nu},$$

$\mathbf{w}_{\mu\nu}$ is the projector operator to transverse space

$$\mathbf{w}_{\mu}{}^{\nu}(\mathbf{k_1},\ldots,\mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1}\cdots\mathbf{k_{D_P}}\,\nu}_{\mathbf{k_1}\ldots\mathbf{k_{D_P}}\,\mu}}{\mathbf{\Delta}(\mathbf{k_1},\ldots,\mathbf{k_{D_P}})}$$

$$\mathbf{w}_{\mu}{}^{\mu} = \mathbf{D_T}\,,\;\mathbf{k_i^{\mu}}\,\mathbf{w}_{\mu\nu} = \mathbf{0}\,,\;\mathbf{w}^{\mu}{}_{\alpha}\mathbf{w}^{\alpha\nu} = \mathbf{w}^{\mu\nu}.$$

Use NV (dual) coordinates in the physical space:

$$\mathbf{l}^\mu = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{l}\cdot\mathbf{k_i})\,\mathbf{v_i}^\mu + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\;\mathbf{n_i}^\mu \;.$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{i=1}}^{\mathbf{D_P}} \mathbf{k_i}^\mu \mathbf{v_i}^\nu + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} \mathbf{n_i}^\mu \mathbf{n_i}^\nu$$

$$\mathbf{n_r}\cdot\mathbf{n_s} = \delta_{\mathbf{rs}}\,,\ \mathbf{k_i}\cdot\mathbf{n_r} = \mathbf{0}\,,\ \mathbf{v_i}\cdot\mathbf{n_r} = \mathbf{0},\qquad \mathbf{w}^{\mu\nu} = \sum_{\mathbf{r=1}}^{\mathbf{D_T}} \mathbf{n_r}^\mu \mathbf{n_r}^\nu,$$

$\mathbf{w}_{\mu\nu}$ is the projector operator to transverse space

$$\mathbf{w}_\mu{}^\nu(\mathbf{k_1},\ldots,\mathbf{k_{D_P}}) \equiv \frac{\delta^{\mathbf{k_1}\cdots\mathbf{k_{D_P}}\nu}_{\mathbf{k_1}\ldots\mathbf{k_{D_P}}\mu}}{\mathbf{\Delta}(\mathbf{k_1},\ldots,\mathbf{k_{D_P}})}$$

$$\mathbf{w}_\mu{}^\mu = \mathbf{D_T}\,,\ \mathbf{k_i}^\mu\,\mathbf{w}_{\mu\nu} = \mathbf{0}\,,\ \mathbf{w}^\mu{}_\alpha \mathbf{w}^{\alpha\nu} = \mathbf{w}^{\mu\nu}.$$

The dual coordinates of the loop momentum vector provide us the reduction at the integrand level:

$$\mathbf{l}\cdot\mathbf{k_i} = \frac{\mathbf{1}}{\mathbf{2}}\left[\mathbf{d_i} - \mathbf{d_{i-1}} - \left(\mathbf{q_i^2} - \mathbf{m_i^2}\right) + \left(\mathbf{q_{i-1}^2} - \mathbf{m_{i-1}^2}\right)\right]$$

$$\mathbf{l}^\mu = \mathbf{V_R}^\mu + \frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i=1}}^{\mathbf{D_P}} (\mathbf{d_i} - \mathbf{d_{i-1}})\,\mathbf{v_i}^\mu + \sum_{\mathbf{i=1}}^{\mathbf{D_T}} (\mathbf{l}\cdot\mathbf{n_i})\,\mathbf{n_i}^\mu \;,$$

$$\mathbf{V_R}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i=1}}^{\mathbf{D_P}} \Big((\mathbf{q_i^2} - \mathbf{m_i^2}) - (\mathbf{q_{i-1}^2} - \mathbf{m_{i-1}^2})\Big)\,\mathbf{v_i}^\mu$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\, \mathcal{I}_{\mathbf{3}} \quad , \quad \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\, \mathcal{I}_{\mathbf{3}} \;\;,\;\; \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2},\, i \in [0,1,2]$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\,\mathcal{I}_{\mathbf{3}} \quad , \quad \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}}, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]\,, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]$$

$$\mathbf{l}^{\mu} = \mathbf{v_1^{\mu}}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^{\mu}}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^{\mu}} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \;\; \mathbf{v_2^{\mu}} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1},\mathbf{2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\mu}}\,\mathbf{q_i^{\nu}} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\nu}}\,\mathbf{q_i^{\mu}}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^{\mathbf{2}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{2}}}\,\mathcal{I}_{\mathbf{3}} \quad , \quad \mathcal{I}_{\mathbf{3}} = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}}, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]\,, \;\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}, \, i \in [0,1,2]$$

$$\mathbf{l}^{\mu} = \mathbf{v_1^{\mu}}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^{\mu}}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^{\mu}} = \frac{\delta^{\mu \mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}}, \;\; \mathbf{v_2^{\mu}} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1},\mathbf{2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\mu}}\mathbf{q_i^{\nu}} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^{\nu}}\,\mathbf{q_i^{\mu}}$$

$$\mathbf{2}(\mathbf{d_0} + \mathbf{m_0^2}) = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1},\mathbf{2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w})$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;= \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;= \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;\; = \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

Example 1: **Reduction of triangle scalar integrand to bubble integrands in D=2 dimension**

$$\mathbf{I_3} = \int \frac{\mathrm{d}^2\mathbf{l}}{(2\pi)^2}\,\mathcal{I}_3 \;\;,\;\; \mathcal{I}_3 = \frac{1}{\mathbf{d_0 d_1 d_2}},\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]\,,\; \mathbf{d_i} = (\mathbf{l}+\mathbf{q_i})^2 - \mathbf{m_i^2},\, i \in [0,1,2]$$

$$\mathbf{l}^\mu = \mathbf{v_1^\mu}(\mathbf{l}\cdot\mathbf{q_1}) + \mathbf{v_2^\mu}(\mathbf{l}\cdot\mathbf{q_2})\,, \quad \mathbf{v_1^\mu} = \frac{\delta^{\mu\mathbf{q_2}}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},\;\; \mathbf{v_2^\mu} = \frac{\delta^{\mathbf{q_1}\mu}_{\mathbf{q_1 q_2}}}{\mathbf{\Delta_2}},$$

$$\mathbf{l}\cdot\mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}}(\mathbf{d_i} - \mathbf{d_0} - \mathbf{r_i})\,, \quad \mathbf{r_i} = \mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}\,, \quad \mathbf{i} = \mathbf{1,2}\,,$$

$$\mathbf{g}^{\mu\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\mu q_i^\nu} = \sum_{\mathbf{j=1}}^{\mathbf{2}} \mathbf{v_i^\nu q_i^\mu}$$

$$\mathbf{2(d_0 + m_0^2)} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})\mathbf{r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}_\mu \sum_{\mathbf{i=1}}^{\mathbf{2}} \mathbf{v_i^\mu r_i} = \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \mathbf{l}\cdot\mathbf{w}$$

$$= \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{v_i})(\mathbf{d_i} - \mathbf{d_0}) - \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{l}\cdot\mathbf{q_i})(\mathbf{v_i}\cdot\mathbf{w}) \;= \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{i=1}}^{\mathbf{2}} (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_i} - (\mathbf{2l}\cdot\mathbf{v_i} - \mathbf{w}\cdot\mathbf{v_i})\mathbf{d_0} + \frac{\mathbf{1}}{\mathbf{2}}\mathbf{w^2}$$

$$\frac{\mathbf{1}}{\mathbf{d_0 d_1 d_2}} = \frac{\mathbf{1}}{(\mathbf{4m_0^2 - w^2})} \left\{ \frac{\mathbf{2(l\cdot v_1) - (w\cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l\cdot v_2) - (w\cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4 + (2l - w)\cdot(v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{\mathbf{d_0 d_1 d_2}} = \frac{1}{(4\mathbf{m_0^2} - \mathbf{w^2})} \left\{ \frac{\mathbf{2(l \cdot v_1)} - \mathbf{(w \cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l \cdot v_2)} - \mathbf{(w \cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4} + \mathbf{(2l - w)} \cdot \mathbf{(v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{\mathbf{d_0 d_1 d_2}} = \frac{1}{(4\mathbf{m_0^2} - \mathbf{w^2})} \left\{ \frac{\mathbf{2(l \cdot v_1) - (w \cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l \cdot v_2) - (w \cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4 + (2l - w) \cdot (v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

In the right hand side the terms proportional to $\mathbf{l}_\mu$ depend only on $\mathbf{l}_\mathbf{T}^\mu$ and integrate to zero.

The inflow momenta for the three bubble denominators are different

$$\mathbf{q_2 \cdot v_1 = 0}\,,\ \mathbf{q_1 \cdot v_2 = 0}\,,\ \mathbf{(q_2 - q_1) \cdot (v_1 + v_2) = 0}$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{d_0 d_1 d_2} = \frac{1}{(4m_0^2 - w^2)} \left\{ \frac{2(l \cdot v_1) - (w \cdot v_1)}{d_0 d_2} + \frac{2(l \cdot v_2) - (w \cdot v_2)}{d_0 d_1} - \frac{4 + (2l - w) \cdot (v_1 + v_2)}{d_1 d_2} \right\}.$$

In the right hand side the terms proportional to l_μ depend only on l_T^μ and integrate to zero.

The inflow momenta for the three bubble denominators are different

$$q_2 \cdot v_1 = 0\,,\ q_1 \cdot v_2 = 0\,,\ (q_2 - q_1) \cdot (v_1 + v_2) = 0$$

$$I_{3;012} = \frac{(-1)}{(4m_0^2 - w^2)} \{(w \cdot v_1) I_{2;02} + (w \cdot v_2) I_{2;01} + (4 - w \cdot (v_1 + v_2)) I_{2;12}\}$$

Reduction of triangle scalar integrand to bubble integrands in D=2 dimension

$$\frac{1}{\mathbf{d_0 d_1 d_2}} = \frac{1}{(\mathbf{4m_0^2 - w^2})} \left\{ \frac{\mathbf{2(l \cdot v_1) - (w \cdot v_1)}}{\mathbf{d_0 d_2}} + \frac{\mathbf{2(l \cdot v_2) - (w \cdot v_2)}}{\mathbf{d_0 d_1}} - \frac{\mathbf{4 + (2l - w) \cdot (v_1 + v_2)}}{\mathbf{d_1 d_2}} \right\}.$$

In the right hand side the terms proportional to $\mathbf{l}_\mu$ depend only on $\mathbf{l}_\mathbf{T}^\mu$ and integrate to zero.

The inflow momenta for the three bubble denominators are different

$$\mathbf{q_2 \cdot v_1 = 0\,,\ q_1 \cdot v_2 = 0\,,\ (q_2 - q_1) \cdot (v_1 + v_2) = 0}$$

$$\mathbf{I_{3;012} = \frac{(-1)}{(4m_0^2 - w^2)} \{(w \cdot v_1) I_{2;02} + (w \cdot v_2) I_{2;01} + (4 - w \cdot (v_1 + v_2)) I_{2;12}\}}$$

Exercise:

Any N-point scalar one-loop integrand function, for N > D, where D is the dimensionality of space-time, can be written as a linear combination of the D-point scalar and vector integrand functions.

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^{\mathbf{2}}}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2 - m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l + k})^{\mathbf{2}} - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^{\mathbf{2}} = \mathbf{1}$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^{\mathbf{2}}}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2 - m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l + k})^{\mathbf{2}} - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^{\mathbf{2}} = \mathbf{1}$$

Our aim is to reduce it using VN method.

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^{\mathbf{2}}}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1 = l^2 - m_1^2}\,,\ \mathbf{d_2 = (l + k)^2 - m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^{\mathbf{2}} = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})}, \qquad (\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})},$$

$$(\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

$$\mathbf{l}^\mu = (\mathbf{l} \cdot \mathbf{n})\mathbf{n}^\mu + (\mathbf{l} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}^\mu + \mathbf{n}^\mu_\epsilon (\mathbf{l} \cdot \mathbf{n}_\epsilon)$$

$$\mathbf{l}^2_{(\mathbf{2})} = \mathbf{d_1} + \mathbf{m_1^2} - \mu^2, \quad \mathbf{2\, l \cdot k} = \mathbf{d_2} - \mathbf{d_1} - \mathbf{r_1^2},$$

$$\mu^2 = (\mathbf{n}_\epsilon \cdot \mathbf{l})^2$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})},$$

$$(\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

$$\mathbf{l}^\mu = (\mathbf{l} \cdot \mathbf{n})\mathbf{n}^\mu + (\mathbf{l} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}^\mu + \mathbf{n}^\mu_\epsilon (\mathbf{l} \cdot \mathbf{n}_\epsilon)$$

$$\mathbf{l}^2_{(\mathbf{2})} = \mathbf{d_1} + \mathbf{m_1^2} - \mu^2, \quad \mathbf{2\, l \cdot k} = \mathbf{d_2} - \mathbf{d_1} - \mathbf{r_1^2},$$

$$\mu^2 = (\mathbf{n}_\epsilon \cdot \mathbf{l})^2$$

Result of the reduction:

$$\frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} = -\frac{(\lambda^2 + \mu^2)}{\mathbf{d_1 d_2}} + \frac{\mathbf{1}}{\mathbf{4k^2}} \left[\frac{\mathbf{r_1^2} - \mathbf{2\, l \cdot k}}{\mathbf{d_1}} + \frac{\mathbf{r_2^2} + \mathbf{2\, l \cdot k} + \mathbf{2k^2}}{\mathbf{d_2}} \right].$$

$$\mathbf{r_1^2} = \mathbf{k^2} + \mathbf{m_1^2} - \mathbf{m_2^2}, \quad \mathbf{r_2^2} = \mathbf{k^2} + \mathbf{m_2^2} - \mathbf{m_1^2},$$

$$\lambda^2 = \frac{\mathbf{k^4} - \mathbf{2k^2}(\mathbf{m_1^2} + \mathbf{m_2^2}) + (\mathbf{m_1^2} - \mathbf{m_2^2})^2}{\mathbf{4k^2}}$$

Example 2.: **Reduction of a rank-two two point function in** $\mathbf{D = 2 - 2\epsilon}$

$$\mathcal{I}(\mathbf{k}, \mathbf{m_1}, \mathbf{m_2}) = \frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}}, \qquad \mathbf{d_1} = \mathbf{l^2} - \mathbf{m_1^2}\,,\ \mathbf{d_2} = (\mathbf{l} + \mathbf{k})^2 - \mathbf{m_2^2}\,,\ \hat{\mathbf{n}} \cdot \mathbf{k} = \mathbf{0}, \mathbf{k^2} \neq \mathbf{0}\,,\ \text{and}\ \hat{\mathbf{n}}^2 = \mathbf{1}$$

Our aim is to reduce it using VN method.

The integral is UV divergent. We regularize it dimensionally $\mathbf{D = 2 - 2\epsilon}$

$$\mathbf{n}^\mu = \frac{\mathbf{k}^\mu}{\sqrt{\mathbf{k^2}}}, \quad \mathbf{n^2} = \mathbf{1},\ \mathbf{n}^\mu \mathbf{n}^\nu + \hat{\mathbf{n}}^\mu \hat{\mathbf{n}}^\nu = \mathbf{g}^{\mu\nu}_{(\mathbf{2})},$$

$$(\hat{\mathbf{n}} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - (\mathbf{n} \cdot \mathbf{l})^2 = \mathbf{l}^2_{(\mathbf{2})} - \frac{(\mathbf{l} \cdot \mathbf{k})^2}{\mathbf{k^2}}$$

$$\mathbf{l}^\mu = (\mathbf{l} \cdot \mathbf{n})\mathbf{n}^\mu + (\mathbf{l} \cdot \hat{\mathbf{n}})\hat{\mathbf{n}}^\mu + \mathbf{n}^\mu_\epsilon (\mathbf{l} \cdot \mathbf{n}_\epsilon)$$

$$\mathbf{l}^2_{(\mathbf{2})} = \mathbf{d_1} + \mathbf{m_1^2} - \mu^2, \quad \mathbf{2\, l \cdot k} = \mathbf{d_2} - \mathbf{d_1} - \mathbf{r_1^2},$$

$$\mu^2 = (\mathbf{n}_\epsilon \cdot \mathbf{l})^2$$

Result of the reduction:

$$\frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} = -\frac{(\lambda^2 + \mu^2)}{\mathbf{d_1 d_2}} + \frac{\mathbf{1}}{\mathbf{4k^2}} \left[\frac{\mathbf{r_1^2} - \mathbf{2\, l \cdot k}}{\mathbf{d_1}} + \frac{\mathbf{r_2^2} + \mathbf{2\, l \cdot k} + \mathbf{2k^2}}{\mathbf{d_2}} \right].$$

$$\mathbf{r_1^2} = \mathbf{k^2} + \mathbf{m_1^2} - \mathbf{m_2^2}, \quad \mathbf{r_2^2} = \mathbf{k^2} + \mathbf{m_2^2} - \mathbf{m_1^2},$$

$$\lambda^2 = \frac{\mathbf{k^4} - \mathbf{2k^2}(\mathbf{m_1^2} + \mathbf{m_2^2}) + (\mathbf{m_1^2} - \mathbf{m_2^2})^2}{\mathbf{4k^2}}$$

It follows the parametrization:

$$\frac{(\hat{\mathbf{n}} \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} = \frac{\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}} \cdot \mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon \cdot \mathbf{l})^2}{\mathbf{d_1 d_2}} + \frac{\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n} \cdot \mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}} \cdot \mathbf{l})}{\mathbf{d_1}} + \frac{\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n} \cdot \mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}} \cdot \mathbf{l})}{\mathbf{d_2}}.$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^{\mathbf{2}} = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^{\mathbf{2}}\right] + \left[\mathbf{a}_{1,0} + \mathbf{a}_{1,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{1,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a}_{2,0} + \mathbf{a}_{2,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{2,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a}_{1,0} + \mathbf{a}_{1,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{1,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a}_{2,0} + \mathbf{a}_{2,1}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a}_{2,2}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1} + \mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0} + \mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}} + \mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}} + \mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l}) + \mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1} + \mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1}+\mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

$$\frac{(\hat{\mathbf{n}}\cdot\mathbf{l_1})^\mathbf{2}}{\mathbf{d_2(l_1)}} - \frac{\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l_1})^\mathbf{2}}{\mathbf{d_2(l_1)}} = \mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \frac{\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})}{\mathbf{d_2(l_1)}}\mathbf{d_1(l_1)}$$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1}+\mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

$$\frac{(\hat{\mathbf{n}}\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} - \frac{\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} = \mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \frac{\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})}{\mathbf{d_2(l_1)}}\mathbf{d_1(l_1)}$$

$$\mathbf{l_1} = \gamma_\mathbf{1}\mathbf{n} + \gamma_\mathbf{2}\hat{\mathbf{n}}\,, \quad \gamma_\mathbf{1}^\mathbf{2} + \gamma_\mathbf{2}^\mathbf{2} = \mathbf{m_1^2}$$

Choose $\gamma_\mathbf{1} = \mathbf{0}, \gamma_\mathbf{2} = \pm\mathbf{m_1}$

Choose $\gamma_\mathbf{2} = \mathbf{0}, \gamma_\mathbf{1} = \mathbf{m_1}$

Connection to “unitarity” : evaluate coefficients with “on-shell” loop momenta

$$(\hat{\mathbf{n}}\cdot\mathbf{l})^2 = \left[\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l})^2\right] + \left[\mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_2} \quad + \left[\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})\right]\mathbf{d_1}$$

Calculate $\mathbf{b_0}, \mathbf{b_1}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{n}_\epsilon\cdot\mathbf{l} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\hat{\mathbf{n}}$$

$$\mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^+} = -\lambda^\mathbf{2}, \quad \mathbf{b_0} + \mathbf{b_1}\hat{\mathbf{n}}\cdot\mathbf{l_c^-} = -\lambda^\mathbf{2}$$

$$\alpha_\mathbf{c} = -\frac{\mathbf{r_1^2}}{\mathbf{2}\sqrt{\mathbf{k^2}}}, \quad \beta_\mathbf{c} = \lambda$$

$$\mathbf{b_0} = -\lambda^\mathbf{2}\,, \ \mathbf{b_1} = \mathbf{0}$$

Calculate $\mathbf{b_2}$ assuming $\mathbf{d_1(l)} = \mathbf{d_2(l)} = \mathbf{0}$ and $\mathbf{l}\cdot\hat{\mathbf{n}} = \mathbf{0}$:

$$\mathbf{l_c^\pm} = \alpha_\mathbf{c}\mathbf{n} \pm \mathbf{i}\beta_\mathbf{c}\mathbf{n}_\epsilon$$

$$\mathbf{0} = (\mathbf{1}+\mathbf{b_2})\lambda^\mathbf{2}$$

$$\mathbf{b_2} = -\mathbf{1}$$

Calculate tadpole coefficients $\mathbf{a_{1,0}}.\mathbf{a_{1,1}}, \mathbf{a_{1,2}}$ assuming $\mathbf{d_1(l)} = \mathbf{0}$:

$$\frac{(\hat{\mathbf{n}}\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} - \frac{\mathbf{b_0}+\mathbf{b_1}(\hat{\mathbf{n}}\cdot\mathbf{l})+\mathbf{b_2}(\mathbf{n}_\epsilon\cdot\mathbf{l_1})^2}{\mathbf{d_2(l_1)}} = \mathbf{a_{1,0}}+\mathbf{a_{1,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{1,2}}(\hat{\mathbf{n}}\cdot\mathbf{l}) + \frac{\mathbf{a_{2,0}}+\mathbf{a_{2,1}}(\mathbf{n}\cdot\mathbf{l})+\mathbf{a_{2,2}}(\hat{\mathbf{n}}\cdot\mathbf{l})}{\mathbf{d_2(l_1)}}\mathbf{d_1(l_1)}$$

$$\mathbf{l_1} = \gamma_\mathbf{1}\mathbf{n} + \gamma_\mathbf{2}\hat{\mathbf{n}}\,, \quad \gamma_\mathbf{1}^\mathbf{2} + \gamma_\mathbf{2}^\mathbf{2} = \mathbf{m_1^2}$$

$$\mathbf{a_{1,0}} \pm \mathbf{a_{1,2}}\mathbf{m_1} = \frac{\mathbf{m_1^2}+\lambda^\mathbf{2}}{\mathbf{r_1^2}} = \frac{\mathbf{r_1^2}}{\mathbf{4k^2}}$$

Choose $\gamma_\mathbf{1} = \mathbf{0}, \gamma_\mathbf{2} = \pm\mathbf{m_1}$

$$\mathbf{a_{1,2}} = \mathbf{0}\,, \mathbf{a_{1,0}} = \mathbf{r_1^2}/(\mathbf{4k^2})\,,$$

Choose $\gamma_\mathbf{2} = \mathbf{0}, \gamma_\mathbf{1} = \mathbf{m_1}$

$$\mathbf{a_{1,1}} = -(\mathbf{4k^2})^{\mathbf{-1/2}}$$

OPP reduction *(Ossola, Papadopoulos, Pittau)*

First we made direct NV reduction of the loop integrand.

Next we have pointed out a generic parametric form of the loop integrand and fitted the parameters with the help of on-shell values of the loop momenta solving first double cut and single cut conditions (iteratively).

The loop integration can be easily carried out:

$$\mathbf{I(k, m_1^2, m_2^2)} = \int \frac{\mathbf{d^{2-2\epsilon}}}{\mathbf{i(2\pi)^{2-2\epsilon}}} \mathcal{I}(\mathbf{k, m_1, m_2}) = \mathbf{b_0\, I_2(k^2, m_1^2, m_2^2) + a_{1,0}\, I_1(m_1^2) + a_{2,0}\, I_1(m_2^2) + \frac{b_2}{4\pi}}$$

Exercise:

Calculate the photon self-energy in D=2 QED (Scwinger-model) using NV reduction at the integrand level.

OPP reduction: general case

OPP reduction: general case

✦ **Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.**

OPP reduction: general case

- Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.

- "Integrand of a diagram has fully ordered external legs:
  Ordered amplitudes given as sums of diagrams.
  N different l-dependent scalar propagators. Momentum inflow to the loop.

  This gives unique prescription of the integrand function as a function of the loop momentum modulo overall shift.

OPP reduction: general case

✦ Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.

✦ "Integrand of a diagram has fully ordered external legs:
Ordered amplitudes given as sums of diagrams.
N different l-dependent scalar propagators. Momentum inflow to the loop.

This gives unique prescription of the integrand function as a function of the loop momentum modulo overall shift.



OPP reduction: general case

✦ Parametric integral over the loop momentum. Any integrand is decomposed in terms a few known functions.

✦ "Integrand of a diagram has fully ordered external legs:
Ordered amplitudes given as sums of diagrams.
N different l-dependent scalar propagators. Momentum inflow to the loop.

This gives unique prescription of the integrand function as a function of the loop momentum modulo overall shift.

✦ For 4D external kinematics, the integrand of any one-loop Feynman amplitude with arbitrary number of external legs can always be written in linear combination of penta, quadru-,triple-,double- and single-pole terms



Ordered amplitudes have well defined integrand

$$\mathcal{I}_N(p_1,p_2,\ldots,p_N,l)=\frac{\mathcal{N}(p_1,p_2,\ldots,p_N;l)}{d_1 d_2\cdots d_N}$$



$$\mathcal{I}_N(p_1,p_2,\ldots,p_N,l)$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms





The number of terms with k denominators is $\binom{\mathbf{N}}{\mathbf{k}}$







The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d^D l}}{(2\pi)^{\mathrm{D}}} \frac{\mathrm{Num(l)}}{\prod_{\mathrm{i}} \mathrm{d_i(l)}} = \int \frac{\mathrm{d^D l}}{(2\pi)^{\mathrm{D}}} \frac{1}{\prod_{\mathrm{i}} \mathrm{d_i(l)}} \times \Bigg\{ \sum_{\mathrm{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathrm{e}}_{\mathrm{i_1,i_2,i_3,i_4,i_5}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2,i_3,i_4,i_5}]} \mathrm{d_j(l)}$$

$$+ \sum_{\mathrm{i_1,i_2,i_3,i_4}} \tilde{\mathrm{d}}_{\mathrm{i_1,i_2,i_3,i_4}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2,i_3,i_4}]} \mathrm{d_j(l)}$$

$$+ \sum_{\mathrm{i_1,i_2,i_3}} \tilde{\mathrm{c}}_{\mathrm{i_1,i_2,i_3}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2,i_3}]} \mathrm{d_j(l)} + \sum_{\mathrm{i_1,i_2}} \tilde{\mathrm{b}}_{\mathrm{i_1,i_2}}(\mathrm{l}) \prod_{\mathrm{j} \neq [\mathrm{i_1,i_2}]} \mathrm{d_j(l)} + \sum_{\mathrm{i_1}} \tilde{\mathrm{a}}_{\mathrm{i_1}}(\mathrm{l}) \prod_{\mathrm{j} \neq \mathrm{i_1}} \mathrm{d_j(l)} \Bigg\}.$$

$$\mathbf{d_i(l) = (l + q_i)^2 - m_i^2 \, , i = 0, \ldots, 4 \, , \; q_0 = 0}$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathbf{1}}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} \times \Bigg\{ \sum_{\mathbf{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathbf{e}}_{\mathbf{i_1,i_2,i_3,i_4,i_5}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4,i_5]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3,i_4}} \tilde{\mathbf{d}}_{\mathbf{i_1,i_2,i_3,i_4}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3}} \tilde{\mathbf{c}}_{\mathbf{i_1,i_2,i_3}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1,i_2}} \tilde{\mathbf{b}}_{\mathbf{i_1,i_2}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}_{\mathbf{i_1}}(\mathbf{l}) \prod_{\mathbf{j \neq i_1}} \mathbf{d_j}(\mathbf{l}) \Bigg\}.$$

$$\mathbf{d_i}(\mathbf{l}) = (\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}\,, \mathbf{i} = \mathbf{0}, \ldots, \mathbf{4}\,, \;\; \mathbf{q_0} = \mathbf{0}$$

$$\mathrm{Num}(\mathbf{l}) = \mathbf{N_5}(\mathbf{l}) = \prod_{\mathbf{i=1}}^{\mathbf{5}} \mathbf{u_i} \cdot \mathbf{l}, \qquad \mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{4}} (\mathbf{l} \cdot \mathbf{q_i}) \mathbf{v_i^{\mu}} + (\mathbf{l} \cdot \mathbf{n}_{\epsilon}) \mathbf{n}_{\epsilon}^{\mu} \quad \mathbf{l} \cdot \mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left(\mathbf{d_i} - \mathbf{d_0} - (\mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}) \right)$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathbf{1}}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} \times \Bigg\{ \sum_{\mathbf{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathbf{e}}_{\mathbf{i_1,i_2,i_3,i_4,i_5}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4,i_5]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3,i_4}} \tilde{\mathbf{d}}_{\mathbf{i_1,i_2,i_3,i_4}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3}} \tilde{\mathbf{c}}_{\mathbf{i_1,i_2,i_3}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1,i_2}} \tilde{\mathbf{b}}_{\mathbf{i_1,i_2}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}_{\mathbf{i_1}}(\mathbf{l}) \prod_{\mathbf{j \neq i_1}} \mathbf{d_j}(\mathbf{l}) \Bigg\}.$$

$$\mathbf{d_i}(\mathbf{l}) = (\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}\,, \mathbf{i} = \mathbf{0}, \ldots, \mathbf{4}\,, \;\; \mathbf{q_0} = \mathbf{0}$$

$$\mathrm{Num}(\mathbf{l}) = \mathbf{N_5}(\mathbf{l}) = \prod_{\mathbf{i=1}}^{\mathbf{5}} \mathbf{u_i} \cdot \mathbf{l}, \qquad \mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{4}} (\mathbf{l} \cdot \mathbf{q_i}) \mathbf{v_i^{\mu}} + (\mathbf{l} \cdot \mathbf{n}_{\epsilon}) \mathbf{n}_{\epsilon}^{\mu} \quad \mathbf{l} \cdot \mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left(\mathbf{d_i} - \mathbf{d_0} - (\mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}) \right)$$

$$\mathbf{N_5}(\mathbf{l}) = \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{u_5} \cdot \mathbf{l}) = \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{d_j} - \mathbf{d_0}) \; - \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{q_j^2} - \mathbf{m_j^2} + \mathbf{m_0^2})$$

The integrand can be decomposed to pentagon, box, triangle, bubble and tadpole terms

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2\pi})^{\mathbf{D}}} \frac{\mathbf{1}}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} \times \Bigg\{ \sum_{\mathbf{i_1,i_2,i_3,i_4,i_5}} \tilde{\mathbf{e}}_{\mathbf{i_1,i_2,i_3,i_4,i_5}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4,i_5]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3,i_4}} \tilde{\mathbf{d}}_{\mathbf{i_1,i_2,i_3,i_4}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3,i_4]}} \mathbf{d_j}(\mathbf{l})$$

$$+ \sum_{\mathbf{i_1,i_2,i_3}} \tilde{\mathbf{c}}_{\mathbf{i_1,i_2,i_3}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2,i_3]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1,i_2}} \tilde{\mathbf{b}}_{\mathbf{i_1,i_2}}(\mathbf{l}) \prod_{\mathbf{j \neq [i_1,i_2]}} \mathbf{d_j}(\mathbf{l}) + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}_{\mathbf{i_1}}(\mathbf{l}) \prod_{\mathbf{j \neq i_1}} \mathbf{d_j}(\mathbf{l}) \Bigg\}.$$

$$\mathbf{d_i}(\mathbf{l}) = (\mathbf{l} + \mathbf{q_i})^{\mathbf{2}} - \mathbf{m_i^2}\,, \mathbf{i} = \mathbf{0}, \ldots, \mathbf{4}\,, \;\; \mathbf{q_0} = \mathbf{0}$$

$$\mathrm{Num}(\mathbf{l}) = \mathbf{N_5}(\mathbf{l}) = \prod_{\mathbf{i=1}}^{\mathbf{5}} \mathbf{u_i} \cdot \mathbf{l}, \qquad \mathbf{l}^{\mu} = \sum_{\mathbf{i=1}}^{\mathbf{4}} (\mathbf{l} \cdot \mathbf{q_i}) \mathbf{v_i^{\mu}} + (\mathbf{l} \cdot \mathbf{n}_{\epsilon}) \mathbf{n}_{\epsilon}^{\mu} \quad \mathbf{l} \cdot \mathbf{q_i} = \frac{\mathbf{1}}{\mathbf{2}} \left(\mathbf{d_i} - \mathbf{d_0} - (\mathbf{q_i^2} - \mathbf{m_i^2} + \mathbf{m_0^2}) \right)$$

$$\mathbf{N_5}(\mathbf{l}) = \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{u_5} \cdot \mathbf{l}) = \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{d_j} - \mathbf{d_0}) \; - \frac{\mathbf{1}}{\mathbf{2}} \sum_{\mathbf{j=1}}^{\mathbf{4}} (\mathbf{u_5} \cdot \mathbf{v_j}) \left(\prod_{\mathbf{i}}^{\mathbf{4}} \mathbf{u_i} \cdot \mathbf{l} \right) (\mathbf{q_j^2} - \mathbf{m_j^2} + \mathbf{m_0^2})$$

$$\tilde{\mathbf{e}}_{\mathbf{01234}}(\mathbf{l}) = \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{01234}}$$

Parameter counting

The numerators are simple polynomials of the loop momentum components of the corresponding trivial space.

For a given 5,-4-,3-,2-,1 denominator 1, 5, 10, 10, 5 parameters;

Pentagon (rank five): $\mathbf{D_P = 4\,, D_T = 0 + 1\,, l_T^2 = (ln_\epsilon)^2} = \text{constant terms} + \mathcal{O}(\mathbf{d_i})$

Box (rank four): $\mathbf{D_P = 3\,, D_T = 1 + 1\,, l_T^2 = (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Triangle (rank three): $\mathbf{D_P = 2\,, D_T = 3 + 1\,, l_T^2 = (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Bubble (rank two): $\mathbf{D_P = 1\,, D_T = 3 + 1\,, l_T^2 = (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Tadpole (rank one): $\mathbf{D_P = 0\,, D_T = 4 + 1\,, l_T^2 = (ln_1)^2 + (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Parameter counting

The numerators are simple polynomials of the loop momentum components of the corresponding trivial space.

For a given 5,-4-,3-,2-,1 denominator 1, 5, 10, 10, 5 parameters; $l_T^2 = l^2 - l_P^2\,, \quad l_P^\mu = (lq_i)v_i^\mu$

Pentagon (rank five): $\mathbf{D_P = 4\,, D_T = 0+1\,, l_T^2 = (ln_\epsilon)^2} = \text{constant terms} + \mathcal{O}(\mathbf{d_i})$

Box (rank four): $\mathbf{D_P = 3\,, D_T = 1+1\,, l_T^2 = (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Triangle (rank three): $\mathbf{D_P = 2\,, D_T = 3+1\,, l_T^2 = (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Bubble (rank two): $\mathbf{D_P = 1\,, D_T = 3+1\,, l_T^2 = (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Tadpole (rank one): $\mathbf{D_P = 0\,, D_T = 4+1\,, l_T^2 = (ln_1)^2 + (ln_2)^2 + (ln_3)^2 + (ln_4)^2 + (ln_\epsilon)^2} = \text{const} + \mathcal{O}(\mathbf{d_i})$

Parameter counting

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

Parameter counting

$$\tilde{\mathbf{e}}_{\mathbf{01234}}(\mathbf{l}) = \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{01234}}$$

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathbf{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

Parameter counting

$$\tilde{\mathbf{e}}_{\mathbf{01234}}(\mathbf{l}) = \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{01234}}$$

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathbf{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\begin{aligned}\tilde{\mathbf{c}}_{\mathbf{012}}(\mathbf{l}) = \tilde{\mathbf{c}}_{\mathbf{0}} &+ \tilde{\mathbf{c}}_{\mathbf{1}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}}) + \tilde{\mathbf{c}}_{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{c}}_{\mathbf{3}}((\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}})^{\mathbf{2}} - (\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}})^{\mathbf{2}}) + \tilde{\mathbf{c}}_{\mathbf{4}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}})(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{c}}_{\mathbf{5}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}})^{\mathbf{3}} \\ &+ \tilde{\mathbf{c}}_{\mathbf{6}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}})^{\mathbf{3}} + \tilde{\mathbf{c}}_{\mathbf{7}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{c}}_{\mathbf{8}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{3}}) + \tilde{\mathbf{c}}_{\mathbf{9}}(\mathbf{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathbf{l}\cdot\mathbf{n}_{\mathbf{4}})\end{aligned}$$

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

$$\tilde{d}_{0123}(l) = \tilde{d}_0 + \tilde{d}_1(l\cdot n_4) + \tilde{d}_2(l\cdot n_\epsilon)^2 + \tilde{d}_3(l\cdot n_\epsilon)^2(l\cdot n_4) + \tilde{d}_4(l\cdot n_\epsilon)^4,$$

$$\begin{aligned}\tilde{c}_{012}(l) = \tilde{c}_0 &+ \tilde{c}_1(l\cdot n_3) + \tilde{c}_2(l\cdot n_4) + \tilde{c}_3((l\cdot n_3)^2 - (l\cdot n_4)^2) + \tilde{c}_4(l\cdot n_3)(l\cdot n_4) + \tilde{c}_5(l\cdot n_3)^3 \\ &+ \tilde{c}_6(l\cdot n_4)^3 + \tilde{c}_7(l\cdot n_\epsilon)^2 + \tilde{c}_8(l\cdot n_\epsilon)^2(l\cdot n_3) + \tilde{c}_9(l\cdot n_\epsilon)^2(l\cdot n_4)\end{aligned}$$

$$\begin{aligned}\tilde{b}_{01}(l) = \tilde{b}_0 &+ \tilde{b}_1(l\cdot n_2) + \tilde{b}_2(l\cdot n_3) + \tilde{b}_3(l\cdot n_4) + \tilde{b}_4((l\cdot n_2)^2 - (l\cdot n_4)^2) + \tilde{b}_5((l\cdot n_3)^2 - (l\cdot n_4)^2) \\ &+ \tilde{b}_6(l\cdot n_2)(l\cdot n_3) + \tilde{b}_7(l\cdot n_3)(l\cdot n_4) + \tilde{b}_8(l\cdot n_2)(l\cdot n_4) + \tilde{b}_9(l\cdot n_\epsilon)^2,\end{aligned}$$

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

$$\tilde{d}_{0123}(l) = \tilde{d}_0 + \tilde{d}_1(l\cdot n_4) + \tilde{d}_2(l\cdot n_\epsilon)^2 + \tilde{d}_3(l\cdot n_\epsilon)^2(l\cdot n_4) + \tilde{d}_4(l\cdot n_\epsilon)^4,$$

$$\begin{aligned}\tilde{c}_{012}(l) = \tilde{c}_0 &+ \tilde{c}_1(l\cdot n_3) + \tilde{c}_2(l\cdot n_4) + \tilde{c}_3((l\cdot n_3)^2 - (l\cdot n_4)^2) + \tilde{c}_4(l\cdot n_3)(l\cdot n_4) + \tilde{c}_5(l\cdot n_3)^3 \\ &+ \tilde{c}_6(l\cdot n_4)^3 + \tilde{c}_7(l\cdot n_\epsilon)^2 + \tilde{c}_8(l\cdot n_\epsilon)^2(l\cdot n_3) + \tilde{c}_9(l\cdot n_\epsilon)^2(l\cdot n_4)\end{aligned}$$

$$\begin{aligned}\tilde{b}_{01}(l) = \tilde{b}_0 &+ \tilde{b}_1(l\cdot n_2) + \tilde{b}_2(l\cdot n_3) + \tilde{b}_3(l\cdot n_4) + \tilde{b}_4((l\cdot n_2)^2 - (l\cdot n_4)^2) + \tilde{b}_5((l\cdot n_3)^2 - (l\cdot n_4)^2) \\ &+ \tilde{b}_6(l\cdot n_2)(l\cdot n_3) + \tilde{b}_7(l\cdot n_3)(l\cdot n_4) + \tilde{b}_8(l\cdot n_2)(l\cdot n_4) + \tilde{b}_9(l\cdot n_\epsilon)^2,\end{aligned}$$

$$\tilde{a}_i(l) = \tilde{a}_0 + \tilde{a}_1(l\cdot n_1) + \tilde{a}_2(l\cdot n_2) + \tilde{a}_3(l\cdot n_3) + \tilde{a}_4(l\cdot n_4)$$

Parameter counting

$$\tilde{e}_{01234}(l) = \tilde{e}^{(0)}_{01234}$$

$$\tilde{d}_{0123}(l) = \tilde{d}_0 + \tilde{d}_1(l\cdot n_4) + \tilde{d}_2(l\cdot n_\epsilon)^2 + \tilde{d}_3(l\cdot n_\epsilon)^2(l\cdot n_4) + \tilde{d}_4(l\cdot n_\epsilon)^4,$$

$$\begin{aligned}\tilde{c}_{012}(l) = \tilde{c}_0 &+ \tilde{c}_1(l\cdot n_3) + \tilde{c}_2(l\cdot n_4) + \tilde{c}_3((l\cdot n_3)^2 - (l\cdot n_4)^2) + \tilde{c}_4(l\cdot n_3)(l\cdot n_4) + \tilde{c}_5(l\cdot n_3)^3 \\ &+ \tilde{c}_6(l\cdot n_4)^3 + \tilde{c}_7(l\cdot n_\epsilon)^2 + \tilde{c}_8(l\cdot n_\epsilon)^2(l\cdot n_3) + \tilde{c}_9(l\cdot n_\epsilon)^2(l\cdot n_4)\end{aligned}$$

$$\begin{aligned}\tilde{b}_{01}(l) = \tilde{b}_0 &+ \tilde{b}_1(l\cdot n_2) + \tilde{b}_2(l\cdot n_3) + \tilde{b}_3(l\cdot n_4) + \tilde{b}_4((l\cdot n_2)^2 - (l\cdot n_4)^2) + \tilde{b}_5((l\cdot n_3)^2 - (l\cdot n_4)^2) \\ &+ \tilde{b}_6(l\cdot n_2)(l\cdot n_3) + \tilde{b}_7(l\cdot n_3)(l\cdot n_4) + \tilde{b}_8(l\cdot n_2)(l\cdot n_4) + \tilde{b}_9(l\cdot n_\epsilon)^2,\end{aligned}$$

$$\tilde{a}_i(l) = \tilde{a}_0 + \tilde{a}_1(l\cdot n_1) + \tilde{a}_2(l\cdot n_2) + \tilde{a}_3(l\cdot n_3) + \tilde{a}_4(l\cdot n_4)$$

The coefficients $\tilde{a}_0, \ldots, \tilde{e}_{01234}$ are independent from the loop momenta and in all numerator functions we can replace $(l\cdot n_i)$ with $(l_T\cdot n_i)$

Note that the integration over the transverse space is trivial

$$\int d^{D_1}l_\perp \delta(l_\perp^2 - \mu_0^2)\,(l_\perp^\mu, l_\perp^\mu l_\perp^\nu) = \int d^{D_1}l_\perp \delta(l_\perp^2 - \mu_0^2)\left(0,\ \frac{g_\perp^{\mu\nu}}{D_1}\, l_\perp^2\right), \qquad D_1 = D - 1$$

Only the constant terms and some of the terms depending on $(\mathbf{l} \cdot \mathbf{n}_\epsilon)^2$ give non-vanishing integrals

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}\mathbf{i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}\mathbf{i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}\mathbf{i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1}\mathbf{i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

Only the constant terms and some of the terms depending on $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ give non-vanishing integrals

$$\mathbf{I_N} = \int \frac{\mathrm{d}^{\mathbf{D}}\mathbf{l}}{(\mathbf{2}\pi)^{\mathbf{D}}} \frac{\mathrm{Num}(\mathbf{l})}{\prod_{\mathbf{i}} \mathbf{d_i}(\mathbf{l})} = \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \tilde{\mathbf{e}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4},\mathbf{i_5}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4 i_5}}$$
$$+ \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \tilde{\mathbf{d}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \mathbf{I}_{\mathbf{i_1 i_2 i_3 i_4}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \tilde{\mathbf{c}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \mathbf{I}_{\mathbf{i_1 i_2 i_3}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \tilde{\mathbf{b}}^{(\mathbf{0})}_{\mathbf{i_1},\mathbf{i_2}} \mathbf{I}_{\mathbf{i_1 i_2}} + \sum_{\mathbf{i_1}} \tilde{\mathbf{a}}^{(\mathbf{0})}_{\mathbf{i_1}} \mathbf{I}_{\mathbf{i_1}} + \mathcal{R}$$

Non-vanishing master integrals with $(\mathbf{l}\cdot\mathbf{n}_\epsilon)^2$ factors in the numerator

$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = -\frac{D-4}{2} I^{D+2}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^4}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} = \frac{(D-2)(D-4)}{4} I^{D+4}_{i_1 i_2 i_3 i_4},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2} d_{i_3}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2 i_3},$$
$$\int \frac{\mathrm{d}^D l}{(i\pi)^{D/2}} \frac{s_e^2}{d_{i_1} d_{i_2}} = -\frac{(D-4)}{2} I^{D+2}_{i_1 i_2}.$$

$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3 i_4} = 0,$$
$$\lim_{D\to 4} \frac{(D-4)(D-2)}{4} I^{(D+4)}_{i_1 i_2 i_3 i_4} = -\frac{1}{3},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2 i_3} = \frac{1}{2},$$
$$\lim_{D\to 4} \frac{(D-4)}{2} I^{(D+2)}_{i_1 i_2} = -\frac{m^2_{i_1} + m^2_{i_2}}{2} + \frac{1}{6}(q_{i_1} - q_{i_2})^2$$

they contribute to the rational terms

$$\mathcal{R} = -\sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3},\mathbf{i_4}} \frac{\tilde{\mathbf{d}}^{(\mathbf{4})}_{\mathbf{i_1 i_2 i_3 i_4}}}{\mathbf{6}} + \sum_{\mathbf{i_1},\mathbf{i_2},\mathbf{i_3}} \frac{\tilde{\mathbf{c}}^{(\mathbf{7})}_{\mathbf{i_1 i_2 i_3}}}{\mathbf{2}} + \sum_{\mathbf{i_1},\mathbf{i_2}} \left[\frac{\mathbf{m^2_{i_1}} + \mathbf{m^2_{i_2}}}{\mathbf{2}} - \frac{(\mathbf{q_{i_1}} - \mathbf{q_{i_2}})^{\mathbf{2}}}{\mathbf{6}} \right] \tilde{\mathbf{b}}^{(\mathbf{9})}_{\mathbf{i_1 i_2}}.$$

Example: Projecting out individual (quadrupole) coefficients in D-dimension

$$\mathcal{I}_N(p_1,p_2,\ldots,p_N,l) = \frac{\mathcal{N}(p_1,p_2,\ldots,p_N;l)}{d_1 d_2 \cdots d_N} =$$

$$= \sum_{i_1\le i_2\le i_3\le i_4\le i_5\le n} \frac{\bar{e}_{i_1 i_2 i_3 i_4 i_5}(l)}{d_{i_1} d_{i_2} d_{i_3} d_{i_4} d_{i_5}} +$$

$$\sum_{1\le i_1<i_2<i_3<i_4\le N} \frac{\bar{d}_{i_1 i_2 i_3 i_4}(l)}{d_{i_1} d_{i_2} d_{i_3} d_{i_4}} + \sum_{1\le i_1<i_2<i_3\le N} \frac{\bar{c}_{i_1 i_2 i_3}(l)}{d_{i_1} d_{i_2} d_{i_3}} + \sum_{1\le i_1<i_2\le N} \frac{\bar{b}_{i_1 i_2}(l)}{d_{i_1} d_{i_2}} + \sum_{1\le i_1\le N} \frac{\bar{a}_{i_1}(l)}{d_{i_1}}$$

Denote:

$$\mathbf{RR(l) = Residuum_{0123}(Integrand)} \\ \mathbf{- \text{ pentagon contributions } = \tilde{d}_{0123}(l)}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathrm{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathrm{n}_\epsilon)^2 + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathrm{n}_\epsilon)^2(\mathrm{l}\cdot\mathrm{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathrm{n}_\epsilon)^4,$$

$$\mathrm{l}^\mu = \mathrm{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathrm{n}_4^\mu + \sin\phi\ \mathrm{n}_\epsilon^\mu), \qquad \mathrm{V}^\mu = -\frac{1}{2}\sum_{\mathrm{i}}^{3} \mathrm{v}_\mathrm{i}^\mu \left(\mathrm{q}_\mathrm{i}^2 - \mathrm{m}_\mathrm{i}^2 + \mathrm{m}_0^2\right),$$

- Choose $\sin\phi = 0, \cos\phi = \pm 1$ denote $\mathrm{l}_\pm^\mu = \mathrm{V}^\mu \pm \mathrm{l}_\perp \mathrm{n}_4^\mu,$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

- Choose $\mathrm{l}_{\epsilon}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

calculate the residuum of the integrand for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

- Choose $\mathrm{l}_{\epsilon}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\mathrm{l}_{\perp}^{\mathbf{2}} = \frac{\mathrm{Num}(\mathrm{l}_{\epsilon}) - \tilde{\mathbf{d}}_{\mathbf{0}}}{\mathrm{l}_{\perp}^{\mathbf{2}}}.$$

Comments on the rational part:

The origin is UV divergent tensor integrals. Reduction requires regularization.
After reduction: D-dimensional finite tensor integrals. In the limiting case D=4
they provide finite constants independent from the kinematics.

It may happen that the numerator has manifest dependence on (D-4) coming from polarization sum. These terms can only contribute if it appears in a term leading to UV divergent integrals. There are only few UV divergent one-loop Feynman diagrams even for large number of external particles (OPP).

Sophisticated recursion relations (BDK) as an option in Black Hat

Comment on N=4 sYM amplitudes

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

- N=1 sYM scattering amplitudes have no rational parts

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

- N=1 sYM scattering amplitudes have no rational parts

BDDK theorem (1994) for one loop amplitudes:

The maximum number of loop momentum in the numerator of Feynman-diagrams is reduced by one for N=1 and by four for N=4 .

Comment on N=4 sYM amplitudes

- N = 4 sYM scattering amplitudes are free from UV divergences.
  n-particle one loop amplitudes in N = 4 are built out of only boxes.
  No triangles, no bubbles, no rational parts

- N=1 sYM scattering amplitudes have no rational parts

BDDK theorem (1994) for one loop amplitudes:

The maximum number of loop momentum in the numerator of Feynman-diagrams is reduced by one for N=1 and by four for N=4 .

Excercise (not easy):

Find proper parametrization for the bubble numerator $\tilde{\mathrm{b}}_{\mathbf{01}}(\mathrm{l})$ in case of light-like inflow momentum.

Lecture 3: Unitarity method and amplitudes

Constraints from Unitarity: $M^\dagger - M = -iM^\dagger M$



$$-i\,\mathrm{Disc}\;A_4(1,2,3,4)\Big|_{s-\mathrm{cut}} = \int \frac{d^4p}{(2\pi)^4}\, 2\pi\delta^{(+)}(\ell_1^2 - m^2)\, 2\pi\delta^{(+)}(\ell_2^2 - m^2)$$
$$\times\, A_4^{\mathrm{tree}}(-\ell_1, 1, 2, \ell_2)\, A_4^{\mathrm{tree}}(-\ell_2, 3, 4, \ell_1)\,,$$

Imaginary part of NLO an amplitude is calculated from tree amplitudes.

- Non-linear relation, iterative in the coupling.
- Iterative in amplitudes. Building blocks are amplitudes and not Feynman diagrams
- Manifestly gauge invariant.

Unitarity and Cutkosky rule:

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s = (p_1 + p_2)^2, t = (p_1 + p_3)^2,}$

$$\mathbf{s = 4E^2_{cm}, \quad t = -\frac{(s - 4m^2)}{2}\,(1 - \cos\theta_{cms})}$$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity:

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,}$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^* \langle n|T|p_1,p_2\rangle}$$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed θ_{cms} ?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^* \langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed θ_{cms} ?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,}$ $\qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

More external particles?

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

More external particles?

In perturbation theory simple recipe to calculate discontinuities across branch cuts of multi-leg, multi-loop Feynman diagrams:

Unitarity and Cutkosky rule:

Scattering amplitudes of scalar particles are functions of the external momenta $\mathbf{A_N(\{p_i\})}$

In case of 2 to 2 scattering with equal masses the scattering amplitude $\mathbf{A_4(s,t)}$ depends on Lorenz-invariant Mandelstam-variables s,t , $\mathbf{s=(p_1+p_2)^2, t=(p_1+p_3)^2,}$

$$\mathbf{s=4E_{cm}^2,} \qquad \mathbf{t=-\frac{(s-4m^2)}{2}\,(1-\cos\theta_{cms})}$$

The scattering amplitude as analytic function of s at fixed $\theta_{\rm cms}$?

Unitarity: $\mathbf{S^\dagger S=1,\ S=1+iT,} \qquad \mathbf{2\,Im\,T=-i(T-T^\dagger)=T^\dagger T=TT^\dagger}$

$$\mathbf{2\,Im\,\langle p_3,p_4|T|p_1,p_2\rangle=\sum_n \langle n|T|p_3,p_4\rangle^*\langle n|T|p_1,p_2\rangle}$$

$$\mathbf{\langle p_3,p_4|T|p_1,p_2\rangle=(2\pi)^4\delta(p_1+p_2-p_3-p_4)M(s,t)}$$

As s increases :contributions form more and more intermediate states.

$\mathbf{M(s,t)}$ has branch cuts in the complex s-plane at $\mathbf{(nm)^2}$, $\mathbf{(n=2,3,\ldots)}$

More external particles?

In perturbation theory simple recipe to calculate discontinuities across branch cuts of multi-leg, multi-loop Feynman diagrams:

Cutkosky rule: $\mathbf{1/(p^2-m^2+i\delta)\to(-2\pi i)\delta(p^2-m^2)\theta(p_0)}$

Cutkosky rule in case of the bubble integral:

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2}, \qquad D_1 = l^2 - m^2 + i0, \quad D_2 = (p-l)^2 - m^2 + i0$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2}, \qquad D_1 = l^2 - m^2 + i0, \quad D_2 = (p-l)^2 - m^2 + i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s = p^2$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2}, \qquad D_1 = l^2 - m^2 + i0, \quad D_2 = (p-l)^2 - m^2 + i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s = p^2$

$$\epsilon_l = \sqrt{\vec{l}^2 + m^2}$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$$D_1:\qquad a_1)\ l_0=\epsilon_l-i0,\qquad b_1)\ l_0=-\epsilon_l+i0,$$

$$D_2:\qquad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$$

Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$$D_1:\qquad a_1)\ l_0=\epsilon_l-i0,\qquad\qquad b_1)\ l_0=-\epsilon_l+i0,$$

$$D_2:\qquad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$$



Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$$D_1:\qquad a_1)\ l_0=\epsilon_l-i0,\qquad\qquad b_1)\ l_0=-\epsilon_l+i0,$$

$$D_2:\qquad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$$

No discontinuity in p_0 if the poles a_2,b_2 are far from the contour



Cutkosky rule in case of the bubble integral:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2},\qquad D_1=l^2-m^2+i0,\quad D_2=(p-l)^2-m^2+i0$$

Integrate over $\mathbf{l_0}$ and calculate the discontinuity in $s=p^2$

Work in the reference frame: $p=(p_0,\vec{0})$ and $p_0>0$

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\int_C\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)}.\qquad \epsilon_l=\sqrt{\vec{l}^2+m^2}$$

The integrand has four poles in l_0 :

$D_1:\quad a_1)\ l_0=\epsilon_l-i0,\qquad b_1)\ l_0=-\epsilon_l+i0,$

$D_2:\quad a_2)\ l_0=p_0+\epsilon_l-i0,\qquad b_2)\ l_0=p_0-\epsilon_l+i0,$

No discontinuity in p_0 if the poles a_2,b_2 are far from the contour

The distance of the poles a_1,b_2 can vanish if $p_0>2m$ when these poles can pinch the contour.



One can avoid pinching the contour by moving the first pole above the contour.
To compensate the difference for the integral we add an integral over a closed small circle around the first pole

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

$$\mathcal{I}^{(2)} = \int \frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\left[\int_{C'}\frac{\mathrm{d}l_0}{(2\pi)}\ldots + \oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\ldots\right]$$

$$\oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)} = \int\frac{\mathrm{d}l_0}{(2\pi)}\frac{(-2\pi i)\delta_+(l^2-m^2)}{((l_0-p_0)^2-\epsilon_l^2)}$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \mathcal{I}_2(p^2+i\delta)-\mathcal{I}_2(p^2-i\delta) = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\times\left[\frac{1}{(l_0-p_0-i\delta)^2-\epsilon_l^2}-\mathrm{c.c}\right]$$

$$= \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\left[\frac{1}{(l-p)^2-m^2+i\delta}-\mathrm{c.c.}\right]$$

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

$$\mathcal{I}^{(2)} = \int \frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\left[\int_{C'}\frac{\mathrm{d}l_0}{(2\pi)}\ldots + \oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\ldots\right]$$

$$\oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)} = \int\frac{\mathrm{d}l_0}{(2\pi)}\frac{(-2\pi i)\delta_+(l^2-m^2)}{((l_0-p_0)^2-\epsilon_l^2)}$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \mathcal{I}_2(p^2+i\delta)-\mathcal{I}_2(p^2-i\delta) = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\times\left[\frac{1}{(l_0-p_0-i\delta)^2-\epsilon_l^2}-\mathrm{c.c}\right]$$

$$= \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\left[\frac{1}{(l-p)^2-m^2+i\delta}-\mathrm{c.c.}\right]$$

$$\frac{1}{x+i\delta} = \mathrm{P}\left[\frac{1}{x}\right] - i\pi\delta(x)$$

Discontinuity of the bubble integral = discontinuity of the pinch contribution:

$$\mathcal{I}^{(2)} = \int \frac{\mathrm{d}^3\vec{l}}{(2\pi)^3}\left[\int_{C'}\frac{\mathrm{d}l_0}{(2\pi)}\ldots + \oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\ldots\right]$$

$$\oint_{\gamma_{a_1}}\frac{\mathrm{d}l_0}{(2\pi)}\frac{1}{(l_0^2-\epsilon_l^2)((l_0-p_0)^2-\epsilon_l^2)} = \int\frac{\mathrm{d}l_0}{(2\pi)}\frac{(-2\pi i)\delta_+(l^2-m^2)}{((l_0-p_0)^2-\epsilon_l^2)}$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \mathcal{I}_2(p^2+i\delta)-\mathcal{I}_2(p^2-i\delta) = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\times\left[\frac{1}{(l_0-p_0-i\delta)^2-\epsilon_l^2}-\mathrm{c.c}\right]$$

$$= \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\left[\frac{1}{(l-p)^2-m^2+i\delta}-\mathrm{c.c.}\right]$$

$$\frac{1}{x+i\delta} = \mathrm{P}\left[\frac{1}{x}\right] - i\pi\delta(x)$$

$$\mathrm{Disc}\,[\mathcal{I}_2] = \int\frac{\mathrm{d}^4l}{(2\pi)^4}(-2\pi i)\delta_+(l^2-m^2)\;(-2\pi i)\delta_+((l-p)^2-m^2).$$

Cutkosky rule, generalizations from the case of the bubble integral:

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2} = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \int \mathrm{d}\alpha_1 \mathrm{d}\alpha_2 \frac{\delta(1 - \sum \alpha_i)}{D^2}, \qquad D = \alpha_1 D_1 + \alpha_2 D_2$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2, m^2) = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \frac{1}{D_1 D_2} = \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \int \mathrm{d}\alpha_1 \mathrm{d}\alpha_2 \frac{\delta(1 - \sum \alpha_i)}{D^2}, \qquad D = \alpha_1 D_1 + \alpha_2 D_2$$

Necessary condition for singularity: $D_1(l^*) = 0\,,\ D_2(l^*) = 0$ and so $D(l^*) = 0$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

$$\mathcal{I}(\{p_i\},\{m_i\})=\int\prod_{k=1}^{L}\frac{d^4l_k}{(2\pi)^4}\frac{\mathcal{N}(\{l_i\}\{p_i\})}{\prod_{i=1}^{N}D_i}\qquad D_i=q_i^2-m_i^2,i\in N$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

$$\mathcal{I}(\{p_i\},\{m_i\})=\int\prod_{k=1}^{L}\frac{d^4l_k}{(2\pi)^4}\,\frac{\mathcal{N}(\{l_i\}\,\{p_i\})}{\prod_{i=1}^{N}D_i}\qquad D_i=q_i^2-m_i^2,i\in N$$

$$D_i=q_i^2-m_i^2=0,\quad (i=1,2,...,r\leq 4L)$$

$$\sum_{i\subset\text{loop j}}\alpha_i^{(j)}q_i=0\quad\text{for every loop.}$$

Cutkosky rule, generalizations from the case of the bubble integral:

Landau equations:

$$\mathcal{I}^{(2)}(p^2,m^2)=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\frac{1}{D_1D_2}=\int\frac{\mathrm{d}^4l}{(2\pi)^4}\int\mathrm{d}\alpha_1\mathrm{d}\alpha_2\frac{\delta(1-\sum\alpha_i)}{D^2},\qquad D=\alpha_1D_1+\alpha_2D_2$$

Necessary condition for singularity: $D_1(l^*)=0\,,\ D_2(l^*)=0$ and so $D(l^*)=0$

$$D(l)=(l_0-a(\vec{l},p,m))(l_0-b(\vec{l},p,m)),\qquad \left.\frac{\partial D}{\partial l_0}\right|_{l_0=a\ \text{or}\ b}=0$$

Since quadratic equation, the sufficient condition also requires that two poles coincide: $\left.\frac{\partial D}{\partial l^\mu}\right|_{l=l^*}=0$

$$\mathcal{I}(\{p_i\},\{m_i\})=\int\prod_{k=1}^{L}\frac{d^4l_k}{(2\pi)^4}\frac{\mathcal{N}(\{l_i\}\{p_i\})}{\prod_{i=1}^{N}D_i}\qquad D_i=q_i^2-m_i^2, i\in N$$

$$D_i=q_i^2-m_i^2=0,\quad (i=1,2,...,r\leq 4L)$$

$$\sum_{i\subset\text{loop j}}\alpha_i^{(j)}q_i=0\quad\text{for every loop.}$$

Excercise: Work out the Landau equations and solve them for the self energy and two different internal mass and find the location of the branch cut.

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

$$\text{Disc}\,[\mathcal{I}] = \int \prod_{k=1}^{L} \frac{d^4 l_k}{(2\pi)^4} \; \frac{\mathcal{N}(\{l_i\}\,\{p_i\}) \prod_{i=1}^{r} (-2\pi i)\delta^{(+)}(q_i^2 - m_i^2)}{\prod_{j=r+1}^{N} (q_j^2 - m_j^2)}$$

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

$$\mathrm{Disc}\,[\mathcal{I}] = \int \prod_{k=1}^{L} \frac{d^4 l_k}{(2\pi)^4} \; \frac{\mathcal{N}(\{l_i\}\,\{p_i\}) \prod_{i=1}^{r} (-2\pi i)\delta^{(+)}(q_i^2 - m_i^2)}{\prod_{j=r+1}^{N} (q_j^2 - m_j^2)}$$

Unitarity: Non-linear relations between scattering amplitudes.
It can be used to compute the discontinuities of scattering amplitudes at a given order in PT in terms of amplitudes at lower order.

It is built into perturbation theory in even for external off-shell lines and external lines with on-shell complex momenta.
If the Landau equations have solutions, the Cutkosky formula provide the correct singularities.

The diagrammatic version of the Cutkosky formula also can be applied to scattering amplitudes, leading to generalized unitarity relations. E.g. triple cut.

Generalizations of the Cutkosky rule to L loop Feynman-diagrams with N denominators:

$$\text{Disc}\,[\mathcal{I}] = \int \prod_{k=1}^{L} \frac{d^4 l_k}{(2\pi)^4} \; \frac{\mathcal{N}(\{l_i\}\,\{p_i\}) \prod_{i=1}^{r} (-2\pi i)\delta^{(+)}(q_i^2 - m_i^2)}{\prod_{j=r+1}^{N} (q_j^2 - m_j^2)}$$

Unitarity: Non-linear relations between scattering amplitudes.
It can be used to compute the discontinuities of scattering amplitudes at a given order in PT in terms of amplitudes at lower order.

It is built into perturbation theory in even for external off-shell lines and external lines with on-shell complex momenta.
If the Landau equations have solutions, the Cutkosky formula provide the correct singularities.

The diagrammatic version of the Cutkosky formula also can be applied to scattering amplitudes, leading to generalized unitarity relations. E.g. triple cut.

$$\text{Disc}_{l \to l_{n_1 n_2 n_3}} \left[A_n^{1-\text{loop}}\right] = \sum_{\text{states}} \int \frac{\mathrm{d}^4 l}{(2\pi)^4} \prod_{i=1}^{3} (-2\pi i)\delta_+(l_i^2 - m_i^2)$$

$$\times A_{n_1}^{\text{tree}}(-l_3, l_1) A_{n_2}^{\text{tree}}(-l_1, l_2) A_{n_3}^{\text{tree}}(-l_2, l_3),$$

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:



Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.



Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})} + \frac{\mathbf{l}_\mu \mathbf{b}_\nu + \mathbf{b}_\mu \mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}} = \sum_{\mathbf{i=1}}^{\mathbf{D_s-2}} \mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}} \mathbf{l}_\mu \mathbf{\Gamma}^\mu + \mathbf{m} = \sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}} \mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})} + \frac{\mathbf{l}_\mu \mathbf{b}_\nu + \mathbf{b}_\mu \mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}} = \sum_{\mathbf{i=1}}^{\mathbf{D_s-2}} \mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}} \mathbf{l}_\mu \mathbf{\Gamma}^\mu + \mathbf{m} = \sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}} \mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1 n_2 n_3 n_4}}} \left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right] = \sum_{\mathrm{states}}$$

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})}+\frac{\mathbf{l}_\mu\mathbf{b}_\nu+\mathbf{b}_\mu\mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}}=\sum_{\mathbf{i=1}}^{\mathbf{D_s-2}}\mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}}\mathbf{l}_\mu\mathbf{\Gamma}^\mu+\mathbf{m}\;=\sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}}\mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1}}\mathbf{n_2}\mathbf{n_3}\mathbf{n_4}}\;\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right]=\sum_{\mathrm{states}}\;\mathcal{A}_{\mathbf{n_1}}^{\mathrm{tree}}(-\mathbf{l_4},\mathbf{l_1})\mathcal{A}_{\mathbf{n_2}}^{\mathrm{tree}}(-\mathbf{l_1},\mathbf{l_2})\mathcal{A}_{\mathbf{n_3}}^{\mathrm{tree}}(-\mathbf{l_2},\mathbf{l_3})\mathcal{A}_{\mathbf{n_4}}^{\mathrm{tree}}(-\mathbf{l_3},\mathbf{l_4}),$$

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})}+\frac{\mathbf{l}_\mu\mathbf{b}_\nu+\mathbf{b}_\mu\mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}}=\sum_{\mathbf{i=1}}^{\mathbf{D_s-2}}\mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}}\mathbf{l}_\mu\mathbf{\Gamma}^\mu+\mathbf{m}\;=\sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}}\mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1}}\mathbf{n_2}\mathbf{n_3}\mathbf{n_4}}\;\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right]=\sum_{\mathrm{states}}\;\mathcal{A}_{\mathbf{n_1}}^{\mathrm{tree}}(-\mathbf{l_4},\mathbf{l_1})\mathcal{A}_{\mathbf{n_2}}^{\mathrm{tree}}(-\mathbf{l_1},\mathbf{l_2})\mathcal{A}_{\mathbf{n_3}}^{\mathrm{tree}}(-\mathbf{l_2},\mathbf{l_3})\mathcal{A}_{\mathbf{n_4}}^{\mathrm{tree}}(-\mathbf{l_3},\mathbf{l_4}),$$

Key point: two independent expressions for the same cut amplitude

Generalized cuts of the integrands of one loop amplitudes are a given in terms of tree amplitudes:

The numerator of the propagators can be replaced with their axial-gauge values.

$$-\mathbf{g}_{\mu\nu}^{(\mathbf{D_s})} + \frac{\mathbf{l}_\mu \mathbf{b}_\nu + \mathbf{b}_\mu \mathbf{l}_\nu}{\mathbf{l}\cdot\mathbf{b}} = \sum_{\mathbf{i=1}}^{\mathbf{D_s-2}} \mathbf{e}_\mu^{(\mathbf{i})}(\mathbf{l})\mathbf{e}_\nu^{(\mathbf{i})}(\mathbf{l}),$$

$$\sum_{\mu=\mathbf{1}}^{\mathbf{D}} \mathbf{l}_\mu \mathbf{\Gamma}^\mu + \mathbf{m} = \sum_{\mathbf{i=1}}^{\mathbf{2^{(D_s-2)/2}}} \mathbf{u}^{(\mathbf{i})}(\mathbf{l})\overline{\mathbf{u}}^{(\mathbf{i})}(\mathbf{l}),$$



$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}_{\mathbf{n_1 n_2 n_3 n_4}}}\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right] = \sum_{\mathrm{states}} \mathcal{A}_{\mathbf{n_1}}^{\mathrm{tree}}(-\mathbf{l_4},\mathbf{l_1})\mathcal{A}_{\mathbf{n_2}}^{\mathrm{tree}}(-\mathbf{l_1},\mathbf{l_2})\mathcal{A}_{\mathbf{n_3}}^{\mathrm{tree}}(-\mathbf{l_2},\mathbf{l_3})\mathcal{A}_{\mathbf{n_4}}^{\mathrm{tree}}(-\mathbf{l_3},\mathbf{l_4}),$$

Key point: two independent expressions for the same cut amplitude

$$\mathrm{Res}_{\mathbf{l}\to\mathbf{l}*=\mathbf{l}_{\mathbf{n_1 n_2 n_3 n_4}}}\left[\mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}\right] = \sum_{\mathbf{n_5}} \frac{\tilde{\mathbf{e}}_{\mathbf{n_1 n_2 n_3 n_4 n_5}}(\mathbf{l}^*)}{\mathbf{d}_{\mathbf{n_5}}(\mathbf{l}^*)} + \tilde{\mathbf{d}}_{\mathbf{n_1 n_2 n_3 n_4}}(\mathbf{l}^*)$$

Projecting out individual quadrupole coefficients in D-dimension:

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^{\mathbf{4}},$$

$$\mathrm{l}^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathbf{n}_4^\mu + \sin\phi\ \mathbf{n}_\epsilon^\mu), \qquad \mathbf{V}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_\mathbf{i}^\mu \left(\mathbf{q}_\mathbf{i}^\mathbf{2} - \mathbf{m}_\mathbf{i}^\mathbf{2} + \mathbf{m}_\mathbf{0}^\mathbf{2}\right),$$

$$\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2 + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^4,$$

$$\mathrm{l}^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathbf{n}_4^\mu + \sin\phi\ \mathbf{n}_\epsilon^\mu), \qquad \mathbf{V}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_\mathbf{i}^\mu \left(\mathbf{q}_\mathbf{i}^\mathbf{2} - \mathbf{m}_\mathbf{i}^\mathbf{2} + \mathbf{m}_\mathbf{0}^\mathbf{2}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_\pm^\mu = \mathbf{V}^\mu \pm \mathrm{l}_\perp \mathbf{n}_4^\mu,$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_\mathbf{0} = \frac{\mathrm{RR}(\mathrm{l}_+) + \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_\mathbf{1} = \frac{\mathrm{RR}(\mathrm{l}_+) - \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}\mathrm{l}_\perp}$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_{\mathbf{4}}) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_{\epsilon})^{\mathbf{4}},$$

$$\mathrm{l}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}(\cos\phi\ \mathbf{n}_{\mathbf{4}}^{\mu} + \sin\phi\ \mathbf{n}_{\epsilon}^{\mu}), \qquad \mathbf{V}^{\mu} = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_{\mathbf{i}}^{\mu}\left(\mathbf{q}_{\mathbf{i}}^{\mathbf{2}} - \mathbf{m}_{\mathbf{i}}^{\mathbf{2}} + \mathbf{m}_{\mathbf{0}}^{\mathbf{2}}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_{\pm}^{\mu} = \mathbf{V}^{\mu} \pm \mathrm{l}_{\perp}\mathbf{n}_{\mathbf{4}}^{\mu},$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_{\mathbf{0}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) + \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_{\mathbf{1}} = \frac{\mathrm{RR}(\mathrm{l}_{+}) - \mathrm{RR}(\mathrm{l}_{-})}{\mathbf{2}\mathrm{l}_{\perp}}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_{\pm} = \mathbf{V} \pm \mathrm{l}_{\perp}(\mathbf{n}_{\mathbf{4}} + \mathbf{n}_{\epsilon})/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_{\mathbf{2}} + \tilde{\mathbf{d}}_{\mathbf{4}}\frac{\mathrm{l}_{\perp}^{\mathbf{2}}}{\mathbf{2}} = \mathrm{l}_{\perp}^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) + \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \mathbf{2}\,\tilde{\mathbf{d}}_{\mathbf{0}}\right), \tilde{\mathbf{d}}_{\mathbf{3}} = \sqrt{\mathbf{2}}\,\mathrm{l}_{\perp}^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_{+}) - \mathrm{RR}(\tilde{\mathrm{l}}_{-}) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_{\mathbf{1}}\mathrm{l}_{\perp}\right).$$

- Choose $\mathrm{l}_{\epsilon}^{\mu} = \mathbf{V}^{\mu} + \mathrm{l}_{\perp}$

Projecting out individual quadrupole coefficients in D-dimension:

$$\tilde{\mathbf{d}}_{\mathbf{0123}}(\mathrm{l}) = \tilde{\mathbf{d}}_{\mathbf{0}} + \tilde{\mathbf{d}}_{\mathbf{1}}(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{2}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2 + \tilde{\mathbf{d}}_{\mathbf{3}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^2(\mathrm{l}\cdot\mathbf{n}_4) + \tilde{\mathbf{d}}_{\mathbf{4}}(\mathrm{l}\cdot\mathbf{n}_\epsilon)^4,$$

$$\mathrm{l}^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp(\cos\phi\ \mathbf{n}_4^\mu + \sin\phi\ \mathbf{n}_\epsilon^\mu), \qquad \mathbf{V}^\mu = -\frac{\mathbf{1}}{\mathbf{2}}\sum_{\mathbf{i}}^{\mathbf{3}} \mathbf{v}_\mathbf{i}^\mu \left(\mathbf{q}_\mathbf{i}^\mathbf{2} - \mathbf{m}_\mathbf{i}^\mathbf{2} + \mathbf{m}_\mathbf{0}^\mathbf{2}\right),$$

- Choose $\sin\phi = \mathbf{0}, \cos\phi = \pm\mathbf{1}$ denote $\mathrm{l}_\pm^\mu = \mathbf{V}^\mu \pm \mathrm{l}_\perp \mathbf{n}_4^\mu,$

  calculate the numerator for these values

$$\tilde{\mathbf{d}}_\mathbf{0} = \frac{\mathrm{RR}(\mathrm{l}_+) + \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}}, \quad \tilde{\mathbf{d}}_\mathbf{1} = \frac{\mathrm{RR}(\mathrm{l}_+) - \mathrm{RR}(\mathrm{l}_-)}{\mathbf{2}\mathrm{l}_\perp}$$

- Choose $\cos\phi = \sin\phi = \pm\mathbf{1}/\sqrt{\mathbf{2}},$ denote $\tilde{\mathrm{l}}_\pm = \mathbf{V} \pm \mathrm{l}_\perp(\mathbf{n}_4 + \mathbf{n}_\epsilon)/\sqrt{\mathbf{2}}$

$$\tilde{\mathbf{d}}_\mathbf{2} + \tilde{\mathbf{d}}_\mathbf{4}\frac{\mathrm{l}_\perp^\mathbf{2}}{\mathbf{2}} = \mathrm{l}_\perp^{-\mathbf{2}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_+) + \mathrm{RR}(\tilde{\mathrm{l}}_-) - \mathbf{2}\,\tilde{\mathbf{d}}_\mathbf{0}\right), \tilde{\mathbf{d}}_\mathbf{3} = \sqrt{\mathbf{2}}\,\mathrm{l}_\perp^{-\mathbf{3}}\left(\mathrm{RR}(\tilde{\mathrm{l}}_+) - \mathrm{RR}(\tilde{\mathrm{l}}_-) - \sqrt{\mathbf{2}}\tilde{\mathbf{d}}_\mathbf{1}\mathrm{l}_\perp\right).$$

- Choose $\mathrm{l}_\epsilon^\mu = \mathbf{V}^\mu + \mathrm{l}_\perp$

$$\tilde{\mathbf{d}}_\mathbf{2} + \tilde{\mathbf{d}}_\mathbf{4}\mathrm{l}_\perp^\mathbf{2} = \frac{\mathrm{Num}(\mathrm{l}_\epsilon) - \tilde{\mathbf{d}}_\mathbf{0}}{\mathrm{l}_\perp^\mathbf{2}}.$$

The parameters are fixed by linear algebraic equations

$$\mathrm{Res}_{ij\cdots k}\left[F(l)\right] \equiv \left[d_i(l)d_j(l)\cdots d_k(l)F(l)\right]\Big]_{l=l_{ij\cdots k}} .$$

$$\bar{d}_{ijkl}(l) = \mathrm{Res}_{ijkl}\left(\mathcal{A}_N(l)\right) \qquad d_i=d_j=d_k=d_l=0 \qquad \text{two solutions}$$

$$\bar{c}_{ijk}(l) = \mathrm{Res}_{ijk}\left(\mathcal{A}_N(l) - \sum_{l\neq i,j,k}\frac{\bar{d}_{ijkl}(l)}{d_i d_j d_k d_l}\right) \qquad d_i=d_j=d_k=0 \qquad \text{infinite \# of solutions}$$

$$\bar{b}_{ij}(l) = \mathrm{Res}_{ij}\left(\mathcal{A}_N(l) - \sum_{k\neq i,j}\frac{\bar{c}_{ijk}(l)}{d_i d_j d_k} - \frac{1}{2!}\sum_{k,l\neq i,j}\frac{\bar{d}_{ijkl}(l)}{d_i d_j d_k d_l}\right) \qquad d_i=d_j=0$$

infinite # of solutions

unitarity: the residues factorize into the products of tree amplitudes

we fully reconstruct the integrand in terms of product of tree amplitudes

no Feynman diagrams

Unitarity method: one-loop amplitudes from tree amplitudes + scalar integral functions

- Decompose the amplitude in terms of basic set of scalar integral functions and read out the coefficients using unitarity cuts (‘98) (BDK)

- Consider the integrand, the amplitude is parametric integral over the loop momentum OPP(‘06) (EGK (’07))

- Use generalized cuts, read out the coefficients in terms of tree amplitudes at cut-momenta (complex) BCF/BDK(’05)

- Rational part is obtained by carrying out the algorithm in two different integer D>4 dimensions GKM (08) (see also Badger,BDK, OPP)

Unitarity method: one-loop amplitudes from tree amplitudes + scalar integral functions

- Decompose the amplitude in terms of basic set of scalar integral functions and read out the coefficients using unitarity cuts (‘98) (BDK)

- Consider the integrand, the amplitude is parametric integral over the loop momentum OPP(‘06) (EGK (’07))

- Use generalized cuts, read out the coefficients in terms of tree amplitudes at cut-momenta (complex) BCF/BDK(’05)

- Rational part is obtained by carrying out the algorithm in two different integer D>4 dimensions GKM (08) (see also Badger,BDK, OPP)

Bern, Dixon, Kosower (BDK) ; Ellis, Giele, ZK, Melnikov (EGKM); Britto, Cachazo, Feng (BCF)

Managing color of amplitudes

Unitarity: instead of Feynman diagrams amplitudes
Tree and one loop Feynman diagrams: color and space time part

$$\mathcal{A}_{\mathbf{n}}^{\mathrm{tree}}(\{\mathbf{k}\},\{\mathbf{h}\},\{\mathbf{a}\}),\ \mathcal{A}_{\mathbf{n}}^{1-\mathrm{loop}}(\{\mathbf{k}\},\{\mathbf{h}\},\{\mathbf{a}\})$$

➡ Color decomposition of amplitudes with the help of a basis color space
(T-based, F-based, color flow based}

$$[T^a,T^b]=-F^a_{bc}T^c\,,\qquad \mathrm{Tr}(T^aT^b)=\delta_{ab}.$$

$$[F^a,F^b]=-F^a_{bc}F^c\,,\quad F^a_{bc}=-i\sqrt{2}f^{abc}\,,\quad \mathrm{Tr}(F^aF^b)=2N_c\delta_{ab}.$$
$$\left(F^{a_1}\right)_{a_2a_3}=-\frac{1}{2N_c}\mathrm{Tr}\left([F^{a_1},F^{a_2}]F^{a_3}\right)=-\mathrm{Tr}\left([T^{a_1},T^{a_2}]T^{a_3}\right).$$

$$(T^a)^{\bar{j}_1}_{i_1}(T^a)^{\bar{j}_2}_{i_2}=(\delta)^{\bar{j}_2}_{i_1}(\delta)^{\bar{j}_1}_{i_2}-\frac{1}{N_c}(\delta)^{\bar{j}_1}_{i_1}(\delta)^{\bar{j}_2}_{i_2}$$

$$\begin{aligned}\left(F^{a_2}F^{a_3}\ldots F^{a_{(n-2)}}F^{a_{(n-1)}}\right)_{a_1a_n}&=\frac{1}{2N_c}\mathrm{Tr}\left([[\ldots[[F^{a_1},F^{a_2}],F^{a_3}],\ldots,F^{a_{n-2}}]\,[F^{a_{n-1}},F^{a_n}]\right)\\&=\mathrm{Tr}\left([[\ldots[[T^{a_1},T^{a_2}],T^{a_3}],\ldots,T^{a_{n-2}}]\,[T^{a_{n-1}},T^{a_n}]\right).\end{aligned}$$

Color ordered n-gluon tree sub-amplitudes

$$\mathcal{A}_{\mathbf{n}}^{\text{tree}} = \frac{\mathbf{g_s^{n-2}}}{\mathbf{2N_c}} \sum_{\sigma \in \mathbf{S_n/Z_n}} \mathrm{Tr}\left(\mathbf{F^{a_{\sigma(1)}} F^{a_{\sigma(2)}} F^{a_{\sigma(3)}} \dots F^{a_{\sigma(n)}}}\right) \mathbf{A}_{\mathbf{n},\sigma}^{\text{tree}},$$

(n-1)! color ordered sub-amplitudes

$$\mathcal{A}_{\mathbf{n}}^{\text{tree}} = \mathbf{g_s^{n-2}} \sum_{\sigma \in \mathbf{S_n/Z_n}} \mathrm{Tr}\left(\mathbf{T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} T^{a_{\sigma(3)}} \dots T^{a_{\sigma(n)}}}\right) \mathbf{A}_{\mathbf{n},\sigma}^{\text{tree}}$$

$$\mathbf{A}_{\mathbf{n},\sigma}^{\text{tree}} = \mathbf{m_n}\left(\mathbf{g_{\sigma(1)}, g_{\sigma(2)}, g_{\sigma(3)}, \dots, g_{\sigma(n)}}\right)$$

Some properties of sub-amplitudes:

$$\mathbf{m_n}(\mathbf{g_1, g_2, g_3, \dots, g_n}) = \mathbf{m_n}(\mathbf{g_2, g_3, \dots, g_n, g_1})$$ (cyclic identity)

$$\mathbf{m_n}(\mathbf{g_1, g_2, g_3, \dots, g_{n-1}, g_n}) = (-1)^{\mathbf{n}} \mathbf{m_n}(\mathbf{g_n, g_{n-1}, \dots, g_2, g_1}),$$ (reflection identity)

$$\mathbf{m_n}(\mathbf{1}, \underline{\mathbf{2}, \dots \mathbf{n_1}}, \overline{\mathbf{n_1+1}, \dots, \mathbf{n}}) \equiv \sum_{\sigma(\mathbf{n})} \mathbf{m_n}(\mathbf{g_1, g_{\sigma(2)}, g_{\sigma(3)}, \dots, g_{\sigma(n)}}) = \mathbf{0}$$ (Abelian identity)

~ Kleiss-Kuijf relations

$$\mathcal{A}_{\mathbf{n}}^{\text{tree}}(\mathbf{1,2,3,\dots,n}) = \mathbf{g_s^{n-2}} \sum_{\sigma = \mathcal{P}(\mathbf{2,3,\dots,n-1})} \left(\mathbf{F^{a_{\sigma(2)}} \dots F^{a_{\sigma(n-1)}}}\right)_{\mathbf{a_1 a_n}} \mathbf{m_n}(\mathbf{g_1, g_{\sigma(2)}, g_{\sigma(3)}, \dots, g_{\sigma(n-1)}, g_n}).$$

(n-2)! color ordered sub-amplitudes (see also BCJ relations)

Unitary color basis: on each pole of the tree amplitude the color factor of a given colorless amplitude also factorizes

three gluon vertex



$$i\frac{g}{\sqrt{2}}\left((k_1-k_2)_{\mu_3}g_{\mu_1\mu_2}+(k_2-k_3)_{\mu_1}g_{\mu_2\mu_3}+(k_3-k_1)_{\mu_2}g_{\mu_3\mu_1}\right.$$

four gluon vertex



$$i\frac{g^2}{2}(2g_{\mu_1\mu_3}g_{\mu_2\mu_4}-g_{\mu_1\mu_4}g_{\mu_2\mu_3}-g_{\mu_1\mu_2}g_{\mu_3\mu_4})$$

gluon-quark-antiquark vertex



sign depends on orientation

Color stripped Feynman rules for color ordered sub-amplitudes

three gluon vertex



$$i\frac{g}{\sqrt{2}}\left((k_1-k_2)_{\mu_3}g_{\mu_1\mu_2}+(k_2-k_3)_{\mu_1}g_{\mu_2\mu_3}+(k_3-k_1)_{\mu_2}g_{\mu_3\mu_1}\right)$$

four gluon vertex



$$i\frac{g^2}{2}(2g_{\mu_1\mu_3}g_{\mu_2\mu_4}-g_{\mu_1\mu_4}g_{\mu_2\mu_3}-g_{\mu_1\mu_2}g_{\mu_3\mu_4})$$

gluon-quark-antiquark vertex



sign depends on orientation

Color ordered n-gluon one-loop sub-amplitudes

$$\mathcal{A}_{\mathbf{n}}^{1-\text{loop}} = \mathbf{g}_{\mathbf{s}}^{\mathbf{n}} \mathbf{c}_{\mathbf{\Gamma}} \sum_{\sigma \in \mathbf{S}_{\mathbf{n-1}}} \mathbf{A}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g}_{\sigma(\mathbf{2})}, \ldots, \mathbf{g}_{\sigma(\mathbf{n})}),$$

$$\mathbf{c}_{\mathbf{\Gamma}} = \frac{\mathbf{1}}{(\mathbf{4\pi})^{\mathbf{2-\epsilon}}} \frac{\mathbf{\Gamma(1+\epsilon)\Gamma^2(1-\epsilon)}}{\mathbf{\Gamma(1-2\epsilon)}}$$

Consider a double cut between external line k and k+1 and n and 1

$$\begin{aligned}\text{Im}_{(\mathbf{k},\mathbf{n})}\left[\mathbf{A}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_n})\right] &= \{\mathbf{a_1 a_2 \cdots a_k}\}_{\mathbf{vu}} \{\mathbf{a_{k+1} \cdots a_n}\}_{\mathbf{uv}} \\ &\quad \times \mathbf{m_{k+2}}(\mathbf{g_v}, \mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_k}, \mathbf{g_u}) \mathbf{m_{n-k+2}}(\mathbf{g_u}, \mathbf{g_{k+1}}, \ldots, \mathbf{g_n}, \mathbf{g_v}) \\ &= \{\mathbf{a_1 a_2 \cdots a_n}\} \, \text{Im}_{(\mathbf{k},\mathbf{n})}\left[\mathbf{m}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_n})\right]\end{aligned}$$

$$\mathcal{A}_{\mathbf{n}}^{1-\text{loop}} = \mathbf{g}_{\mathbf{s}}^{\mathbf{n}} \mathbf{c}_{\mathbf{\Gamma}} \sum_{\mathcal{P}(\mathbf{2},\cdots,\mathbf{n})/\mathcal{R}} \text{Tr}(\mathbf{F}^{\mathbf{a_1}}, \ldots, \mathbf{F}^{\mathbf{a_n}}) \mathbf{m}_{\mathbf{n}}^{(\mathbf{1})}(\mathbf{g_1}, \mathbf{g_2}, \ldots, \mathbf{g_n})$$

one loop color order sub-amplitude

A reflection transformation is factored out. The cyclic property and reflection symmetry remain valid. The number of independent one-loop amplitudes is (n-1)!/2.

Color ordered n-gluon one-loop sub-amplitudes for large N_c

Decomposition in T-basis:

$$\mathcal{A}_{n}^{1-\text{loop}} = g_s^n N_c c_\Gamma \sum_{\mathcal{P}(2,\cdots,n)/\mathcal{R}} \text{Tr}(T^{a_1},\ldots,T^{a_n}) m_{1,n}^{(1)}(g_1,g_2,\ldots,g_n)$$

$$+g_s^n c_\Gamma \sum_{r=2}^{[n/2]+1} \left(\sum_{\mathcal{P}(2,\cdots,n)/\mathcal{Z}_{r-1}\times\mathcal{Z}_{n-r+1}} \text{Tr}\left(T^{a_1},\ldots,T^{a_{r-1}}\right) \text{Tr}\left(T^{a_r},\ldots,T^{a_n}\right) m_{2,n}^{(1)}\left(g_1,g_2,\ldots,g_n\right) \right)$$

- The single-trace color structures have an explicit factor of Nc out.
- They dominate in the large Nc limit.
- The planar L-loop color decomposition formula remains the same .
- The decomposition remains the same also for the N=4sYM theory.
- T-based color decomposition is preferred.

Two particle unitarity gives color decomposition of a quark-loop to n-gluon amplitudes

$$\mathcal{A}_{n;n_f}^{1-\text{loop}} = g_s^n c_\Gamma n_f \sum_{\sigma\in S_{n-1}} \text{Tr}(T^{a_1} T^{a_{\sigma(2)}} \ldots T^{a_{\sigma n}})\, m_{n;n_f}^{(1)}\left(g_1, g_{\sigma(2)},\ldots,g_{\sigma(n)}\right),$$

$\bar{q}q + (n-2)g$ amplitudes and fully ordered primitive amplitudes

Color factors of Feynman diagrams $\quad \left(\mathbf{T}^{\mathbf{b_1}} \dots \mathbf{T}^{\mathbf{b_k}} \dots\right)_{\mathbf{j\bar{\imath}}} \times \left(\mathbf{F}^{\mathbf{a_1}} \dots \mathbf{F}^{\mathbf{a_r}}\right)_{\mathbf{b_1 a_{r+1}}} \dots \left(\mathbf{F}^{\mathbf{a_p}} \dots \mathbf{F}^{\mathbf{a_t - 1}}\right)_{\mathbf{b_k a_t}} \dots$

$$\mathcal{A}_{\mathbf{n}}^{\mathrm{tree}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \mathbf{g_3}, \dots, \mathbf{g_n}) = \mathbf{g_s^{n-2}} \sum_{\sigma \in \mathbf{S_{n-2}}} \left(\mathbf{T}^{\mathbf{a_{\sigma(3)}}} \mathbf{T}^{\mathbf{a_{\sigma(4)}}} \dots \mathbf{T}^{\mathbf{a_{\sigma(n)}}}\right)_{\mathbf{i_2 \bar{\imath}_1}} \mathbf{m_n}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \mathbf{g_{\sigma(3)}}, \dots, \mathbf{g_{\sigma(n)}}).$$

(n-2)! colorless color ordered tree sub-amplitudes

the quark labels do not participate in the permutation sum

Decomposition in mixed basis such that the quark is also in the permutation sum

$$\mathcal{A}_{\mathbf{n}}^{\mathrm{tree}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \mathbf{g_3}, \dots, \mathbf{g_n}) = \mathbf{g_s^{n-2}} (-\mathbf{1})^{\mathbf{n}} \sum_{\mathbf{k=3}}^{\mathbf{n}} \sum_{\mathcal{P}(\mathbf{4},\dots,\mathbf{n})} \left(\mathbf{T^y}\, \mathbf{T}^{\mathbf{a_{k+1}}} .. \mathbf{T}^{\mathbf{a_n}}\right)_{\mathbf{i_2 \bar{\imath}_1}} \mathrm{Tr}\left(\mathbf{F}^{\mathbf{a_4}} \dots \mathbf{F}^{\mathbf{a_k}}\right)_{\mathbf{a_3 y}}$$
$$\times \tilde{\mathbf{m}}_{\mathbf{n}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{g_n}, \dots, \mathbf{g_{k+1}}, \mathbf{q_2}, \mathbf{g_k}, \dots, \mathbf{g_3})$$

Color ordered tree “left primitive amplitudes”

$$\tilde{\mathbf{m}}_{\mathbf{n}}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{g_n}, .., \mathbf{g_{(k+1)}}, \mathbf{q_2}, \mathbf{g_k}, .., \mathbf{g_3}) = (-\mathbf{1})^{\mathbf{k}} \mathbf{m_n}(\bar{\mathbf{q}}_{\mathbf{1}}, \mathbf{q_2}, \underline{\mathbf{k}, ..., \mathbf{3}}, \overline{(\mathbf{k+1}), .., \mathbf{n}})$$

Excercise: derive this relation using commutator identities

When anti-quark is in the permutation sum: “right primitive amplitude”

This mixed basis is “unitary”

$\bar{\mathbf{q}}\mathbf{q}+(\mathbf{n}-\mathbf{2})\mathbf{g}$ colorless one-loop fully ordered left primitive amplitude



$$\mathcal{A}_{\mathbf{n}}^{1-\text{loop}}(\bar{\mathbf{q}}_{\mathbf{1}},\mathbf{q}_{\mathbf{2}},\mathbf{g}_{\mathbf{3}},..,\mathbf{g}_{\mathbf{n}})=\mathbf{g}_{\mathbf{s}}^{\mathbf{n}}\mathbf{c}_{\Gamma}\sum_{\mathbf{p=2}}^{\mathbf{n}}\sum_{\sigma\in\mathbf{S}_{\mathbf{n-2}}}(\mathbf{T}^{\mathbf{x_2}}\mathbf{T}^{\mathbf{a}_{\sigma_3}}..\mathbf{T}^{\mathbf{a}_{\sigma_p}}\mathbf{T}^{\mathbf{x_1}})_{\mathbf{i_2\bar{\imath}_1}}(\mathbf{F}^{\mathbf{a}_{\sigma_{p+1}}}..\mathbf{F}^{\mathbf{a}_{\sigma_n}})_{\mathbf{x_1x_2}}$$
$$\times(-\mathbf{1})^{\mathbf{n}}\tilde{\mathbf{m}}_{\mathbf{n}}^{(\mathbf{1})}(\bar{\mathbf{q}}_{\mathbf{1}},\mathbf{g}_{\sigma(\mathbf{p})},..,\mathbf{g}_{\sigma(\mathbf{3})},\mathbf{q}_{\mathbf{2}},\mathbf{g}_{\sigma(\mathbf{n})},..,\mathbf{g}_{\sigma(\mathbf{p+1})})$$

Color ordered amplitudes for n-gluons and primitive amplitudes
for $\bar{\mathbf{q}}\mathbf{q}+(\mathbf{n}-\mathbf{2})\mathbf{g}$ can be calculated using colorless Feynman rules
Berends-Giele recursion relations

Comment: Leading color is good approximations for gluons only

Amplitudes with multiple quarks

Colorless ordered amplitudes, primitive amplitudes, parent diagrams

$$\mathcal{B}^{\rm tree}(\bar{\mathbf{q}}_1,\mathbf{q}_2,\bar{\mathbf{Q}}_3,\mathbf{Q}_4,\mathbf{g}_5)=\mathbf{g}_s^3\Big[(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_1}\delta_{\mathbf{i}_2\bar{\imath}_3}\mathbf{B}^{\rm tree}_{\mathbf{5};1}+\frac{\mathbf{1}}{\mathbf{N_c}}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_1}\delta_{\mathbf{i}_4\bar{\imath}_3}\mathbf{B}^{\rm tree}_{\mathbf{5};2}$$
$$+(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_3}\delta_{\mathbf{i}_4\bar{\imath}_1}\mathbf{B}^{\rm tree}_{\mathbf{5};3}+\frac{\mathbf{1}}{\mathbf{N_c}}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_3}\delta_{\mathbf{i}_2\bar{\imath}_1}\mathbf{B}^{\rm tree}_{\mathbf{5};4}\Big],$$

$$\mathcal{B}^{1-\rm loop}(\bar{\mathbf{q}}_1,\mathbf{q}_2,\bar{\mathbf{Q}}_3,\mathbf{Q}_4,\mathbf{g}_5)=\mathbf{g}_s^5\Big[\mathbf{N_c}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_1}\delta_{\mathbf{i}_2\bar{\imath}_3}\mathbf{B}_{\mathbf{5};1}+(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_1}\delta_{\mathbf{i}_4\bar{\imath}_3}\mathbf{B}_{\mathbf{5};2}+\mathbf{N_c}(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_2\bar{\imath}_3}\delta_{\mathbf{i}_4\bar{\imath}_1}\mathbf{B}_{\mathbf{5};3}$$
$$+(\mathbf{T}^{\mathbf{a}_5})_{\mathbf{i}_4\bar{\imath}_3}\delta_{\mathbf{i}_2\bar{\imath}_1}\mathbf{B}_{\mathbf{5};4}\Big]$$

$$\mathbf{B}_{\mathbf{5};\mathbf{i}}=\mathbf{B}^{[\mathbf{1}]}_{\mathbf{5};\mathbf{i}}+\frac{\mathbf{n_f}}{\mathbf{N_c}}\mathbf{B}^{[\mathbf{1/2}]}_{\mathbf{5};\mathbf{i}},\qquad \mathbf{i}=\mathbf{1},\mathbf{2},\mathbf{3},\mathbf{4},$$

$$\mathbf{B}^{[1/2]}_{5;1}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{g}_5,\mathbf{Q}_4,\bar{\mathbf{Q}}_3,\mathbf{q}_2),\quad \mathbf{B}^{[1/2]}_{5;2}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{Q}_4,\bar{\mathbf{Q}}_3,\mathbf{q}_2,\mathbf{g}_5),$$
$$\mathbf{B}^{[1/2]}_{5;3}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{Q}_4,\bar{\mathbf{Q}}_3,\mathbf{g}_5,\mathbf{q}_2),\quad \mathbf{B}^{[1/2]}_{5;4}=-\mathbf{A}^{[1/2]}_{\mathbf{L}}(\bar{\mathbf{q}}_1,\mathbf{Q}_4,\mathbf{g}_5,\bar{\mathbf{Q}}_3,\mathbf{q}_2).$$



Amplitudes with multiple quarks

Colorless ordered amplitudes, primitive amplitudes, parent diagrams

More and more complicated.....

End of color management

Singular behavior of one loop primitive amplitudes

n-gluon:

$$\mathbf{m_n^{(1)}(g_1,g_2,\ldots g_n)} = -\sum_{\mathbf{i=1}}^{\mathbf{n}} \left[\frac{\mathbf{1}}{\epsilon^{\mathbf{2}}} + \frac{\mathbf{1}}{\epsilon}\left(\frac{\mathbf{11}}{\mathbf{3n}} + \mathbf{L_{i,i+1}}\right)\right] \mathbf{m_n^{(0)}(g_1,g_2,\ldots g_n)},$$

$$\mathbf{L_{kn}} = \ln\left(\mu^{\mathbf{2}}/(-\mathbf{s_{kn}} - \mathbf{i0})\right)$$

$\mathbf{\bar{q}q + (n-2)g}$:

$$\mathbf{\tilde{m}_n^{(1)}(\bar{q}_n, g_{k+1}, \ldots, g_{n-1}, q_2, g_3, \ldots g_k)} =$$

$$\mathbf{\tilde{m}_n(\bar{q}_n, g_{k+1}, \ldots, g_{n-1}, q_2, g_3, \ldots g_k)} \left[-\frac{\mathbf{k}}{\epsilon^{\mathbf{2}}} - \frac{\mathbf{1}}{\epsilon}\left(\frac{\mathbf{3}}{\mathbf{2}} + \sum_{\mathbf{i=1}}^{\mathbf{k-1}} \mathbf{L_{i\,i+1}} + \mathbf{L_{kn}}\right)\right]$$

Color is eliminated. Important for testing the calculations.
Primitive tree amplitude is calculated with Berends-Giele recursion relations based on color stripped Feynman rules or BCFW recursion relations